

Solomon Press
Statistics S1
Paper I
(Mark Scheme)

**All exam papers are issued free to students for education purpose only.
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations
Advanced Subsidiary / Advanced Level

Statistics
Module S1

Paper I

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong & Chris Huffer

© *Solomon Press*



These sheets may be copied for use solely by the purchaser's institute.

S1 Paper I – Marking Guide


1. (a) (i) normal A1
 (ii) e.g. producer must ensure that most bottles contain at least 75 cl B1
- (b) (i) discrete uniform A1
 (ii)
- | | | | | |
|----------|---------------|---------------|---------------|---------------|
| x | 1 | 2 | 3 | 4 |
| $P(X=x)$ | $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ | $\frac{1}{4}$ |
- mean = $\frac{5}{2}$ (symmetry) A1
 $E(X^2) = \sum x^2 P(x) = \frac{1}{4} + 1 + \frac{9}{4} + 4 = \frac{15}{2}$ M1 A1
 $\text{Var}(X) = \frac{15}{2} - (\frac{5}{2})^2 = \frac{5}{4}$ M1 A1 **(8)**
-

2. (a) $P(A \cap B) = P(A) \times P(B) = 2P(B) \times P(B) = 2[P(B)]^2$ M2
 $\therefore 2[P(B)]^2 = \frac{1}{8}; \therefore [P(B)]^2 = \frac{1}{16}; \therefore P(B) = \frac{1}{4}$ M2 A1
- (b) $P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{1}{2} + \frac{1}{4} - \frac{1}{8} = \frac{5}{8}$ M2 A1
- (c) A and B independent $\therefore A$ and B' independent
 $\therefore P(A|B') = P(A) = \frac{1}{2}$ M1 A1 **(10)**
-

3. (a) $12.7 + 5.8 = 18.5$ minutes A1
- (b) $P(X < 12.7) = 0.25; P(Z < \frac{12.7 - \mu}{\sigma}) = 0.25$ M1
 $\frac{12.7 - \mu}{\sigma} = -0.67; 12.7 - \mu = -0.67\sigma$ M1 A1
 $P(X < 18.5) = 0.75; P(Z < \frac{18.5 - \mu}{\sigma}) = 0.75$ M1
 $\frac{18.5 - \mu}{\sigma} = 0.67; 18.5 - \mu = 0.67\sigma$ M1 A1
 solve simul. giving $\mu = 15.6, \sigma = 4.3284$; so $\mu = 15.6, \sigma^2 = 18.7$ M1 A1
- (c) e.g. would expect normal dist. and mean and variance seem close to actual values so seems a fairly suitable model B2 **(11)**
-

4. (a) median = $15.5^{\text{th}} = \frac{31+32}{2} = 31.5$ M1 A1
 $Q_1 = 7.75^{\text{th}} = 20$ A1
 $Q_3 = 23.25^{\text{th}} = 45.5$ A1
- (b)  B3
- (c)  B3
- (d) e.g. similar range, youngest and oldest both a bit higher for E
 median of M lower meaning younger students on average
 IQR of M smaller meaning student ages more similar
 E roughly symmetrical, M +vely skewed B4 **(14)**
-

5. (a) $(0.6 \times 0.5 \times 0.7) + (0.6 \times 0.5 \times 0.3) + (0.4 \times 0.5 \times 0.3) = 0.36$ M3 A1
- (b) $P(W = 0) = 0.4 \times 0.5 \times 0.7 = 0.14$
 $P(W = 3) = 0.6 \times 0.5 \times 0.3 = 0.09$
 $P(W = 1) = 1 - (0.14 + 0.36 + 0.09) = 0.41$
- | | | | | |
|------------|------|------|------|------|
| w | 0 | 1 | 2 | 3 |
| $P(W = w)$ | 0.14 | 0.41 | 0.36 | 0.09 |
- (c) $E(W) = \sum wP(w) = 0 + 0.41 + 0.72 + 0.27 = 1.4$ M1 A1
 $E(W^2) = \sum w^2P(w) = 0 + 0.41 + 1.44 + 0.81 = 2.66$ M1 A1
 $\text{Var}(W) = 2.66 - 1.4^2 = 0.7$ M1 A1
- (d) e.g. unlikely to be valid as result of each match will probably raise or lower confidence changing probability of success in the next match B2 (16)



- (b) $S_{TT} = 137 - \frac{7^2}{10} = 132.1$ M1
 $S_{AA} = 2172.66 - \frac{143.8^2}{10} = 104.816$ M1
 $S_{TA} = 20.7 - \frac{7 \times 143.8}{10} = -79.96$ M1
 $r = \frac{-79.96}{\sqrt{132.1 \times 104.816}} = -0.6795$ M1 A1
 e.g. fairly strong -ve correlation so belief seems reasonable B1
- (c) $q = \frac{-79.96}{132.1} = -0.60530$ M1 A1
 $p = \frac{143.8}{10} - (-0.60530 \times \frac{7}{10}) = 14.804$ M1 A1
 $A = 14.8 - 0.605T$
- (d) line on graph above B2 (16)

Total (75)