



*Research Article*

## Microwave Assisted Synthesis, Characterization and Biological studies on 1,3,4-Thiadiazinan-5-one derivatives

S. Srinivasan, R. Girija\*

Department of Chemistry, Queen Mary's College (Autonomous), Chennai, India.

\*Corresponding author's e-mail: [ss2201phd@gmail.com](mailto:ss2201phd@gmail.com)

### Abstract

Heterocyclic compounds play an important role as the main sources of lead molecules of agrochemicals. The 1,3,4-thiadiazinan-5-one nucleus is one of the most important and well known heterocyclic nuclei, which is a common and integral feature of a variety of natural products and medicinal agents. To find the novel lead compounds with various biological activities a series of thiadiazinan-5-one were rationally designed and synthesized. In this study an effort is made to synthesize 1,3,4-Thiadiazinan-5-one derivatives through the condensation reaction of Aryl aldehydes, substituted hydrazine derivatives and thioacetic acid using green solvent which is an efficient, green and facile three-component reaction. Using this method, 80% yield of thiadiazinan-5-one was obtained, while in conventional method, the yield only exceeded upto 50%. The chemical structures of the synthesized compounds were characterized by considering the data of their elemental analysis as well as their spectral data using, FT-IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR and Mass spectra. The compounds are found to possess good biological activity. Especially, compounds were bearing the sulfur and nitrogen atoms showed activity than those bearing the oxygen atom.

**Keywords:** Microwave; Aryl aldehyde; Thioacetic acid; Hydrazine; Antimicrobial activity.

### Introduction

In recent years green chemistry, which is an environmental friendly processes, has become an alternative to traditional synthetic approach which plays an important role in organic synthesis. Due to environmental awareness, the development of environmentally benign organic reactions has become a crucial and demanding research area in modern organic chemical research [1].

Microwave irradiation can be used as a facile method for the construction of a wide variety of thiadiazinone derivatives. Multi-component reactions (MCRs) are a promising and vital field of chemistry because the synthesis of complicated molecules can be achieved in a very fast, efficient and time saving manner without the isolation of any intermediate. These reactions offer a wide range of possibilities for the efficient construction of highly complex molecules in a single step. In recent years, the discovery of MCRs has become an increasingly active area of research yielding novel chemical scaffolds for drug discovery. Thus the development of new multi-component reaction is

a popular area of research in current organic chemistry [2-4]. 1,2,4-triazoles fused with 1,3,4-thiadiazines are found to possess diverse applications in the field of medicine [5,6]. Triazolo-thiadiazines are reported to show a broad spectrum of pharmacologically important properties like antifungal [7], antibacterial [8], antiviral [9], anthelmintic [10], antitumor [11], anti-inflammatory [12], antitubercular [13], diuretics [14], anticancer [15] and hypoglycemic agents [16]. 1,3,4-Thiadiazines constitute an important class of heterocycles, which has attracted much interest because of their wide range of biological activities. Considerable attention has been focused on these compounds because many of these derivatives are important matrix metalloproteinase inhibitors. They show excellent cardio tonic and hypertensive activities. These can be used for treatment of tumors and acquired immune deficiency syndrome (AIDS) [17].

Thiadiazepines are reported for their potent anti-microbial, anti-fungal activity and metalloproteinase inhibition. The literature survey has shown that some fused thiadiazepines exhibit antidepressant, central nervous

depressant, bactericidal, fungicidal, anti-cancer activity. Recently 1,4,5-dibenzolo thiadiazepine has been reported to show good neuroprotective properties against neurodegenerative diseases without anticholinergic effects [18].

In recent years, interest in thiadiazines has increased due to the high biological activity and broad spectrum action of their derivatives [19-20]. Many thiadiazines have been discovered with possible applications in medical practice as sedatives, antianxiety agents, antiasthmatic agents, anticonvulsants, myorelaxants, coronary vasodilators, and spasmolytics. Synthesis of the 1,3,4-thiadiazine system employing a reaction of  $\alpha$ -bromoacetophenone with thiosemicarbazide was first reported by Bose [21]. In continuation of the synthesis of novel heterocyclic compounds and studying their activities, it is interesting to synthesize compounds containing thiadiazine groups in their structure.

### Materials and methods

All the chemicals and the reagents used in the study were synthetic grade purity. Benzaldehyde, phenyl hydrazine, hydrazine hydrate, thioacetic acid and various related chemicals were purchased from Scientific Chemicals Pvt. Ltd.

Solvents used were purified by distillation. All substance prepared for studies were purified by crystallization using appropriate solvents and established procedures. Melting points were determined in open capillary tubes and are uncorrected. The purity of the compounds was confirmed and checked by Thin layer chromatography (TLC). All the reactions were monitored using ether:chloroform (4:1 v/v) as a solvent system. The spots were visualized by using iodine vapour. Full form (IR) spectra were recorded on FTIR-8300 Shimadzu spectrometer. Full form (NMR) spectra were recorded on Jeol-FXQ (90 MHz), Jeol GSX (400 MHz) and DPX 200 (200 MHz). Mass spectra were recorded on Jeol- JMS- DX 303hf.

### General method for the synthesis of substituted 1,3,4-thiadiazinan-5-one derivatives

Equal molar of substituted Aryl aldehydes, substituted hydrazine derivatives and thioacetic acid (1:1:1) is treated on microwave irradiation for 2-3 min afforded the formation of the red orange solid mass of thiadiazinan-5-one derivative of an excellent yield (85-92%). Various substituted thiadiazinan-5-one were prepared and are reported in Fig. 1.




Fig. 1(a). Synthesis of 2,4-substituted 1,3,4-thiadiazinan-5-one




Fig. 1(b). Synthesis of 2-substituted 1,3,4-thiadiazinan-5-one




Fig. 1(c). Synthesis of 2,4-substituted 1,3,4-thiadiazinan-5-one

Table 1. Physical data of substituted 1,3,4-thiadiazinan-5-one

Compound	R	R'	Yield %	M.P <sup>o</sup> C
1a	H	H	84	162
1b	NO <sub>2</sub>	H	78	158
1c	H	Cl	80	153
1d	H	OH	82	150
1e	OMe	OH	69	140
2a	H	H	88	159
2b	NO <sub>2</sub>	H	76	148
2c	H	OH	82	160
2d	OMe	OH	86	144
3a	H	H	73	156
3b	H	OH	79	165
3c	H	Cl	77	156
3d	NO <sub>2</sub>	H	83	162
3e	H	Cl	88	158
3f	OMe	OH	72	148
4a	H	H	75	154
4b	H	OH	78	163
4c	OMe	OH	79	168
4d	H	NO <sub>2</sub>	87	153
4e	H	Cl	81	146

## Results and discussion

Thiadiazinan-5-one plays a major role in drug industry because of its applications continued in the effort to developed new thiadiazinan-5-one derivatives. Pharmaceutical industries are facing constraints regarding environmental and energy saving aspects. In this study, we used an excellent synthetic method for the synthesis of thiadiazinan-5-one derivatives through microwave technique. Synthesis of the compounds was accomplished through only one step, aryl aldehydes with substituted hydrazine and thioacetic acid in equimolar quantities in the presence of green solvent were treated with microwave irradiation method.

Generally, the reaction that employed aromatic aldehydes bearing electron-withdrawing or electron donating functional groups at different positions produced the corresponding products thiadiazinan-5-one in good to excellent yields. Further research and applications of the reactions are in progress in our laboratories. It is believed that this method is highly useful for the synthesis of biologically potent highly substituted thiadiazinan-5-one derivatives (Table 1). Syntheses of the compounds were carried out under mild reaction condition and the products were obtained in high yields. The resulted compounds were monitored by TLC. The structures of the newly synthesized compounds were deduced from the analysis of their <sup>1</sup>H and <sup>13</sup>C NMR, IR spectra, and confirmed by elemental analysis.

### 2,4-diphenyl-1,3,4-thiadiazinan-5-one 1a

Benzaldehyde (1 g, 0.0094 m), phenyl hydrazine (1.4 g, 0.0094 m) with Thioacetic acid (1:1:1) (0.8 g, 0.0094 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.17 (s, 1H, NH), 2.66-2.80 (s, 2H, CH<sub>2</sub>), 5.20 (s, 1H, METHINE), 7.12-7.64 (m, Ar-H). <sup>13</sup>C-NMR: 35.80, 76.40, 127.20, 128.48, 129.52, 130.17, 130.50, 134.9, 170.65. IR: 3250, 3085, 1724, 1567, 812 cm<sup>-1</sup>. Mass: M/Z 268.

### 2-(3-nitrophenyl)-4-phenyl-1,3,4-thiadiazinan-5-one 1b

A mixture of 3-nitrobenzaldehyde (1 g, 0.0066 m), phenyl hydrazine (1.03 g, 0.0066 m) with thioacetic acid (0.6 g, 0.0066 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives.

<sup>1</sup>H-NMR: 2.18 (s, 1H, NH), 2.38-2.39 (s, 2H, CH<sub>2</sub>), 4.90 (s, 1H, METHINE), 7.30-7.94 (m, Ar-H). <sup>13</sup>C-NMR: 21.19, 30.46, 76.74, 112.97, 113.21, 119.17, 120.64, 122.52, 129.04, 130.18, 133.87, 135.52, 168.80. IR: 3324, 3066, 1652, 1550, 1527, 750 cm<sup>-1</sup>. Mass: M/Z 314.

#### **2-(4-chlorophenyl)-4-phenyl-1,3,4-thiadiazinan-5-one 1c**

The microwave irradiation of equal molar 4-chlorobenzaldehyde (1 g, 0.0070 m), phenyl hydrazine (1.1 g, 0.0070 m) and with thioacetic acid (0.6 g, 0.0070 m) in ethanol is treated for 4 min. The formation of reddish brown crude confirmed thiadiazinan-5-one derivatives with excellent yields. <sup>1</sup>H-NMR: 2.13 (s, 1H, NH), 2.68-3.09 (s, 2H, CH<sub>2</sub>), 5.12 (s, 1H, METHINE), 7.17-8.00 (m, Ar-H). <sup>13</sup>C-NMR: 21.21, 31.46, 77.75, 113.09, 121.04, 124.89, 127.72, 128.17, 129.20, 131.55, 133.14, 140.12, 170.80. IR: 3320, 3077, 1696, 1546, 810, 728 cm<sup>-1</sup>. Mass: M/Z 303.

#### **2-(4-hydroxyphenyl)-4-phenyl-1,3,4-thiadiazinan-5-one 1d**

4-hydroxy benzaldehyde (1 g, 0.0081 m), phenyl hydrazine (1.2 g, 0.0081 m) with Thioacetic acid (1:1:1) (0.8 g, 0.0081 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.20 (s, 1H, NH), 3.52-3.61 (s, 2H, CH<sub>2</sub>), 3.94 (s, 1H, METHINE), 5.60 (s, OH), 6.89-7.52 (m, Ar-H). <sup>13</sup>C-NMR: 21.10, 55.29, 76.75, 114.07, 115.96, 119.90, 121.90, 129.03, 129.47, 130.24, 131.13, 132.33, 156.90, 167.55. IR: 3417, 3100, 1666, 1532, 1370, 717 cm<sup>-1</sup>. Mass: M/Z 285.

#### **2-(4-hydroxy-3-methoxyphenyl)-4-phenyl-1,3,4-thiadiazinan-5-one 1e**

A mixture of vanillin (1 g, 0.0066 m), phenyl hydrazine (1.03 g, 0.0066 m) with thioacetic acid (0.6 g, 0.0066 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.07 (s, 1H, NH), 3.03-3.13 (s, 2H, CH<sub>2</sub>), 4.90 (s, 1H, METHINE), 6.95-7.82 (m, Ar-H). <sup>13</sup>C-NMR: 36.80, 76.75, 121.04, 124.80, 127.72, 128.17, 130.50, 131.64, 133.15, 148.70, 165.80. IR: 3328, 3034, 1672, 1548, 1352, 848 cm<sup>-1</sup>. Mass: M/Z 314.

#### **2-phenyl-1,3,4-thiadiazinan-5-one 2a**

The microwave irradiation of equal molar Benzaldehyde (1 g, 0.0094 m), hydrazine (0.9 g, 0.0094 m) and with thioacetic acid (0.8g, 0.0094 m) in ethanol is treated for 4 min. The formation of reddish brown crude confirmed thiadiazinan-5-one derivatives with excellent yields. <sup>1</sup>H-NMR: 2.26 (s, 1H, NH), 3.28-3.30 (s, 2H, CH<sub>2</sub>), 4.39 (s, 1H, METHINE), 7.24-7.98 (m, Ar-H). <sup>13</sup>C-NMR: 44.06, 77.06, 125.75, 126.54, 127.13, 128.54, 129.11, 130.12, 131.19, 132.87, 170.08. IR: 3332, 3085, 1692, 1580, 760 cm<sup>-1</sup>. Mass: M/Z 192.

#### **2-(3-nitro phenyl)-1,3,4-thiadiazinan-5-one 2b**

3-nitro benzaldehyde (1 g, 0.0066 m), hydrazine (0.7 g, 0.0066 m) with Thioacetic acid (1:1:1) (0.6 g, 0.0066 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.32 (s, 1H, NH), 3.12-3.25 (s, 2H, CH<sub>2</sub>), 4.43 (s, 1H, METHINE), 7.07-7.81 (m, Ar-H). <sup>13</sup>C-NMR: 39.20, 79.12, 118.72, 121.02, 123.80, 126.72, 127.17, 131.50, 132.64, 133.14, 148.30, 170.80. IR: 3318, 3026, 1620, 1525, 813 cm<sup>-1</sup>. Mass: M/Z 238.

#### **2-(4-hydroxyphenyl)-1,3,4-thiadiazinan-5-one 2c**

A mixture of 4-hydroxy benzaldehyde (1 g, 0.0081 m), hydrazine (0.8 g, 0.0081 m) with thioacetic acid (0.7 g, 0.0081 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.08 (s, 1H, NH), 3.10-3.23 (s, 2H, CH<sub>2</sub>), 4.71 (s, 1H, METHINE), 5.5 (s, OH), 6.61-7.56 (m, Ar-H). <sup>13</sup>C-NMR: 38.46, 78.07, 115.80, 121.30, 128.60, 130.20, 131.61, 132.80, 134.70, 135.52, 168.31. IR: 3231, 3020, 1690, 1517, 768 cm<sup>-1</sup>. Mass: M/Z 209.

#### **2-(4-hydroxy-3-methoxyphenyl)-1,3,4-thiadiazinan-5-one 2d**

The microwave irradiation of equal molar vanillin (1 g, 0.0064 m), hydrazine (0.68 g, 0.0064 m) and with thioacetic acid (0.6 g, 0.0064 m) in ethanol is treated for 4 min. The formation of reddish brown crude confirmed thiadiazinan-5-one derivatives with excellent yields. <sup>1</sup>H-NMR: 2.31 (s, 1H, NH), 3.28-3.32 (s, 2H, CH<sub>2</sub>), 5.17

(s, 1H, METHINE), 5.40 (s, OH) 6.50-7.68 (m, Ar-H). <sup>13</sup>C-NMR: 28.19, 35.20, 55.02, 67.30, 116.80, 122.50, 127.17, 129.30, 132.82, 133.80, 135.02, 144.8, 151.8, 178.10. IR: 3229, 3018, 1712, 1561, 778 cm<sup>-1</sup>. Mass: M/Z 240.

#### **4-(2,4-dinitrophenyl)-2-phenyl-1,3,4-thiadiazinan-5-one 3a**

Benzaldehyde (1 g, 0.0094 m), 2,4-dinitro phenylhydrazine (1.7 g, 0.0094 m) with Thioacetic acid (1:1:1) (0.8 g, 0.0094 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.28 (s, 1H, NH), 3.02-3.15 (s, 2H, CH<sub>2</sub>), 4.87 (s, 1H, METHINE), 7.01-7.75 (m, Ar-H). <sup>13</sup>C-NMR: 44.06, 77.06, 125.75, 126.54, 127.06, 127.64, 128.28, 129.24, 130.12, 131.19, 172.90. IR: 3326, 3085, 1698, 1562, 1365, 812 cm<sup>-1</sup>. Mass: M/Z

#### **2-(4-hydroxyphenyl)-4-(2,4-dinitro phenyl)-1,3,4-thiadiazinan-5-one 3b**

A mixture of 4-hydroxy benzaldehyde (1 g, 0.0081 m), 2,4-dinitro phenylhydrazine (1.5 g, 0.0081 m) with thioacetic acid (0.7 g, 0.0081 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.36 (s, 1H, NH), 2.68-2.82 (s, 2H, CH<sub>2</sub>), 4.37 (s, 1H, METHINE), 6.61-7.28 (m, Ar-H). <sup>13</sup>C-NMR: 31.93, 76.75, 113.02, 121.04, 124.80, 127.70, 128.14, 130.50, 131.64, 133.15, 133.87, 146.62, 165.60. IR: 3390, 3066, 1712, 1571, 742 cm<sup>-1</sup>. Mass: M/Z 365.

#### **4-(2,4-dinitrophenyl)-2-(3-nitro phenyl)-1,3,4-thiadiazinan-5-one 3c**

3-nitrobenzaldehyde (1 g, 0.0066 m), 2,4-dinitro phenylhydrazine (1.2 g, 0.0066 m) with Thioacetic acid (1:1:1) (0.6 g, 0.0066 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.02 (s, 1H, NH), 2.82-3.12 (s, 2H, CH<sub>2</sub>), 4.32 (s, 1H, METHINE), 6.98-7.98 (s, Ar-H). <sup>13</sup>C-NMR: 30.46, 76.75, 121.04, 124.80, 127.60, 128.10, 130.14, 131.52, 133.11, 140.12, 169.10. IR:

3326, 3033, 1692, 1562, 1523, 770 cm<sup>-1</sup>. Mass: M/Z 394.

#### **2-(4-chlorophenyl)-4-(2,4-dinitro phenyl)-1,3,4-thiadiazinan-5-one 3d**

A mixture of 4-chloro benzaldehyde (1 g, 0.0071 m), 2,4-dinitro phenylhydrazine (1.3 g, 0.0071 m) with thioacetic acid (0.6 g, 0.0071 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.21 (s, NH), 3.23-3.28 (s, 2H, CH<sub>2</sub>), 4.57 (s, METHINE), 7.01-7.51 (m, Ar-H). <sup>13</sup>C-NMR: 32.33, 74.21, 115.96, 117.20, 119.90, 121.81, 127.04, 129.30, 129.47, 130.79, 131.13, 132.33, 152.09, 168.20. IR: 3330, 3073, 1714, 1542, 1502, 832 cm<sup>-1</sup>. Mass: M/Z 383.

#### **2-(4-hydroxy-3-methoxyphenyl)-4-(2,4-dinitro phenyl)-1,3,4-thiadiazinan-5-one 3e**

The microwave irradiation of equal molar vanillin (1 g, 0.0064 m), 2,4-dinitro phenylhydrazine (1.2 g, 0.0064 m) with thioacetic acid (0.5 g, 0.0064 m) in ethanol, is treated for 4 min. The formation of reddish brown crude confirmed thiadiazinan-5-one derivatives with excellent yields. <sup>1</sup>H-NMR: 2.98 (s, 1H, NH), 3.12-3.15 (s, 2H, CH<sub>2</sub>), 3.63, 4.91 (s, 1H, METHINE), 5.02 (1H, OH), 6.40-7.38 (m, Ar-H). <sup>13</sup>C-NMR: 34.17, 77.50, 125.25, 126.54, 127.13, 128.54, 129.11, 130.50, 131.64, 132.08, 141.20, 144.12, 167.20. IR: 3378, 3012, 1698, 1532, 1512, 812 cm<sup>-1</sup>. Mass: M/Z 396.

#### **4-(4-chlorophenyl)-2-phenyl-1,3,4 thiadiazinan-5-one 4a**

Benzaldehyde (1 g, 0.0094 m), 4-chlorophenylhydrazine (1.3 g, 0.0094 m) with thioacetic acid (1:1:1) (0.8 g, 0.0094 m) and added ethanol were treated on microwave irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.12 (s, 1H, NH), 3.33-3.41 (s, 2H, CH<sub>2</sub>), 5.10 (s, 1H, METHINE), 6.88-7.38 (m, Ar-H). <sup>13</sup>C-NMR: 28.09, 75.09, 118.09, 120.03, 123.00, 127.10, 128.08, 129.02, 139.20, 164.70. IR: 3218, 3032, 1665, 1520, 917, 760 cm<sup>-1</sup>. Mass: M/Z 291

#### 4-(4-chlorophenyl)-2-(4-hydroxyphenyl)-1,3,4-thiadiazinan-5-one 4b

A mixture of 4-hydroxybenzaldehyde (1 g, 0.0081 m), 4-chlorophenylhydrazine (1.1 g, 0.0081 m) with thioacetic acid (0.7 g, 0.0081 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.08 (s, 1H, NH), 3.01-3.12 (s, 2H, CH<sub>2</sub>), 4.73 (s, 1H, METHINE), 5.20, 6.60-7.58 (m, Ar-H). <sup>13</sup>C-NMR: 35.80, 75.40, 116.89, 123.03, 129.01, 130.10, 131.08, 133.22, 150.20, 164.02. IR: 3317, 3018, 1698, 1515, 1021, 797 cm<sup>-1</sup>. Mass: M/Z 308.

#### 4-(4-chlorophenyl)-2-(4-hydroxy-3-methoxyphenyl)-1,3,4-thiadiazinan-5-one 4c

The microwave irradiation of equal molar vanillin (1 g, 0.0064 m), 4-chlorophenylhydrazine (0.9 g, 0.0064 m) with thioacetic acid (0.5 g, 0.0064 m) in ethanol, is treated for 4 min. The formation of reddish brown crude confirmed thiadiazinan-5-one derivatives with excellent yields. <sup>1</sup>H-NMR: 2.21 (s, 1H, NH), 3.10-3.22 (s, 2H, CH<sub>2</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 5.12 (s, 1H, METHINE), 5.09, 6.40-7.52 (m, Ar-H). <sup>13</sup>C-NMR: 40.21, 55.03, 78.04, 119.20, 122.05, 123.01, 125.60, 129.87, 133.08, 142.20, 163.02. IR: 3351, 3081, 1712, 1530, 917, 817 cm<sup>-1</sup>. Mass: M/Z 323.

#### 4-(4-chlorophenyl)-2-(3-nitrophenyl)-1,3,4-thiadiazinan-5-one 4d

3-nitrobenzaldehyde (1 g, 0.0066 m), 4-chlorophenylhydrazine (0.9 g, 0.0066 m) with Thioacetic acid (1:1:1) (0.6 g, 0.0066 m) and added ethanol were treated on microwave

irradiation for 2-3 min. The formation of the red orange solid mass of thiadiazinan-5-one derivatives is confirmatory. <sup>1</sup>H-NMR: 2.08 (s, 1H, NH), 2.87-2.92 (s, 2H, CH<sub>2</sub>), 4.71 (s, 1H, METHINE), 6.82-7.71 (m, Ar-H). <sup>13</sup>C-NMR: 36.18, 74.40, 119.20, 122.02, 123.08, 127.20, 129.18, 131.62, 132.21, 134.09, 167.30. IR: 3321, 3023, 1712, 1535, 930, 812 cm<sup>-1</sup>. Mass: M/Z 337.

#### 4-(4-chlorophenyl)-2-(4-chlorophenyl)-1,3,4-thiadiazinan-5-one 4e

A mixture of 2-hydroxybenzaldehyde (1 g, 0.0081 m), 4-chlorophenylhydrazine (1.1 g, 0.0081 m) with thioacetic acid (0.7 g, 0.0081 m) and added ethanol is treated on microwave irradiation for 4 min. The crude product is obtained. The crude was obtained as reddish brown which confirms the product as thiadiazinan-5-one derivatives. <sup>1</sup>H-NMR: 2.14 (s, 1H, NH), 2.98-3.12 (s, 2H, CH<sub>2</sub>), 4.94 (s, 1H, METHINE), 6.57-7.85 (m, Ar-H). <sup>13</sup>C-NMR: 41.20, 77.07, 124.09, 125.83, 129.21, 128.60, 131.21, 133.02, 134.80, 143.20, 168.12. IR: 3300, 3042, 1694, 1517, 1010, 787 cm<sup>-1</sup>. Mass: M/Z 308.

#### Antibacterial activity

All the newly synthesized thiadiazin-5-one were evaluated for antibacterial activity by disc diffusion technique against gram positive organism *Staphylococcus aureus* and gram negative organism, *Escherichia coli*. Norflaxin (for bacteria) was used as the standard. 10 µg/µl concentrations of the newly synthesized compounds were prepared using chloroform as the solvent. Disc containing only chloroform was used as negative control. The results of antibacterial activity are listed in table 2.

Table 2. Antimicrobial activity of Ph and Hy sample against the bacteria clinical isolates by disc diffusion method

Sample id	Name of the bacteria	Zone of inhibition (mm)						
		5µg/µl	10µg/µl	15µg/µl	20µg/µl	25µg/µl	30µg/µl	10µg/µl Norflaxin
Ph	<i>E.coli</i>	0	0	0	0	0	0.1	0.5
Ph	<i>Proteus vulgaris</i>	0	0	0	0	0	0	0.1
Ph	<i>S. aureus</i>	0	0	0	0	0	0	0.1
Hy	<i>E.coli</i>	0	0	0	0	0	1.8	1.5
Hy	<i>Proteus vulgaris</i>	0	0	0	0	0	0	0.1

Hy	<i>S. aureus</i>	0	0.1	0.3	0.4	0.5	0.7	1.5
----	------------------	---	-----	-----	-----	-----	-----	-----

### Antifungal activity

The newly synthesized thiadiazin-5-one was evaluated for antifungal activity by disc diffusion technique against *Aspergillum niger*, *Aspergillus fumigatus*, *Klebesiella pneumonia*. 10 µgm/ml disc of Fluconazole (*Aspergillum niger*, *Aspergillus fumigatus*); Norflorin

(*Klebesiella pneumonia*) (Fungi) was used as the standard. Disc containing 10 µg/ml concentrations of the newly synthesized compounds as test and 10 µgm/ml disc of chloroform was used as a negative control. The results of antifungal activity are listed in table 3.

Table 3. Antimicrobial activity of Ph and Hy sample against the fungal clinical isolates by disc diffusion method

Sample Id	Name of the microorganism	Zone of inhibition (mm)						Standard
		5 µg/µl	10 µg/µl	15 µg/µl	20 µg/µl	25 µg/µl	30 µg/µl	
Ph	<i>Aspergillum niger</i>	1.5	1.8	2	2.1	2.2	2.3	1
Hy	<i>Aspergillum niger</i>	1	1.7	1.8	1.9	2	2.1	1.2
Ph	<i>Aspergillum fumigatus</i>	0	0	0	0	0	0.1	1.5
Hy	<i>Aspergillum fumigatus</i>	0	0	0	0	0.5	1	2.5
Ph	<i>Klebesiella pneumonia</i>	0	0	0	0	0	0.1	0.1
Hy	<i>Klebesiella pneumonia</i>	0.1	1	1.3	1.5	1.7	1.8	0.1

Standard: Fluconazole (*Aspergillum niger*, *Aspergillus fumigates*), Norflorin (*Klebesiella pneumonia*)

### Conclusions

In summary, title compounds by one pot, three-component synthesis was developed in practical and green synthetic methods with good yields. These methods uses short reaction time, mild condition with simple work-up procedure and environmentally begin process. Therefore in present work, thiadiazin-5-one was synthesized by condensation of substituted aromatic aldehyde and substituted hydrazine in presence of thioacetic acid and ethanol, as a green solvent. All the target compounds were synthesized in only one step with over all yields 85%-90%, respectively. The structures of these novel targets and part of intermediates were confirmed by <sup>1</sup>H, <sup>13</sup>C NMR and IR spectra.

### References

- Hunasha RD, Satyanarayana D. One pot synthesis of 3-(substituted phenoxyethyl)-6-phenyl/substituted phenoxyethyl-1,2,4-triazolo[3,4-b][1,3,4] thiadiazole derivatives as antimicrobial agents. International Journal of Pharma and Bio Sciences 3 (2012) 183-192.
- Ritter SK. EPA analysis suggests green success. Chemical Engineering News 79 (2001) 27-34.
- Moghanian H, Ebrahimi A. Three Component, One pot synthesis of amido-alkyl naphthols using polyphosphate ester under solvent free conditions. Journal of Saudi Chemical Society 2 (2014) 165-168.
- Nalini V, Girija R. Synthesis and characterization and biological studies of 9-aryl substituted acridinedione derivatives by Hantzsch condensation. International journal of current research 5 (2013) 3076-3081.
- Sanjeeva Reddy Ch, Sanjeeva Rao L, Sunitha B, Nagaraj A. Synthesis and antibacterial activity of N-Substituted-[1,2,4] triazoles and 1,2,4-triazole [3,4-b] [1,3,4] thiadiazines. Indian Journal of Chemistry 54B (2015) 1283-1289.
- Deohate PP. Synthesis, structural study and biological activity of bridgehead nitrogen containing Triazolo-thiadiazine derivatives. Chemical Science Transactions 2 (2013) 556-560.
- Kharate RM, Deohate PP, Beard BN. Synthesis structural study and antimicrobial screening of bridgehead nitrogen containing 1,8, bis-thiadiazino-triazolyl-octanes. Der Pharma Chemical 4 (2012) 2434-2437.
- Deohate PP. Synthesis, structural study and biological activity of bridgehead nitrogen containing Triazolo-thiadiazine

- heterocycles. *Der Pharma Chemical*, 4 (2012) 2042-2046.
9. Mathew V, Keshavayya J, Vaidya VP. Heterocyclic system containing bridgehead nitrogen atom: synthesis and pharmacological activities of some substituted 1,2,4-triazolo[3,4-b]-1,3,4-thiadiazoles, *European Journal of Medicinal Chemistry*, 9 (2006) 1048-1058.
  10. Mathew V, Keshavayya J, Vaidya VP, Giles D. Studies on synthesis and pharmacological activities of 3,6-disubstituted-1,2,4-triazolo[3,4-b]-1,3,4-thiadiazoles and their dihydro analogies. *European Journal of Medicinal Chemistry* 6 (2007) 823-840.
  11. Demirbas N, Demirbas A, Karaoglu SA, Celik E. Synthesis and antimicrobial activities of some new [1,2,4] triazolo [3,4-b] [1,3,4] thiadiazoles and [1,2,4] triazolo [3,4-b] [1,3,4] thiadiazines. *ARKIVOC*, 1 (2005) 75-91.
  12. Kirankumar M, Sunitha K, Bhupathi Reddy K, Jayanthkumar M, Shobha Rani N, Mohan R, Vijayakumar B. Synthesis and biological evaluation of new 10-(6-substituted-[1,2,4]-triazolo-[3,4-b] [1,3,4] thiadiazol-3-ylmethyl)-3-nitro-10H-phenoxazine and 10-(6-substituted -7H-[1,2,4]-triazolo-[3,4-b][1,3,4]-thiaiazol-3-yl methyl)-3-nitro-10H-phenoxazine. *World Journal of Pharmacy and Pharmaceutical Sciences* 2 (2013) 5284-5295.
  13. Sridhara AM, Venugopala Reddy KR, Keshavayya J, Chdanda B. Synthesis and antimicrobial activity of pthalazine substituted 1,2,4-triazolo-[3,4-b]-1,3,4-thiadiazoles and 7H-[1,2,4]triazolo[3,4-b][1,3,4]thiadiazines. *Der Pharma Chemical* 2 (2010) 201-211.
  14. Siddiqui MT, Comrie MH. Antituberculosis agents. V:  $\alpha$ -[5-(5-nitro-2-furyl)-1,3,4-oxadiazol-2-ylthio] acetylhydrazide and related compounds. *Journal of Pharmaceutical Sciences* 6 (1991) 548-550.
  15. Venkata S, Rao C, Rajeswar RV. One-Pot synthesis of 1,3,4-Thiadiazin-5-yl-chromen-2-one Derivatives via Three-Component Reaction. *Synthetic Communications* 42 (2012) 1454-1460.
  16. Schroder J, Henke A, Wenzel H, Brandstetter H, Stammler HG, Stammler A, Pfeiffer WD, Tschesche H. Structure-based design and synthesis of potent matrix metalloproteinase inhibitors derived from a 6H-1,3,4-thiadiazine scaffold. *J Med Chem* 44 (2001) 3231-3243.
  17. Sugawara H, Endoh M. Enantiomer EMD 57439 antagonizes the Ca<sup>2+</sup> sensitizing effect of (p)-enantiomer EMD 57033 on diastolic function but not on systolic function in rabbit ventricular cardiomyocytes. *Jpn J Pharmacol* 80 (1999) 55-65.
  18. Himmel HM, Amos GJ, Wettwer E, Ravens U. Effects of the calcium sensitizer [p]-EMD 60263 and its enantiomer-EMD 60264 on cardiac ionic currents of guinea pig and rat ventricular myocytes. *J Cardiovasc Pharmacol* 33 (1999) 301-308.
  19. Milan C, Valentina Pavic, Maja Molnar, Bojansarkanj, Elizabeta Has-Schon. 2Design and synthesis of some new 1,3,4-thiadiazines with coumarin moieties and their antioxidative and antifungal activity. *Molecules* 19 (2014) 1163-1177.
  20. Sarkanj B, Molnar M, Cacic M, Gille L. 4-Methyl-7-hydroxycoumarin antifungal and antioxidant activity enhancement by substitution with thiosemicarbazide and thiazolidinone moieties. *Food Chem* 139 (2013) 488-495.
  21. Novikova AP, Perova NM, Chupakhin ON. Synthesis and properties of functional derivatives of 1,3,4-thiadiazines and condensed systems based on these compounds. *Khim Geterot Soedin* 11(1991) 1443-1457.

\*\*\*\*\*