

Mercury Contamination of White Cheek Shark (*Carcharhinus dussumieri*): its Relation to the Length and SexSomayeh Torabi Delshad^{1*}, Seyed Abdolmajid Mousavi², Houman Rajabi Islami¹, Abdolrahim Pazira³¹ Department of Fisheries, Science and Research Branch, Islamic Azad University, P.O Box: 14155-4933, Tehran, Iran.² Department of animal Science, Agriculture Faculty, Varamin Branch, Islamic Azad University, Varamin, Iran.³ Department of Fisheries, Boushehr Branch, Islamic Azad University, Boushehr, Iran.* Corresponding author: st.delshad@yahoo.com

Abstract: This study was conducted to assess the total mercury concentration in muscle, liver and fin tissues of whitecheek shark (*Carcharhinus dussumieri*, Carcharhinidae). Approximately 26 percent of all captured Sharks had total mercury concentrations greater than 0.2 mg/kg w.w., 48% contained amounts higher than 0.4 mg/kg w.w. and 22 % had more than 0.6 mg/kg w.w. Only 4 % of all samples had a concentration greater than 0.8 mg/kg w.w. In spite of significant differences in the length distributions of male and female, there were no significant differences in total mercury concentrations of experimental tissues between the genders. In all stations, muscle tissue showed the highest mercury levels (0.73 mg/kg w.w in males and 0.77 mg/kg w.w in females), followed by Liver (0.28 mg/kg w.w. in males and 0.29 mg/kg w.w. in females) and Fins (0.13 mg/kg and 0.16 mg/kg w.w., respectively). No significant correlation was found between the concentration of mercury in tissues with sex and location, although length had an increasing effect on mercury concentration. In addition, length showed strongly positive correlation with mercury concentration of fins. The current study illustrated that consumption of less than 0.5 kg/week from muscle tissue of the whitecheek shark could result in a daily intake of 80 µg Hg which is more than 45 times the maximum intake concentration established by the World Health Organization.

[Torabi Delshad S, Mousavi SA, Rajabi Islami H, Pazira A. **Mercury Contamination of Whitecheek Shark (*Carcharhinus dussumieri*): its Relation to the Length and Sex.** *Biomedicine and Nursing* 2015;1(3):68-74]. ISSN 2379-8211 (print); ISSN 2379-8203 (online); <http://www.nbmedicine.org>. 7. doi:[10.7537/marsbnj010315.07](https://doi.org/10.7537/marsbnj010315.07)

Keywords: Mercury, Whitecheek shark, Muscle, Persian Gulf**1. Introduction**

Mercury (Hg) contamination is a great concern which its toxicity in many coastal ecosystems poses some health hazard for human and wildlife (Branco et al., 2004; Tessier et al., 2007; Saniewska et al., 2010). Methylmercury (CH₃Hg⁺) is the most toxic of the mercury compounds that accounts for more than 95% of organic mercury in aquatic muscle tissues (Jewett et al., 2003; Bloom et al., 2004; Branco et al., 2004; Covelli et al., 2009; Liu et al., 2012). The main routes of acute and chronic Hg exposure include inhalation, dermal absorption, and ingestion (Solis et al., 2000; Moreno et al., 2005; Harper and Harris, 2008; Al-Saleh et al., 2009; Burke et al., 2010). Also, toxic responses such problem as malformations, growth reduction, neurological deficiency in levels of certain enzymes, and renal failure are examples of chemically induced effects of Hg exposure (Lombourdis and Danscher, 2004; Bose-O'Reilly et al., 2010; Rani et al., 2011).

World Health Organization (WHO) had been estimated that around 10×10⁶ kg mercury is annually released all over the world (WHO, 1989). About 50 percent of this amount transport from continents to the coastal area by pathways such as surface runoff, atmospheric deposition mainly fluvial transport, and natural weathering processes (Hanten et al., 1998;

Babiarz et al., 2003; Carvalho et al., 2008; Stern et al., 2012).

Since degradation of mercury compounds in environment has some complications, the most mercury contaminations in food chain come mainly from existing background mercury pollutants (Merritt and Amirbahman, 2009; Nfon et al., 2009; Lavoie et al., 2010). Recently, public health concerns over mercury toxicity have focused on the potential risk associated by relatively low doses of Hg in the environment (Hsiao et al., 2010; Fang et al., 2011; Huang et al., 2011). Several studies have shown that mercury could bioaccumulate, mainly as methylated forms, in the muscle tissues of aquatic organisms (EPA, 2001; Cai et al., 2007; Negrete et al., 2008). Probably due to some difficulties in sample catching of apex predators like sharks from marine, there are scarce researches on acute and chronic effects of mercury in marine environments (Feng et al., 2004).

Because of predatory behavior, long life, and higher trophic levels, sharks exhibit higher mercury concentrations than other marine fishes (Da Silva et al., 2004; Endo et al., 2008). Whitecheek shark (*C. dussumieri*) is one of the most plentiful sharks of the Persian Gulf which has been inhabited in coastal ecosystem, where point sources of mercury are abundant (Henderson et al., 2007). This ecologically

and economically important species live in continental shelf of south the Indian Ocean with preferable depth of 20 to 60 m and fed from cephalopods, crustaceans, and small fishes (Salini et al., 1992, 1994; Cortés, 1999; Henderson et al., 2007).

Meat and fins of whitecheek shark are used for human consumptions and could consequently represent a model for the study of mercury contaminations in marine environments and the possible intake of mercury by human populations (Ferreira et al., 2004). Accordingly, the objective of the present study was to analysis total mercury concentration in muscle, liver and fin tissues of whitecheek shark, *C. dussumieri*, from the Iranian coastal waters of the Persian Gulf. Besides, correlation between mercury levels and body length, age, and sex of sharks was also considered to clarify risks associated by consumption of the marine resources.

2. Material and Methods

Fish sampling was performed from 3 different stations (Genaveh, Boushehr and Dayer) located in northern coast of the Persian Gulf (50-51° N and 27-29° E), within 20 km of the shoreline (Fig. 1). A total of 149 specimens whitecheek shark (*C. dussumieri*) were captured opportunistically by a local trawl fishing boat from November to December 2011. After capture, all sharks were individually weighted, sized and frozen for preservation before being transported to the laboratory. Besides, the age of each sample was evaluated using the correspondent length of maturity recommended by Compagno (1984). Specimens shorter than the maturity length were considered juveniles, and individuals equal to or larger than the maturity length were considered adults.

Tissue samples were taken in the laboratory by necropsies of each specimen. A portion (approximately 20 g) of muscle, liver and fin were segregated using a clean stainless-steel knife. Care was done to ensure that the samples had no contact with dermal layers or other surrounding surfaces during the dissection. After that, shark tissues washed 3 times with tap water and rinsed by deionized water. To reduce the risk of contamination, each sample were immediately placed in a separate plastic zip-lock bag and frozen at -20°C before analysis. Sex was also determined macroscopically and checked by examination of interior reproductive organs.

Analytical Methods

All samples were processed within two months of being captured. Approximately 2 g of experimental tissues (liver, muscle and fin) were freeze-dried at -50°C and their moisture contents determined by weight loss. Then, the samples were ground,

homogenized and sieved ($d \leq 175 \mu\text{m}$) for investigation.

The amount of total mercury (THg) in liver, muscle and fin was separately determined by an advanced mercury analyzer (LECO model AMA 254, USA), which did not require pretreatment or acid-digestion of the sample. In brief, aliquots ranging from 20 to 40 mg of freeze-dried sample were placed into oven of the instrument. After drying, each sample was pyrolyzed at 800 °C under oxygen atmosphere for 3 min and elemental mercury vapor was subsequently collected in a gold net (Au-amalgamator). The net was then heated for liberating and measuring of Hg by atomic absorption spectrometry (AAS). Each sample was analyzed in triplicate for assurance of consistent results.




Fig. 1. Study area and topology of sampling points.

The accuracy of the procedure was examined by analyzing of certified reference materials including dogfish muscle (DORM-2), dogfish liver (DOLT-3), and lobster hepatopancreas (TORT-2) obtained from the National Research Council of Canada. The standard materials were analyzed according to the procedure described for the tissue samples of whitecheek shark. A suitable conformity was found between certified and obtained values, as recovery ranged from 98.3% to 103.4%. All total mercury results were stated as milligrams per kilogram (mg/kg) of wet weight (w.w.).

3. Results

The total number and body length of the whitecheek shark captured from different stations are shown in Table 1. Of total 149 specimens, the most samples were collected from Dayer station (42.3 %) followed by Genaveh (32.8 %) and Bushehr (24.9 %) stations. The lengths of the sharks varied from 65 to 105 cm. The mean body size of the individuals

showed that all specimens were adult (Based on Compagno, 1984). The total shark samples were composed of 59 (41.6 %) males and 88 (58.4 %) females. Body length of the female samples was significantly higher than males in all sampling station ($p < 0.05$), except for Genaveh station which no significant differences was found between males and females. All the shark specimens were showed significant positive correlations between body length and weight ($p < 0.05$). Therefore, only relationship between body length and THg levels is reported here.

Overall variations of mercury concentration in different tissues of whitecheek shark were high and concentrations varied, but were usually high in the study area (Table II). Approximately 26 percent of all Sharks had total mercury concentrations greater than 0.2 mg/kg w.w., 48% contained amount higher than 0.4 mg/kg w.w., and 22 % had more than 0.6 mg/kg w.w. Only 4 % of all samples had a concentration greater than 0.8 mg/kg w.w.

Although females had higher length than males, no significant difference was obtained in total mercury concentrations of experimental tissues between the genders ($p > 0.05$). In all stations, muscle tissue showed the highest mercury levels (Fig. 1), with an average concentration of 0.73 ± 0.28 mg/kg w.w. in males and 0.77 ± 0.27 mg/kg w.w. in females, followed by the Liver (0.28 ± 0.9 mg/kg and 0.29 ± 0.9 mg/kg w.w., respectively). Fins had the least tissue contamination in all cases, with mercury concentrations varied between 0.13 ± 0.05 to 0.16 ± 0.07 mg/kg w.w.

No significant correlations was found between the concentration of mercury in tissues with sex and location, although length had an increasing effect on mercury concentration ($p > 0.05$) (see Table 2). In addition, length showed strongly positive correlation by mercury concentration of fins, whereas other variables like sex and station had not significant effect on fins mercury (Table 3).

Cluster analysis was also applied to provide an outline of similarities between mercury concentrations of different tissues (Fig. 2). It is clear that pattern of mercury level in muscle and fin is closer, whereas kidney concentration was different from the others.

4. Discussions

The current research provides information concerning total mercury levels in muscle, liver, and fin of whitecheek shark, *C. dussumieri*, as a top marine predator caught from northern offshore of the Persian Gulf. Based on the amount of natural mercury in fish (0.15 to 0.2 $\mu\text{g/g}$ wet weight) provided by Johnels et al. (1967), the values found for this species were considered high. Overall concentrations were also high when the results compared to the levels of mercury in the other shark species captured from Florida (Adams and McMichael, 1999), Brazil (Pinho et al., 2002), western equatorial Atlantic Ocean (Ferreira et al., 2004), and western equatorial Pacific Ocean (Endo et al., 2008). The present difference could be related to distinct feeding habits and other species-specific parameters of *C. dussumieri* such as metabolic rate and lifetime.

There are few studies on the segregation of mercury in fish. Coelho et al. (2010) reported mercury content in various tissues of demersal shark and chimaeras from continental shelf and slope waters of southeast Australia. In all specimens, the highest mercury level was recorded in muscle tissue, followed by the heart, gills, liver, and Pancreas. Endo et al. (2008) stated that the average of Hg values in the muscle tissues of tiger and silvertip sharks were higher than that in the corresponding liver tissues. Other shark species have also indicated higher value of total mercury in muscle compared to liver (Taguchi et al., 1979; Lacerda et al., 2000; Adams et al., 2005; Branco et al., 2007; Pethybridge et al., 2010).

Table 1. Number of the fish captured from different station of the Persian Gulf.

	Number			Length (cm)		
	Total	Female	Male	Mean	Female	Male
Genaveh	49	28	21	82.5 ± 7.7	83.8 ± 7.1	$80.2 \pm 8.1^*$
Bushehr	37	22	15	84.7 ± 9.2	90.0 ± 9.2	$79.6 \pm 9.4^*$
Dayer	63	37	26	87.6 ± 9.6	93.5 ± 10.7	$81.3 \pm 8.6^*$

* Shows significant difference at level of p less than 0.05.

Similarly, the present results showed the highest mercury level in the muscle tissue. The liver exhibited comparatively lower mercury level followed by the fin. Current pattern corroborates the hypothesis that mercury is more assimilated from the dietary exposure than the environment, and is easily distributed

throughout the body and preferential accumulated in muscle tissue (Lacerda et al., 2000; Wang, 2002; Coelho et al., 2008).

Boening (2000) cleared that dietary behavior has an important effect on mercury accumulation in top marine predators which the results presented in Table

2 support this view. The mercury content in the diet increases as the shark grows larger (Mathers and Johansen, 1985). Whitecheek shark feed on a wide range of crustaceans but preys cephalopods and small benthic as they became greater in size (Henderson et al., 2007; White et al., 2012). Therefore, higher mercury accumulation in muscle is generally associated by dietary uptake (Branco et al., 2007; Pethybridge et al., 2010), while environmental absorption has lower impact on external organs such the skin and fins (Coelho et al., 2010; Pethybridge et al., 2010). Median concentration of mercury in liver could be explained by demethylation mechanism of mercury occurring in liver tissue for elimination and Neutralization of the harmful effect of mercury in sharks (Palmisano et al., 1995; Storelli et al., 2002; Branco et al., 2007).

The current findings support this view that mercury level in muscle tissue has a linear relationship by size/age of the shark (Green and Knutzen, 2003; Branco et al., 2007; Pethybridge et al., 2010). The continuing accumulation may be explained by strong binding of mercury in muscle to trial groups of proteins whose content expands with age (Sfezer et al., 2003; Storelli et al., 2002). Besides, lower growth rates in adult individuals lead to longer exposure and, therefore, more accumulation of mercury (Pinho et al., 2002). On the other hand, the content of Hg in liver was poorly correlated by individuals' lengths which demonstrate that accumulation in this organ does not proceed essentially by age (Branco et al., 2007). The processes of detoxification in liver could cause the elimination of toxicant forms of mercury arrived to this organ (Storelli et al., 2002).

There was no significant difference in total mercury concentration of whitecheek shark between the stations, although Bushehr station showed a slight higher level of contamination. Similar results of mercury accumulation in specimens between stations may be related to their highly migratory nature and identical absorption through the body.

There was no significant difference in total mercury concentration between genders of whitecheek shark, although females exhibited slightly higher level of contamination than males. Differences in mercury levels of males and females may be caused by factors such as energetic requirements, maturation condition, mercury deposition, and transference to eggs and fetuses as previously reported for the other shark species (Taguchi et al., 1979; Walker, 1988; Endo et al., 2008). In the present work, higher mercury levels were found in males, although there were significantly shorter on average than females ($p < 0.05$). Differences in mercury concentration could also be clarified by higher growth rate in females, resulting younger female with same length and time for mercury exposure than males (Forrester et al., 1972; Lyle, 1984; Monteiro et al., 1991; Licata et al., 2005). The lower mercury levels in females may be explained by viviparous reproduction of *Carcharhinus* species and nutrients transfer to the fetus through the blood (Lyle, 1984; Compagno et al., 1984; Henderson et al., 2007; Holms et al., 2009). Thus, mercury present in the mother will be easily deposited in the fetus and, therefore, reduced in females compared to that the males.

Table 2. Total mercury concentration (mg/kg wet weight) in tissues of whitecheek shark from different stations of the Persian Gulf.

	Muscle				Liver				Fin			
	Female	Male	Max	Min	Female	Male	Max	Min	Female	Male	Max	Min
Genaveh	0.75±0.20 ^{ef}	0.72±0.27 ^c	1.20	0.18	0.26±0.10 ^c	0.25±0.09 ^c	0.41	0.02	0.15±0.09 ^b	0.15±0.06 ^b	0.24	0.01
Bushehr	0.76±0.29 ^{ef}	0.74±0.33 ^c	1.51	0.10	0.33±0.12 ^d	0.29±0.10 ^{cd}	0.58	0.08	0.15±0.07 ^b	0.16±0.05 ^b	0.29	0.07
Dayer	0.77±0.32 ^{ef}	0.71±0.26 ^c	1.35	0.34	0.30±0.10 ^{cd}	0.30±0.07 ^{cd}	0.47	0.06	0.18±0.04 ^b	0.10±0.05 ^a	0.30	0.07
Total	0.77±0.27 ^{ef}	0.73±0.28 ^c	1.51	0.10	0.29±0.10 ^{cd}	0.28±0.09 ^c	0.41	0.02	0.16±0.07 ^b	0.13±0.05 ^{ab}	0.24	0.01

Results are present as Mean±SE. means with different superscript are significantly different at level of $p < 0.05$.

The whitecheek shark is traditionally consumed by people living in the northern offshore of the Persian Gulf and the eaten of this shark could be one of the major origins of Hg exposure in the human foods. The average of total mercury concentration found in the meat (muscle) of whitecheek shark was 0.73 ± 0.27 g/kg wet weight. The joint FAO/WHO Expert Committee on Food Additives (JECFA) recommended the provisional tolerable weekly intake (PTWI) of $1.6 \mu\text{g/kg}$ of body weight for organic mercury equivalent to $300 \mu\text{g}$ of total mercury per person (WHO, 2007). Accordingly, consumption of even less than 0.5 kg muscle tissue of the whitecheek

shark per week could result in a daily intake of $75 \mu\text{g/kg}$ of body weight Hg which is more than 45 times the maximum intake concentration established by the World Health Organization (WHO, 2007; Ferreira et al., 2004). However, lower concentration of THG in the fin let it to be more consumed.

Acknowledgements

The authors are grateful from the staff of the Zakaria-e-Razi Laboratory Complex of Islamic Azad University for their active participation in the production of the experimental place and technical supports. Thanks also to R. Assare for assisting data

analysis and careful maintenance of the fish during experiment.

Corresponding Author:

Somayeh Torabi Delshad

Department of Fisheries, College of agricultural and Natural Resources, Islamic Azad University, Science and Research Branch, P.O. Box: 14515-775, Tehran, Iran.

E-mail: st.delshad@yahoo.com

References

- Adams DH, RH McMichael. Mercury levels in four species of shark from the Atlantic coast of Florida. *Fishery Bulletin* 1999; 97: 372-379.
- Adams DA, GV Onorato. Mercury concentrations in red drum, *Sciaenops ocellatus*, from estuarine and offshore waters of Florida. *Marine Pollution Bulletin* 2005; 50: 291-30.
- Al-Saleh I, N Shinwari, M Al-Amodi. Accumulation of mercury in ovaries of mice after the application of skin-lightening creams. *Biological Trace Element Research* 2009; 131(1): 43-54.
- Babiarz CL, JP Hurley, DP Krabbenhoft, C Gilmour, BA Branfireun. Application of ultrafiltration and stable isotopic amendments to field studies of mercury partitioning to filterable carbon in lake water and overland runoff. *Science of the Total Environment* 2003; 304: 295-303.
- Bloom NS, LM Moretto, P Scopece, P Ugo. Seasonal cycling of mercury and monomethyl mercury in the Venice Lagoon (Italy). *Marine Chemistry* 2004; 91: 85-99.
- Boening DW. Ecological effects, transport, and fate of mercury: A general review. *Chemosphere* 2000; 40: 1335-1351.
- Bose-O'Reilly S, KM McCarty, N Steckling, B Lettmeier. Mercury Exposure and Children's Health. *Current Problems in Pediatric and Adolescent Health Care* 2010; 40: 186-215.
- Branco V, J Canário, C Vale, J Raimundo, C Reis. Total and organic mercury concentrations in muscle tissue of the blue shark (*Prionace glauca* L.1758) from the Northeast Atlantic. *Marine Pollution Bulletin* 2004; 49: 854-874.
- Branco V, C Vale, J Canário, MN dos Santos. Mercury and selenium in blue shark (*Prionace glauca*, K. 1758) and swordfish (*Xiphias gladius*, L. 1758) from two areas of the Atlantic Ocean. *Environmental Pollution* 2007; 150: 373-380.
- Burke JN, CM Bergeron, BD Todd, WA Hopkins. Effects of mercury on behavior and performance of northern two-lined salamanders (*Eurycea bislineata*). *Environmental Pollution* 2010; 158: 3546-3551.
- Cai Y, JR Rooker, GA Gill, JP Turner. Bioaccumulation of mercury in pelagic fishes from the northern Gulf of Mexico. *Canadian Journal of Fisheries and Aquatic Sciences* 2007; 64: 458-469.
- Carvalho CML, AINM Matos, ML Mateus, APM Santos, MCC Batoréu. High-Fish Consumption and Risk Prevention: Assessment of Exposure to Methylmercury in Portugal. *Journal of Toxicology and Environmental Health-Part A* 2008; 71: 1279-1288.
- Coelho JP, AT Reis, S Ventura, ME Pereira, AC Duarte, MA Pardal. Pattern and pathways for mercury lifespan bioaccumulation in *Carcinus maenas*. *Marine Pollution Bulletin* 2008; 56: 1104-1110.
- Coelho JP, H Santos, AT. Reis, J Falcao, ET Rodrigues, ME Pereira, Duarte AC, MA Pardal. Mercury bioaccumulation in the spotted dogfish (*Scyliorhinus canicula*) from the Atlantic Ocean. *Marine Pollution Bulletin* 2010; 60(8): 1372-1375.
- Compagno LJV. FAO species catalogue Vol 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2. FAO Fisheries Synopsis 1984; 4(125): 251-655.
- Cortés E. Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science* 1999; 56: 707-717.
- Covelli S, A Acquavita, R Piani, S Predonzani, C De Vittor. Recent contamination of mercury in an estuarine environment (Marano lagoon, Northern Adriatic, Italy). *Estuarine, Coastal and Shelf Science* 2009; 82: 273-284.
- Da Silva DS, M Lucotte, M Roulet, H Poirier, D Mergler, EO Santos, M Crossa. Trophic structure and bioaccumulation of mercury in fish of three natural lakes of the Brazilian Amazon. *Water, Air, and Soil Pollution* 2005; 165: 77-94.
- Endo T, Y Hisamichi, K Haraguchi, Y Kato, C Ohta, N Koga. Hg, Zn and Cu levels in the muscle and liver of tiger sharks (*Galeocerdo cuvier*) from the coast of Ishigaki Island, Japan: Relationship between metal concentrations and body length. *Marine Pollution Bulletin* 2008; 56: 1774-1780.
- EPA. Water quality criterion for the protection of human health: methylmercury. Office Science and Technology, Office of Water, Washington, 2001; 253 p.
- Fang F, H Wang, Y Lin. Spatial distribution, bioavailability, and health risk assessment of soil Hg in Wuhu urban area, China. *Environmental Monitoring and Assessment* 2011; 179(1-4): 255-265.
- Feng W, M Wang, B Li, J Liu, Z Chai, J Zhao, G Deng. Mercury and trace element distribution in organic tissues and regional brain of fetal rat after in utero and weaning exposure to low dose of inorganic mercury. *Toxicology Letters* 2004; 25: 223-234.
- Ferreira AG, VV Faria, CEV Carvalho, RPT Lessa, FMS Silva. Total Mercury in the Night Shark,

- Carcharhinus signatus*. Brazilian Archives of Biology and Technology 2004; 47(4): 629-634.
24. Forrester CR, KS Ketchen, CC Wong. Mercury content of spiny dogfish (*Squalus acanthias*) in the Strait of Georgia, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 1972; 29: 1487- 1490.
 25. Green NW, J Knutzen. Organohalogenes and metals in marine fish and mussels and some relationships to biological variables at reference localities in Norway. Marine Pollution Bulletin 2003; 46: 362-377.
 26. Hanten RP, RM Neumann, SM Ward. Relationships between concentrations of mercury in largemouth Bass and physical and chemical characteristics of Connecticut Lakes. Transactions of the American Fisheries Society 1998; 127: 807-818.
 27. Harper BL, SG Harris. A possible approach for setting a mercury risk-based action level based on tribal fish ingestion rates. Environmental Research 2008; 107(1): 60-68.
 28. Henderson AC, JL McIlwain, HS Al-Oufi, S Al-Sheili. The Sultanate of Oman shark fishery: Species composition, seasonality and diversity. Fisheries Research 2007; 86: 159-168.
 29. Holmes BH, D Steinke, RD Ward. Identification of shark and ray fins using DNA barcoding. Fisheries Research 2009; 95: 280-288.
 30. Hsiao H, SM Ullrich, TW Tanton. Burdens of mercury in residents of Temirtau, Kazakhstan: II: Verification of methodologies for estimating human exposure to high levels of Hg pollution in the environment. Science of the Total Environment 2010; 408(19): 4033-4044.
 31. Huang C, S Liu, C Hsu, S Lin-Shiau. Neurotoxicological effects of low-dose methylmercury and mercuric chloride in developing offspring mice. Toxicology Letters 2011; 201(3): 196-204.
 32. Huckabee JW, JW Elwood, SG Hildebrand. Accumulation of mercury in freshwater biota. In: The biogeochemistry of mercury in the environment (J.O. Nriagu ed), 1979; pp. 277-302. Elsevier, Amsterdam.
 33. Jackson TA. Biological and environmental control of mercury accumulation by fish in lakes and reservoirs of north Manitoba. Canadian Journal of Fisheries and Aquatic Science 1990; 44: 313-323.
 34. Jewett SC, X Zhang, AS Naidu, JJ Kelley, D Dasher, LK Duffy. Comparison of mercury and methylmercury in north pick and Arctic grayling from western Alaska Rivers. Chemosphere 2003; 50: 383-92.
 35. Johnels AG, T Westermark, W Berg, PI Persson, B Sjostrand. Pike (*Esox lucius* L.) and some other aquatic organisms in Sweden as indicators of mercury contamination in the environment. Oikos 1967; 18: 323-333.
 36. Lacerda LD, HHM Paraquetti, RV Marins, CE Rezende, IR Zalmon, MP Gomes, V Faria. Mercury content in shark species from the southeastern Brazilian Coast. Brazilian Journal of Biology 2000; 60: 571-6.
 37. Lavoie RA, CE Hebert, J Rail, BM Braune, E Yumvihoze, LG Hill, DRS Lean. Trophic structure and mercury distribution in a Gulf of St. Lawrence (Canada) food web using stable isotope analysis. Science of the Total Environment 2010; 408(22): 5529-5539.
 38. Licata P, D Trombetta, M Cristani, C Naccari, D Martino, M Caló, F Naccari. Heavy metals in liver and muscle of Bluefin Tuna (*Thunnus thynnus*) caught in the straits of Messina (Sicily, Italy). Environmental Monitoring and Assessment 2005; 107(1-3): 239-248.
 39. Liu B, H Yan, C Wang, Q Li, SP Guédron, JE Spangenberg, X Feng, J Dominik. Insights into low fish mercury bioaccumulation in a mercury-contaminated reservoir, Guizhou, China. Environmental Pollution 2012; 160: 109-117.
 40. Loumbourdis NS, G Danscher. Autometallographic tracing of mercury in Frog liver. Environmental Pollution 2004; 129: 299-304.
 41. Lyle JM. Mercury concentrations in four carcharhinid and three hammerhead sharks from the coastal waters of the Northern Territory. Australian Journal of Marine and Freshwater Research 1984; 35: 385 - 397.
 42. Mathers RA, PH Johansen. The effects of feeding ecology on mercury accumulation in walleye, *Stizostedion vitreum*, and pike, *Esox lucius*, in Lake Simcoe. Canadian Journal of Zoology 1985; 63: 2006-2012.
 43. Merritt KA, A Amirbahman. Mercury methylation dynamics in estuarine and coastal marine environments-a critical review. Earth-Science Reviews 2009; 96(1-2): 54-66.
 44. Mol JH, SR Joyce, C Lietar, M Verloo. Mercury contamination in freshwater, estuarine, and marine fishes in relation to small-scale gold mining in suriname. Environmental Research 2001; 86: 183-197.
 45. Monteiro LR, EJ Isidro, HD Lopes. Mercury content in relation to sex, size, age and growth in two scorpionfish (*Helicolenus dactylopterus* and *Pontinus kuhlii*) from Azorean waters. Water, Air, and Soil Pollution 1991; 56(1): 359-367.
 46. Moreno T, P Higuera, T Jones, L McDonald, W Gibbons. Size fractionation in mercury-bearing airborne particles (HgPM10) at Almadén, Spain: Implications for inhalation hazards around old mines. Atmospheric Environment 2005; 39(34): 6409-6419.
 47. Negrete JM, JO Verbal, EL Ceballos, LN Benitez. Total mercury and methylmercury concentrations in fish from the Mojana region of Colombia.

- Environmental Geochemistry and Health 2008; 30: 21-30.
48. Nfon E, IT Cousins, O Järvinen, AB Mukherjee, M Verta, D Broman. Trophodynamics of mercury and other trace elements in a pelagic food chain from the Baltic Sea. *Science of the Total Environment* 2009; 407(24): 6267-6274.
 49. Palmisano F, N Cardellicchio, PG Zambonin. Speciation of mercury in dolphin liver: A two-stage mechanism for the demethylation accumulation process and role of selenium. *Marine Environmental Research* 1995; 40: 109-121.
 50. Pethybridge H, D Cossa, ECV Butler. Mercury in 16 demersal sharks from southeast Australia: Biotic and abiotic sources of variation and consumer health implications. *Marine Environmental Research* 2010; 69: 18-26.
 51. Pinho AP, JRD Guimara, AS Martins, PAS Costa, G Olavo, J Valentin. Total mercury in muscle tissue of five shark species from Brazilian offshore waters: effects of feeding habit, sex, and length. *Environmental Research* 2002; 89(3): 250-258.
 52. Rani L, B Basnet, A Kumar. Mercury Toxicity. In: *Encyclopedia of Environmental Health* (O.N. Jerome ed), 2011; pp. 705-712. Elsevier, Burlington.
 53. Salini JP, SJ Blaber, DT Brewer. Diets of sharks from estuaries and adjacent waters of the north-eastern Gulf of Carpentaria, Australia. *Australian Journal of Marine and Freshwater Research* 1992; 43(1): 87-96.
 54. Salini JP, SJM Blaber, DT Brewer. Diets of trawled predatory fish of the Gulf of Carpentaria, Australia, with particular reference to predation on prawns. *Australian Journal of Marine and Freshwater Research* 1994; 45(3): 397-411.
 55. Saniewska D, M Beldowska, J Beldowski, M Saniewski, J Kwasniak, L Falkowska. Distribution of mercury in different environmental compartments in the aquatic ecosystem of the coastal zone of the Southern Baltic Sea. *Journal of Environmental Sciences* 2010; 22: 1144-1150.
 56. Sfezer P, M Domagala-Wieloszewska, J Warzocha, A Garbacik-Wesolowska, T Ciesielski. Distribution and relationships of mercury, lead, cadmium, copper and zinc in perch (*Perca fluviatilis*) from the Pomeranian Bay and Szczecin lagoon, southern Baltic. *Food Chemistry* 2003; 81: 73-83.
 57. Solis MT, E Yuen, PS Cortez, PJ Goebel. Family poisoned by mercury vapor inhalation. *American Journal of Emergency Medicine* 2000; 18: 599-602.
 58. Stern GA, RW Macdonald, PM Outridge, S Wilson, J Chélat, A Cole, H Hintelmann, LL Loseto, A Steffen, F Wang, C Zdanowicz. How does climate change influence arctic mercury? *Science of the Total Environment* 2012; 414: 22-42.
 59. Storelli MM, R Giacomini-Stuffer, GO Marcotrigiano. Mercury accumulation and speciation in muscle tissue of different species of sharks from the Mediterranean Sea, Italy. *Bulletin of Environmental Contamination and Toxicology* 2002; 68: 201-210.
 60. Taguchi M, K Yasuda, S Toda, M Shimizu. Study of metal contents of elasmobranch fishes: part 1-metal concentration in the muscle tissue of dogfish *Squalus mitsukuri*. *Marine Environmental Research* 1979; 2: 239-249.
 61. Tessier E, R Martin-Doimeadios, D Amouroux, A Morin, C Lehnhoff, E Thybaud, E Vindimian, O Donard. Time course transformations and fate of mercury in aquatic model ecosystems. *Water, Air, and Soil Pollution* 2007; 183: 265-281.
 62. Walker T. Mercury concentrations in edible tissues of elasmobranchs, teleosts, crustaceans and molluscs from south-eastern Australian waters. *Australian Journal of Marine and Freshwater Research* 1988; 39: 39-49.
 63. Wang W. Interactions of trace metals and different marine food chains. *Marine Ecology-Progress Series* 2002; 243: 295-309.
 64. White WT. A redescription of *Carcharhinus dussumieri* and *C. sealei*, with resurrection of *C. coatesi* and *C. tjtjot* as valid species (*Chondrichthyes: Carcharhinidae*). *Zootaxa* 2012; 3241: 1-34.
 65. WHO. Evaluation of certain food additives and contaminants. Thirtieth third report of the joint FAO/WHO expert committee on food additives, World Health Organization (WHO, Technical Report Series 776), Geneva, 1989; 80 p.
 66. WHO. Evaluation of certain food additives and contaminants. WHO technical report series, World Health Organization, Geneva, 2007; 225 p.

12/25/2015

It would be a best Wish for any Author to put a publication to be available in Libraries across the world;

Put your Advertisements and/ or Call for Paper to Biomedicine & Nursing by email to nbmeditor@gmail.com , please.

The Biomedicine & Nursing is available in libraries across the world!