

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED

Doc ID: 560251

Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

JAN 18 2019

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO NON-TARGET INVERTEBRATES,
MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS
UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY
THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 6 INSECTICIDE



Agri-Mek[®] SC

Miticide/Insecticide

syngenta[®]

Active Ingredient:

Abamectin¹: 8.0%*

Other Ingredients: 92.0%

Total: 100.0%

¹CAS No. 71751-41-2

*Agri-Mek[®] SC Miticide/Insecticide is formulated as a suspension concentrate and contains 0.7 lb abamectin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1351 EPA Est. 100-NE-001

Product of China
Formulated in USA

SCP 1351A-L1K 1217
4096304

1 quart
(32 fluid ounces)
Net Contents



FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
<p>Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).</p> <p>If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
HOT LINE NUMBER	
<p>For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed or inhaled. Do not breathe vapor or spray mist. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride \geq 14 mils, or Viton® \geq 14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and wildlife.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment wash water or rinsate. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in/or adjacent to the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks to months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of abamectin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers are allowed in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short pants and short-sleeved shirt
- Chemical-resistant gloves made of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride \geq 14 mils, or Viton[®] \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, AND/OR ILLEGAL RESIDUES.

USE INFORMATION

Agri-Mek SC is an aqueous-based soluble concentrate that will control specified pests on the crops listed on this label when the product is applied according to the **Directions for Use**. Thorough coverage of foliage is essential for good mite and insect control.

NOTE:

- **Adjuvant Requirement:** To avoid illegal crop residues, Agri-Mek SC **must always** be mixed with a non-phytotoxic, non-ionic activator type wetting, spreading and/or penetrating spray adjuvant or horticultural oil (not a dormant oil) as specified in the **Directions for Use** for each crop on this label. Non-ionic activator type wetting, spreading and/or penetrating spray adjuvants include non-ionic surfactants (NIS) with at least 75% surface active agent

and crop oil concentrates (COC), vegetable oil concentrates (VOC), methylated seed/vegetable oils (MSO) and organosilicones (OS) with at least 15% emulsifiers/surfactants and include blends of these non-ionic activator type spray adjuvants. Spray adjuvants must be compatible with Agri-Mek SC and must be used at concentrations specified on the **spray adjuvant product label** directions for use for the targeted crop unless more specific directions are provided in the **Directions for Use** for individual crops on this label. **Do not use binder or sticker type adjuvants because these type adjuvants may reduce translaminar movement of the active ingredient into the plant.** SYNGENTA recommends the use of a Chemical Producers and Distributors Association-certified spray adjuvant.

- **Phytotoxicity and Tank Mix Compatibility:** Agri-Mek SC has been tested for phytotoxicity and has a wide margin of safety on a variety of crops. Agri-Mek SC has also been tested and shown to have broad physical, biological, and non-phytotoxic compatibility with many commonly used pesticides, spray adjuvants, and nutritional products. However, since it is not possible to test all potential mixtures, it is recommended that the user contact the supplier of all desired tank mix products for advice and/or pre-test, prior to commercial application, any previously unproven proposed mixtures with Agri-Mek SC. Conduct a jar test for physical compatibility of all components of the proposed mixture using proper concentrations of each mixture component. Pre-test the proposed mixture on a small area of the target crop under environmental and application conditions similar to the intended commercial application to ensure physical, biological, and non-phytotoxic compatibility. **Agricultural products formulated with binder or sticker type components may reduce performance of Agri-Mek SC.**
- **Adjuvant Phytotoxicity Precaution:** Since Agri-Mek SC must always be mixed with a spray adjuvant as instructed in the directions above, and spray adjuvants alone are known to cause phytotoxicity to certain crops under certain environmental conditions, do not use Agri-Mek SC on a spray-adjuvant-sensitive crop unless the spray adjuvant supplier can confirm a known non-phytotoxic labeled use rate for the intended spray adjuvant on the target crop.
- **New York State Restriction:** Do not apply Agri-Mek SC with aircraft in New York State.
- **Seed Treatment:** Unless specifically restricted in the individual **Crop Use Directions** on this label, Agri-Mek SC may be used in the same year abamectin is used as an at-plant seed treatment.

CHEMIGATION – Tuberos and Corm (suppression of thrips)

Sprinkler Irrigation Application Only

Apply Agri-Mek SC at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, rates, and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with Agri-Mek SC applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Agri-Mek SC into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Agri-Mek SC for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

Do not apply Agri-Mek SC through an irrigation system connected to a public water system unless the pesticide label prescribed safety devices for public water systems are in place. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions - Sprinkler Irrigation Applications

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and are capable of being fitted with a system interlock.
- Any alternatives to the above required safety devices must conform to the list of EPA-approved alternative devices.
- Do not apply when wind speed favors drift beyond the area intended for treatment or non-uniform distribution of treated water.
- Do not apply through chemigation systems connected to public water systems.

SPRAY DRIFT

Responsibility

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions.

NOTE: When states have more stringent regulations, they must be observed.

Spray Drift Precautions for Application with Aircraft or Ground Application Equipment

- Apply Agri-Mek SC only when wind velocity favors on-target product deposition (approximately 3 to 10 mph).
- Do not apply with ground application equipment within 25 ft of or with aircraft within 150 ft of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries, or commercial fish farm ponds.
- Do not cultivate within 25 ft of the aquatic area to allow growth of a vegetative filter strip.
- Do not allow this product to drift onto non-target areas. Drift may result in illegal residues or injury to non-target species. Risk of exposure to sensitive areas can be reduced by applying this product when the wind direction is away from the sensitive area.
- Do not apply when the weather conditions may cause drift.
- Avoid application when the temperature is high and/or the humidity is low. These conditions increase the evaporation of spray droplets and the likelihood of drift to aquatic areas.
- Do not apply when wind speed or wind gusts are greater than 15 mph.
- Do not apply when wind speed is below 2 mph because wind direction will vary and there is a high potential for inversion.
- Observe the following precautions when using ground application to spray tree crops or hops in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish ponds:
 - Do not apply Agri-Mek SC when weather conditions favor drift to aquatic areas.
 - Do not apply within 110 ft upwind of aquatic areas or when wind speed is above 8 mph.
 - Spray last 3 rows windward of aquatic areas using nozzles on one side only, with spray directed away from the aquatic areas.
 - Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

Spray Drift Precautions for Aerial Application

Drift Management Requirements

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops.

- **Outermost Nozzle Distance**

The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.

- **Nozzle Direction**

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

- **Maximum Wind Speed**

Do not apply when wind speed is greater than 15 mph.

- **Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions. (See **Wind, Temperature and Humidity**, and **Temperature Inversions**.)

- **Controlling Droplet Size**

- Volume**

- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- Pressure**

- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- Number of Nozzles**

- Use the minimum number of nozzles that provide uniform coverage.

- Nozzle Orientation**

- Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

- Nozzle Type**

- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

- **Boom Length**

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

- **Application Height**

Applications should not be made at a height greater than 10 ft above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

- **Swath Adjustment**

When applications are made with a cross wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind.

- **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

- **Temperature and Humidity**

To compensate for evaporation when applying Agri-Mek SC in low relative humidity, set up equipment to produce larger droplets. Evaporation of droplets is most severe when conditions are both hot and dry.

- **Temperature Inversions**

Agri-Mek SC must not be applied during a temperature inversion because the potential for drift is high. Temperature inversions restrict vertical air mixing, and this causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds that are common during inversions. Temperature inversions are characterized by temperatures that increase with altitude and are common on nights with limited cloud cover and light to no wind. Inversions begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Resistance Management

GROUP	6	INSECTICIDE
--------------	----------	--------------------

Some insect or mite pests are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects or mites on this label.

Agri-Mek SC contains a Group 6 miticide/insecticide (abamectin). Insect or mite biotypes with acquired or inherent resistance to Group 6 miticides/insecticides may eventually dominate the pest population if Group 6 miticides/insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Agri-Mek SC or other Group 6 miticides/insecticides.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect or mite may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 6 miticides/insecticides exclusively for season long control of insect or mite species with more than one generation per crop season.
- For insect or mite species with successive or overlapping generations, apply Agri-Mek SC or other Group 6 miticides/insecticides using a “treatment window” approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated) of the Group 6 miticides/insecticides. Do not exceed the maximum Agri-Mek SC allowed per year.
- Following a treatment window of Group 6 miticides/insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 6 miticides/insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest’s ability to develop resistance to these classes of chemistry.
- If resistance is suspected, do not reapply Agri-Mek SC or other Group 6 miticides/insecticides.

Other Insect or Mite Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect or mite control program.
- Monitoring treated insect or mite populations for loss of field efficacy.
- Using tank-mixtures or premixes with miticides/insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect or Mite Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

CROP USE DIRECTIONS

AVOCADOS

For control of Avocado thrips and Persea mite

NOTE: Aerial vs. Ground Application: Apply Agri-Mek SC using ground application equipment or aircraft. Aerial application is permitted because of its importance to the avocado-growing industry. However, aerial application is not the preferred method of application for the best control of thrips. With aerial application, the resulting level and duration of control of thrips could be less than with ground application.

Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good mite and insect control.

Avocados

Pest	fl oz/A	Instructions
Avocado Thrips (<i>Scirtothrips perseae</i>) Persea mite (<i>Oligonychus perseae</i>)	2.25-4.25 Do not use less than 2.25 fl oz of product/A	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. 1.0% to 4.0% v/v of a horticultural spray oil (not a dormant oil) approved for use on avocados is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • For best results, apply when immature thrips are first observed but before numbers exceed 5 immature thrips per leaf/fruit. • Use 2.25-3.25 fl oz of product /A for low* to moderate* infestations and 3.25-4.25 fl oz of product/A for severe* infestations. • Make a second application, if needed, to maintain control but wait at least 30 days after the first application to retreat. (See Use Restrictions below.) • For Ground Application: If spray volume is greater than 400 gal/A, apply Agri-Mek SC at a rate of 0.5 fl oz per 100 gallons of water.
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	Minimum 1 to 4% v/v	

continued...

*Thrips Infestation Level	Number of immature thrips per leaf/fruit
Low	1-2
Moderate	3-4
Severe	More than 5

Use Restrictions - Avocados

- **PHI:** Do not apply within **14 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on avocados. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.
 - Note:** To prevent crop injury, observe all precautions and restrictions on the horticultural spray oil label. When using horticultural spray oil concentrations above 2.0%, treat a small test area before making a large-scale application.
- **Application Method:** Ground and aerial application are permitted.
- **Amount of Water:** Do not apply in less than 100 gallons of water with ground application equipment or 50 gallons of water with aircraft.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 30 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated orchards.

BEAN, DRY and SUCCULENT

Crops: *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, blackeyed pea, cowpea*, crowder pea, mung bean, and southern pea)

*For use on cowpeas that are grown only for dry seed. Do not allow livestock to graze cowpea forage and do not harvest cowpea forage or hay for use as livestock feed.

For control of *Liriomyza* leafminers and Spider Mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control, apply Agri-Mek SC with ground application equipment. With aerial application, the resulting control of leafminers and spider mites could be less than with ground application.

Bean

Pest	fl oz/A	Instructions
<i>Liriomyza</i> leafminers Spider mites	1.75 – 3.5	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for high infestations.• Leafminers and Spider Mites: Apply when adult leafminer flies or spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions.)

Use Restrictions – Bean

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Bean crop list above. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 10 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb. ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 6 days before repeating application.
- **Number of Applications:** Do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

CANEBERRY (CROP SUBGROUP 13-07A)

Pest	fl oz/A	Instructions
Spider mites	1.75-3.5	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations.• Apply when mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. However, do not apply more than the maximum amount per application or maximum amount per year. (See Use Restrictions).
Broad mite	3.5	

Use Restrictions – Caneberry (Crop Subgroup 13-07A)

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on caneberry. Carefully follow the **Directions for Use** and **Precautions** on the adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 10 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- Application is not permitted from onset of flowering until petal fall is complete.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** Do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

CELERIAC (*Apium graveolens*)

For control of Twospotted spider mite

Apply Agri-Mek SC using conventional ground application sprayers calibrated to deliver sufficient water for thorough coverage. Thorough coverage of the crop canopy is essential for optimum results. Inadequate crop coverage can result in reduced control.

Celeriac

Pest	fl oz/A	Instructions
Twospotted spider mite	3.5	<ul style="list-style-type: none">To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)Apply when mites first appear and repeat as necessary to maintain control. (See Use Restrictions below.)

Use Restrictions – Celeriac

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on celeriac. Carefully follow the **Directions for Use** and **Precautions** on the adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Apply with ground application equipment only.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** After transplanting the celeriac, do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

CITRUS FRUIT CROP GROUP 10-10

Crops in this group are: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliolate orange; unqi fruit; cultivars, varieties, and/or hybrids of these.

For control of Asian citrus psyllid, Broad mite, Citrus bud mite, Citrus leafminer, Citrus rust mite, Citrus thrips, and Twospotted spider mite

Thorough spray coverage is essential for good mite and insect control. Adjust gallons of spray per acre based on size and number of trees per acre as well as density of foliage.

• Ground Application

Apply using conventional ground application sprayers calibrated to deliver sufficient water for thorough coverage (or outside coverage for Asian citrus psyllid, citrus leafminer, or citrus thrips).

• Aerial Application

Aerial application is permitted **only** for control of citrus leafminer and Asian citrus psyllid. When using aerial application, the resulting level and duration of control of these pests could be reduced compared to ground application. It is recommended to use the higher end of the rate range (3.75 to 4.25 fl oz per acre for concentrated sprays) when applying this product by air.

Citrus Fruit Crop Group 10-10

Pest	fl oz/A	Instructions
Asian citrus psyllid	2.25-4.25 ¹	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. It is recommended for best results that a minimum of 0.20% horticultural spray oil (not a dormant oil) be used in the spray mixture or not less than 1.0 gal of horticultural spray oil/A (except when specified differently in footnotes below). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) Asian Citrus Psyllid: Apply Agri-Mek SC to protect newly expanding foliage flush during the spring, summer or fall. Agri-Mek SC will only control adults and nymphs present at the time of application. Use 3.75 to 4.25 fl oz per acre with aerial application. Mites: Apply when mites first appear during spring, summer, and/or fall. Citrus Bud Mite: For best results, time the spray at "bud swell." Citrus Leafminer: Apply to protect new growth during spring, summer, or fall. Use 3.75 to 4.25 fl oz per acre with aerial application. Citrus Thrips: Application targeted for citrus thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun – preferably early to mid-hatch).
Broad mite	2.25-4.25 ¹	
Citrus bud mite	2.25-4.25 ²	
Citrus leafminer	1.0-4.25 ¹	
Citrus rust mite	1.0-4.25 ¹	
Citrus thrips	2.25-4.25 ³	
Twospotted spider mite	2.25-4.25	

¹Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural spray oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 MPH.

²For best results, use a minimum of 500 gal/A of spray mix with a minimum of 0.5% horticultural spray oil (not a dormant oil).

³For best results, use 100-250 gal/A of spray mix applied to achieve outside coverage.

Use Restrictions – Citrus Fruit Crop Group 10-10

- **PHI:** Do not apply within 7 days of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Citrus Fruit Crop Group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Refer to the **Directions for Use** table above for additional spray adjuvant instructions.
- **Application Method:**
 - **To control citrus leafminer and Asian citrus psyllid** - Ground or aerial application is permitted. For aerial application use a minimum of 10 gallons of finished spray volume per acre. For best results, use at least 25 gallons of finished spray volume per acre. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
 - **All other pests** - Ground application only. Do not apply with aircraft.
- **Prohibited Use:** To manage resistance, do not use in citrus nurseries.
- Application is not permitted from onset of flowering until after petal fall is complete.
- **Application Interval:** Wait at least 30 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:**
 - **Do not** apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
 - **Do not** make more than 3 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated orchards.

CORN (SWEET)

For control of Spider Mites (Tetranychid species including Banks Grass Mite)

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control apply Agri-Mek SC with ground application equipment. With aerial application, the resulting control of spider mites could be less than with ground application.

Corn (Sweet)

Pest	fl oz/A	Instructions
Spider mites (Tetranychid species including Banks Grass Mite)	1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for high infestations. • Spider Mites: Apply when spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

footnotes on the following page...

Use Restrictions – Corn (Sweet)

- **PHI:** Do not apply within **7 days** of harvest (7-day PHI) for forage, ears or stover.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on sweet corn. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 10 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 7.0 fl oz/A (or 0.038 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

COTTON

For control of spider mites

Mix with water as indicated below. Thorough coverage of the crop is essential for optimum results. Inadequate coverage may result in reduced control.

Note:

- Agri-Mek SC may be applied with ground application equipment or aircraft.
- For best control of spider mites, apply Agri-Mek SC with ground application equipment. With aerial application, spray coverage and the resulting level and duration of control of mites may be less than with ground application.

Pest	fl oz/A	Instructions
Carmine spider mite Pacific spider mite Strawberry spider mite Twospotted spider mite	1.0-1.25 (Early-Season Cotton - See Instructions) 1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Apply when mites first appear. The use rate depends on the size of the plant and density of the foliage. • The lower use rates of 1.0-1.25 fl oz/A are only recommended if cotton is early season and under 10 inches in height. Do not use less than 1.0 fl oz/A. • West of the Rocky Mountains - the lower use rates may only be used on cotton less than 10 inches in height and applied only with ground equipment. • Repeat application, if needed, to maintain control. See Use Restrictions below.

Use Restrictions

- **PHI:** Do not apply within **20 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a surfactant approved for use on cotton. Carefully follow the **Directions for Use** and **Precautions** on the surfactant label as well as the precautions in official spray guides or on product labels.
- **Application Method:** Ground or aerial application is permitted.
- **Minimum Amount of Water:** Do not apply in fewer than 5 gallons of water per acre. Use sufficient volume to ensure thorough coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application interval:** Wait at least 21 days before repeating application.
- **Maximum Amount per Year:** Do not apply more than 7.0 fl oz/A (or 0.038 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing Restriction:** Do not feed or allow livestock to graze treated cotton.

CUCURBIT VEGETABLES CROP GROUP 9

Crops in this group are: Chayote, Chinese waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourd (includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp., (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo*, includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter Squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), and Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*).

For Control of Leafminers and Spider Mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control of leafminers and spider mites, apply Agri-Mek SC with ground application equipment. With aerial application, the resulting level and duration of control of leafminers and spider mites could be less than with ground application.

Cucurbit Vegetables Crop Group 9

Pest	fl oz/A	Instructions
Leafminers Spider mites	1.75-3.5	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 1.75-2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations.• Apply when adult leafminer flies or spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

footnotes on the following page...

Use Restrictions – Cucurbit Vegetables Crop Group 9

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Cucurbit Vegetables Crop Group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

FRUITING VEGETABLES CROP GROUP 8-10 (EXCEPT CUCURBITS)

Crops in this group are: African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; nonbell pepper; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these

Also for use in Commercially Grown Greenhouse Tomatoes

For control of Broad mite, Colorado potato beetle, *Liriomyza* leafminers, Spider mites, *Thrips palmi*, Tomato pinworm, Tomato psyllid, and Tomato russet mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control of mites and insects, apply Agri-Mek SC with ground application equipment. With aerial application, the resulting level and duration of control of mites and insects could be less than with ground application.

Fruiting Vegetables Crop Group 8-10

Pest	fl oz/A	Instructions
Broad mite Colorado potato beetle <i>Liriomyza</i> leafminers Spider mites Thrips palmi Tomato psyllid Tomato russet mite	1.75-3.5	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) Broad, russet, and spider mites: Apply when mites first appear. Thrips palmi: Apply when thrips are first observed. Tomato pinworm: Application can be made from the time moth activity is detected up to, but no later than, the time when newly emerged larvae are present. Mites, leafminers, Thrips palmi, and Colorado potato beetle: Use 1.75 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations. Repeat application as necessary to maintain control. (See Use Restrictions below.)
Tomato pinworm	3.5	

Use Restrictions – Fruiting Vegetables Crop Group 8-10

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Fruiting Vegetables Crop Group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Prohibited Use:** For resistance management, do not use on fruiting vegetables grown for transplanting.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

Commercially Grown Greenhouse Tomatoes Only

Pest	fl oz/A	Instructions
<i>Liriomyza</i> leafminers Spider mites <i>Thrips palmi</i> Tomato psyllid Tomato russet mite	1.75-3.5	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and <i>Use Restrictions</i> below.) Russet and spider mites: Apply when mites first appear. <i>Thrips palmi</i>: Apply when thrips are first observed. Mites, leafminers, and <i>Thrips palmi</i>: Use 1.75 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations. Repeat application as necessary to maintain control. (See Use Restrictions.)
Tomato pinworm	3.5	

Use Restrictions – Commercially Grown Greenhouse Tomatoes Only

- **PHI:** Do not apply within 1 day of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on greenhouse-grown tomatoes. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only
- **Prohibited Use:** Not for use in nurseries, plant propagation houses, or greenhouses by commercial transplant producers of plants being grown for transplanting
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Season:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per season.
- **Not for use in New York State.**

GRAPES AND SMALL FRUIT VINE CLIMBING SUBGROUP (EXCEPT FUZZY KIWIFRUIT) CROP SUBGROUP 13-07F

Crops in this group are: Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these

For control of Pacific spider mite, Twospotted spider mite, Variegated leafhopper, Western grape leafhopper, Western grapeleaf skeletonizer, and Willamette spider mite

Apply Agri-Mek SC using conventional ground sprayers calibrated to deliver sufficient water for thorough coverage. Thorough coverage is essential for good spider mite and insect control. Do not spray alternate rows. Agri-Mek SC must be applied to both sides of each row for maximum coverage.

Grapes, Crop Subgroup 13-07F

Pest	fl oz/A	Instructions
Pacific spider mite Twospotted spider mite Variegated leafhopper Western grape leafhopper Western grapeleaf skeletonizer Willamette spider mite	1.75-3.5	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) Use 1.75-2.5 fl oz/A for low to moderate infestations and 3.5 fl oz for severe infestations. Repeat application, if needed. (See Use Restrictions below.) Spider mites: Apply when mites first appear but before motiles exceed 5 per leaf. Western grapeleaf skeletonizer: Apply when larvae are first observed. For optimum control, apply shortly after egg hatch. Leafhoppers: Agri-Mek SC provides contact knock-down only.

Use Restrictions – Grapes, Crop Subgroup 13-07F

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the targeted grape variety. Although Agri-Mek SC has been tested in combination with a range of non-ionic activator type wetting, spreading and/or penetrating spray adjuvants for safety to grapes, it is impossible to test Agri-Mek SC with all potential spray adjuvants on all grape varieties under the variety of conditions that could cause crop injury. Therefore, when using Agri-Mek SC, carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment.
- When using an electro-static sprayer, less than 50 gallons of water/A can be used; however, do not use less than 5 gallons of water/A.
- Application is not permitted from onset of flowering until petal fall is complete.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 7.0 fl oz/A (0.038 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated vineyards.

GUAVA

Crops in this group are: guava, feijoa, jaboticaba, wax jambu, starfruit, passionfruit, acerola

For control of Twospotted spider mite

Thorough spray coverage is essential for good mite and insect control. Spray gallonage may vary depending on size and number of trees per acre as well as density of foliage.

Ground Application: Apply using conventional ground application sprayers calibrated to deliver sufficient water for thorough coverage.

Guava

Pest	fl oz/A	Instructions
Twospotted spider mite	2.25 – 4.25 ¹	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. It is recommended for best results that a minimum of 0.20% horticultural spray oil (not a dormant oil) be used in the spray mixture or not less than 1.0 gallon of horticultural spray oil/A (except when specified differently in footnotes below). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 2.25-3.5 fl oz of product/A for low to moderate infestations and 4.25 fl oz of product/A for severe infestations.• Mites: Apply when mites first appear during spring, summer, and/or fall.

¹Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 MPH.

Use Restrictions – Guava

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Guava crop list above. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Refer to the **Directions for Use** table above for additional spray adjuvant instructions.
- **Application Method:** Ground application only.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gallons of water/A may be used; however, do not use less than 5 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.

- **Application Interval:** If a second application is necessary, wait at least 14 days before repeating application.
- **Number of Applications:** Do not make more than 3 applications of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 12.75 fl oz/A (0.07 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

HERB CROP SUBGROUP 19A

Crops in this subgroup are: Angelica, Balm, Basil, Borage, Burnet, Camomile, Catnip, Chervil (dried), Chives, Chives (Chinese), Clary, Coriander (leaf), Cilantro (leaf), Costmary, Culantro (leaf), Curry (leaf), Dill weed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram (*Origanum* species), Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Summer and Winter savory, Sweet bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, and Wormwood.

For control of *Liriomyza* leafminers and Spider Mites

Mix with water as indicated below. Apply thoroughly with ground application equipment as a foliar spray and ensure good upper and lower leaf coverage.

Herb Crop Subgroup 19A

Pest	fl oz/A	Instructions
<i>Liriomyza</i> leafminers Spider Mites	1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 1.75 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations. • Apply when adult leafminer flies or spider mites are first observed and repeat application as necessary to maintain control. (See Use Restrictions below.)

Use Restrictions – Herb Crop Subgroup 19A

- **PHI:** Do not apply within **14 days** of harvest. For **chives**, do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Herb Crop Subgroup 19A. Although Agri-Mek SC has been tested in combination with a range of non-ionic activator type wetting, spreading and/or penetrating spray adjuvants for safety to herbs, it is impossible to test Agri-Mek SC with all potential spray adjuvants on all herb types and varieties under the variety of conditions that could cause crop injury. Therefore, it is important to carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Apply with ground application equipment.
- **Amount of Water:** Use a minimum of 20 gallons of water/A.
- Application is not permitted from onset of flowering until petal fall is complete.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.

continued...

- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per single cutting (harvest).
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

HOPS (Not approved for use in California)

For control of Twospotted spider mite

Mix with water as indicated below. Apply with ground application equipment as a foliar spray and ensure good upper and lower leaf coverage. Adjust gallons of spray per acre based on the size of plants and density of foliage. In any case, thorough coverage is essential for good control of twospotted spider mite. When using concentrate sprays, the amount of water is reduced, but the amount of Agri-Mek SC per acre must remain the same as for dilute sprays.

Hops

Pest	fl oz/A	Instructions
Twospotted spider mite	1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Apply Agri-Mek SC when twospotted spider mites reach treatment thresholds. • For applications at 1/2 trellis growth (6-8 ft height), apply 1.75-3.5 fl oz/A in a minimum of 40 gallons of water/A. • For applications beyond 1/2 trellis growth, do not use less than 3.5 fl oz/A in a minimum of 100 gallons of water/A.

Use Restrictions – Hops

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on hops. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Application Interval:** If a second application is necessary to maintain control, wait at least 21 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product and repeat application of Agri-Mek SC or any other foliar-applied abamectin-containing product only after an alternative miticide with a different mode of action has been used.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Maximum Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 7.0 fl oz/A (or 0.038 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated hops yards.

LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLES) CROP GROUP 4

Crops in this group are: Amaranth (leafy Amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celtuce; Chervil; Chinese celery; Chrysanthemum, edible leaved and garland; Corn salad; Cress, garden and upland (yellow rocket and winter cress); Dandelion; Dock (Sorrel); Endive (escarole); Florence fennel; Lettuce, head and leaf; New Zealand Spinach; Orach; Parsley; Purslane, garden and winter; Radicchio (red chicory); Rhubarb; Spinach; Vine spinach; and Swiss chard.

For Control of Carmine spider mite, *Liriomyza* leafminers, and Twospotted spider mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control of leafminers and spider mites, apply Agri-Mek SC with ground application equipment. With aerial application, the resulting level and duration of control of leafminers and spider mites could be less than with ground application.

Leafy Vegetables Crop Group 4

Pest	fl oz/A	Instructions
Carmine spider mite <i>Liriomyza</i> leafminers Twospotted spider mite	1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 1.75-2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations. • Spider mites: Apply when mites first appear, and repeat as needed to maintain control. (See Use Restrictions below.) • Leafminers: Apply when adult flies are first observed and repeat applications as needed to maintain control. (See Use Restrictions below.)

Use Restrictions – Leafy Vegetables Crop Group 4

- **PHI:** Do not apply within 7 days of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Leafy Vegetables Crop Group. Although Agri-Mek SC has been tested in combination with a range of non-ionic activator type wetting, spreading, and/or penetrating spray adjuvants for safety to leafy vegetables, it is impossible to test Agri-Mek SC with all potential spray adjuvants on all leafy vegetable types and varieties under the variety of conditions that could cause crop injury. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.

continued...

- **Prohibited Use:** For resistance management, do not use on leafy vegetables grown for trans-planting.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

LYCHEE

Crops in this group are: lychee, longan, Spanish lime, rambutan, pulasan

For control of Twospotted spider mites, *Liriomyza* leafminers, and Thrips

Thorough spray coverage is essential for good mite and insect control. Spray gallonage may vary depending on size and number of trees per acre as well as density of foliage.

Ground Application: Apply using conventional ground application sprayers calibrated to deliver sufficient water for thorough coverage.

Pest	fl oz/A	Instructions
Twospotted spider mite <i>Liriomyza</i> leafminers Thrips	2.25 – 4.25 ¹	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. It is recommended for best results that a minimum of 0.20% horticultural spray oil (not a dormant oil) be used in the spray mixture or not less than 1.0 gal of horticultural spray oil/A (except when specified differently in footnotes below). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 2.25-3.5 fl oz of product /A for low to moderate infestations and 4.25 fl oz of product/A for severe infestations. • Mites: Apply when mites first appear during spring, summer, and/or fall. • Leafminers: Apply to protect new growth during spring, summer, or fall. <p style="text-align: right;"><i>continued...</i></p>

Pest	fl oz/A	Instructions
Twospotted spider mite <i>Liriomyza</i> leafminers Thrips (continued)	2.25 – 4.25 ¹	<ul style="list-style-type: none"> • Thrips: Applications targeted for thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun – preferably early to mid-hatch).

¹Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 MPH.

Use Restrictions – Lychee

- **PHI:** Do not apply within **14 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Lychee crop list above. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Refer to the **Directions for Use** table above for additional spray adjuvant instructions.
- **Application Method:** Ground application only.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gallons of water/A may be used; however, do not use less than 5 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 30 days before repeating application.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (0.046 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

MINT (PEPPERMINT AND SPEARMINT)

For control of Twospotted spider mite

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control of spider mites, apply Agri-Mek SC with ground application equipment. With aerial application, the resulting level and duration of control of spider mites could be less than with ground application.

Mint

Pest	fl oz/A	Instructions
Twospotted spider mite	1.75-2.5	<ul style="list-style-type: none">To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)Treat when spider mites first appear.

Use Restrictions – Mint

- **PHI:** Do not apply within **28 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on mint. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions such as high temperatures, use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 2.5 fl oz/A (or 0.014 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product and do not apply more than 3 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 7.75 fl oz/A (or 0.042 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per crop per year.
- **Grazing:** Do not allow livestock to graze or feed treated foliage to livestock.

ONION, BULB CROP SUBGROUP 3-07A

Crops in this group are: onion, bulb including daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.

For control of *Liriomyza* Leafminers and Thrips

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control apply Agri-Mek SC with ground application equipment. With aerial application, the resulting control of leafminers and thrips could be less than with ground application.

Onion, Bulb Crop Subgroup 3-07A

Pest	fl oz/A.	Instructions
<i>Liriomyza</i> leafminers Thrips	1.75-3.5	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for high infestations. Insect control can be reduced if Agri-Mek SC is used in combination with a sticker or binder type product such as Bravo® Weather Stik®. Leafminers: Apply when adult leafminer flies are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.) Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.) Do not use Agri-Mek SC as a rescue treatment for thrips control.

Use Restrictions – Onion, Bulb Crop Subgroup 3-07A

- **PHI:** Do not apply within 30 days of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on bulb onions or on the other intended target crop in the Bulb Crop Subgroup 3-07A. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** Make 2 consecutive applications of Agri-Mek SC, then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional Agri-Mek SC applications. Do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

ONION, GREEN CROP SUBGROUP 3-07B

Crops in this group are: chive, fresh leaves; chive, Chinese, fresh leaves; *Elegans* hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

For control of *Liriomyza* leafminers and Thrips

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control apply Agri-Mek SC with ground application equipment. With aerial application, the resulting control of leafminers and thrips could be less than with ground application.

Onion, Green Crop Subgroup 3-07B

Pest	fl oz/A	Instructions
<i>Liriomyza</i> leafminers Thrips	1.75 – 3.5	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for high infestations.• Insect control can be reduced if Agri-Mek SC is used in combination with a sticker or binder type product such as Bravo® Weather Stik®.• Leafminers: Apply when adult leafminer flies are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions.)• Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions.) Do not use Agri-Mek SC as a rescue treatment for thrips control.

Use Restrictions – Onion, Green Crop Subgroup 3-07B

- **PHI:** Do not apply within **7 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the Use Information section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on green onions or on the other intended target crop in the Bulb Crop Subgroup 3-07B. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.

- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State or CA.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** Make 2 consecutive applications of Agri-Mek SC then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional Agri-Mek SC applications. Do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 14 fl oz/A (or 0.076 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

PAPAYA

Crops in this group are: papaya, star apple, black sapote, sapodilla, canistel, mamey sapote

For control of Twospotted spider mites

Pest	fl oz/A	Instructions
Twospotted spider mite	2.25 -4.25 ¹	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. It is recommended for best results that a minimum of 0.20% horticultural spray oil (not a dormant oil) be used in the spray mixture or not less than 1.0 gallon of horticultural spray oil/A (except when specified differently in footnotes below). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 2.25-3.5 fl oz of product /A for low to moderate infestations and 4.25 fl oz of product/A for severe infestations. • Mites: Apply when mites first appear during spring, summer, and/or fall.

¹Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 MPH.

continued...

Use Restrictions – Papaya

- Do not apply to mango.
- **PHI: Do not apply within 7 days of harvest.**
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Papaya crop list above. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Refer to the **Directions for Use** table above for additional spray adjuvant instructions.
- **Application Method:** Ground application only.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gallons of water/A may be used; however, do not use less than 5 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 14 days before repeating application.
- **Number of Applications:** Do not make more than 3 applications of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 12.75 fl oz/A (0.070 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

PINEAPPLE

For control of Twospotted spider mite

Pest	fl oz/A	Instructions
Twospotted spider mite	2.25 – 4.25	<ul style="list-style-type: none">• To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.)• Use 2.25-3.5 fl oz/A for low to moderate infestations and 4.25 fl oz/A for high infestations.

Use Restrictions – Pineapple

- **PHI:** Do not apply within **16 weeks** of harvest.
- **Application Method:** Ground application only
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on Pineapple. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gallons of water/A may be used; however, do not use less than 5 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 7 days before repeating application.

- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (0.046 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.

POME FRUIT CROP GROUP 11-10

Crops in this group are: Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

For control of European red mite, McDaniel spider mite, Pear psylla, Pear rust mite, Twospotted spider mite, Yellow mite, Tentiform leafminer, and White apple leafhopper

Apply Agri-Mek SC using conventional ground application sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Pome Fruit Crop Group 11-10

Pest	fl oz/A	Instructions
European red mite McDaniel spider mite Pear psylla Pear rust mite Twospotted spider mite Yellow mite Tentiform leafminer White apple leafhopper	2.25-4.25	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. Horticultural spray oil (not a dormant oil) is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Apply Agri-Mek SC when spider mite or insect thresholds are reached. • Make a second application, if needed, to maintain control. (See Use Restrictions below.)
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	Minimum of 1 gal/A	

Use Restrictions – Pome Fruit Crop Group 11-10

- **PHI:** Do not apply within 28 days of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Pome Fruit Crop Group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.
 - Applying the combination of Agri-Mek SC and horticultural spray oil fewer than 14 days before or after applying Captan® or other sulfur-containing products can result in phytotoxicity and crop loss.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.

continued...

- Application is not permitted from onset of flowering until after petal fall is complete.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated orchards.

SOYBEANS

For control of Spider Mites (Tetranychid species)

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft.
- For best control apply Agri-Mek SC with ground application equipment. With aerial application, the resulting control of spider mites could be less than with ground application.

Soybeans

Pest	fl oz/A	Instructions
Spider mites (Tetranychid species)	1.75 - 3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 1.75 – 2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for high infestations. • Spider Mites: Apply when spider mites are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions below.)

Use Restrictions – Soybeans

- **PHI:** Do not apply within **28 days** of harvest (28-day PHI) for soybean seed.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the Use Information section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on soybeans. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 10 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.

- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
- **Application Interval:** Wait at least 7 days before repeating application.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 7.0 fl oz/A (or 0.038 lb ai/A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per year.
- **Do not** allow livestock to graze in treated areas or harvest treated soybean forage, straw or hay as feed for meat or dairy animals. **Do not** feed treated soybean fodder or silage to meat or dairy animals.

STONE FRUIT CROP GROUP 12-12

Crops in this group are: Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

For control of European red mite, Pacific spider mite, and Twospotted spider mite

Apply Agri-Mek SC using conventional ground sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite control.

Stone Fruit Crop Group 12-12

Pest	fl oz/A	Instructions
European red mite Pacific spider mite Twospotted spider mite	2.25-4.25	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant or with a horticultural spray oil (not a dormant oil). Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Apply when mites first appear. • Make a second application, if needed, to maintain control. (See Use Restrictions below.) • A horticultural spray oil (not a dormant oil) can improve efficacy but it could also increase the possibility of phytotoxicity to foliage and fruit. • Use a minimum of 0.25% horticultural spray oil (not a dormant oil) in the dilute spray mixture and not less than 1.0 gallon of horticultural spray oil per acre in the final finished spray.
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	Minimum of 1 gal/A	

continued...

Use Restrictions – Stone Fruit Crop Group 12-12

- **PHI:** Do not apply within **21 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Stone Fruit Crop Group. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. See the **Directions for Use** table above for application instructions. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- Application is not permitted from onset of flowering until after petal fall is complete.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** If a second application is needed, wait at least 21 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated orchards.

STRAWBERRIES, LOW GROWING BERRY CROP SUBGROUP 13-07G

Crops in this group are: Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these.

For control of Strawberry spider mite, Twospotted spider mite, and Carmine spider mite

Mix with water as indicated below. Apply with ground application equipment as a foliar spray and ensure thorough coverage on upper and lower leaf surfaces. Adjust spray volume and nozzle placement to ensure maximum coverage of tops and undersides of leaves.

Strawberries, Low growing berry Crop Subgroup 13-07G

Pest	fl oz/A	Instructions
Carmine spider mite Cyclamen mite ¹ Strawberry spider mite Twospotted spider mite	3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Make 2 applications 7-10 days apart when mites first appear. Repeat this application sequence, if needed, to maintain control. (See Use Restrictions below.) • Cyclamen mite: Apply in sufficient water to obtain good coverage into the crown of the plant.

¹ Suppression

Use Restrictions – Strawberries

- **PHI:** Do not apply within **3 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a surfactant approved for use on strawberries. Do not use binder or sticker type adjuvants. Carefully follow the **Directions for Use** and **Precautions** on the surfactant label as well as the precautions in official spray guides or on product labels.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Prohibited Use:** For resistance management purposes, do not use in strawberry nurseries.
- **Amount of Water:** Do not apply in less than 50 gallons of water/A with conventional ground application equipment. When using electro-static sprayers do not apply in less than 10 gallons of water/A.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Application Interval:** Wait at least 21 days after the second application before repeating application.
- **Maximum Amount per Year:** Do not apply more than 14 fl oz/A (or 0.075 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.

TREE NUTS CROP GROUP 14-12

Crops in this group are: African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these.

For control of European red mite, Pacific spider mite, Strawberry spider mite, and Twospotted spider mite

Apply Agri-Mek SC using conventional ground sprayers calibrated to deliver sufficient water for thorough coverage. Adjust gallons of spray per acre based on size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Tree Nuts Crop Group 14-12

Pest	fl oz/A	Instructions
European red mite Pacific spider mite Strawberry spider mite Twospotted spider mite	2.25-4.25	<ul style="list-style-type: none"> To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant. Horticultural spray oil (not a dormant oil) is recommended. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) Apply Agri-Mek SC when spider mites first appear. Residual spider mite control is greater from spray deposits on newer leaves compared to older leaves. Make a second application, if needed, to maintain control. (See Use Restrictions below.)
Recommended Rate When Using Horticultural Spray Oil (not a dormant oil)	Minimum of 1 gal/A	

Use Restrictions – Tree Nuts Crop Group 14-12

- **PHI:** Do not apply within **21 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC must be mixed with a spray adjuvant as instructed in the Use Information section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Tree Nuts Crop Group 14-12. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in official local spray guides.
- **Application Method:** Ground application only. Do not apply with aircraft.
- **Amount of Water:** Do not apply in less than 40 gallons of water/A.
- Application is not permitted from onset of flowering until after petal fall is complete.
- **Application Interval:** If a second application is necessary, wait at least 21 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Application:** Do not apply more than 4.25 fl oz/A (or 0.023 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per application.
- **Number of Applications:** Do not make more than 2 applications of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Maximum Amount per Year:** Do not apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per year.
- **Grazing:** Do not allow livestock to graze in treated groves/orchards.

TUBEROUS AND CORM VEGETABLES CROP SUBGROUP 1C

Crops in this subgroup are: arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava (bitter and sweet); chayote (root); chufa; dasheen; ginger; leren; potato; sweet potato; taniel; turmeric; yam bean; and yam (true).

For control of Colorado potato beetle, *Liriomyza* leafminers, Potato psyllid, and Spider mites

Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.

Note:

- Apply Agri-Mek SC using ground application equipment or aircraft.
- Agri-Mek SC may be applied through overhead sprinkler chemigation for suppression of thrips.
- For best control of mites, apply Agri-Mek SC with ground application equipment. With aerial application and chemigation, the resulting level and duration of control of insects and spider mites could be less than with ground application.

Tuberous and Corm Vegetables Crop Subgroup 1C

Pests	fl oz/A	Instructions
Colorado potato beetle <i>Liriomyza</i> leafminers Potato psyllid Spider mites	1.75-3.5	<ul style="list-style-type: none"> • To avoid illegal residues, Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of this Agri-Mek SC label and Use Restrictions below.) • Use 1.75-2.5 fl oz/A for low to moderate infestations and 3.5 fl oz/A for severe infestations.
Thrips (suppression)	3.5	<ul style="list-style-type: none"> • Insect and spider mite control can be reduced if Agri-Mek SC is used in combination with a sticker or binder type product such as Bravo® Weather Stik®. • Colorado potato beetle: Make the first application after approximately 50% of the egg masses have hatched and larvae are present. If 2 applications are needed, limit them to a single Colorado potato beetle generation per crop. (See Use Restrictions below.) • <i>Liriomyza</i> leafminers: Make the first application when adult flies are first observed. Repeat applications as needed to maintain control. (See Use Restrictions below.) • Spider mites: Make the first application when mites first appear. Repeat application as needed to maintain control. (See Use Restrictions below.) • Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program. (See Use Restrictions.) Do not use Agri-Mek SC as a rescue treatment for thrips control.

continued...

Use Restrictions – Tuberos and Corm Vegetables Crop Subgroup 1C

- **PHI:** Do not apply within **14 days** of harvest.
- **Adjuvant Requirement:** To avoid illegal residues, Agri-Mek SC **must** be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant as instructed in the **Use Information** section found at the beginning of this Agri-Mek SC label. The spray adjuvant must be approved for use on the intended target crop in the Tuberos and Corm Vegetables Crop Subgroup. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label and in official local spray guides. See the **Directions for Use** table above for application instructions. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground, aerial, or chemigation application is permitted; however, do not apply with aircraft in New York State.
- **Amount of Water:** Do not apply in less than 20 gallons of water/A with ground application equipment. Do not apply in less than 5 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.
- **Maximum Amount per Application:** Do not apply more than 3.5 fl oz/A (or 0.019 lb ai/A) per application.
- **Application Interval:** Wait at least 7 days before repeating application of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliar-applied abamectin-containing product.
- **Maximum Amount per Year:** Do not apply more than 10.25 fl oz/A (or 0.056 lb ai/A) of Agri-Mek SC or any other foliar-applied abamectin-containing product per crop per year.
- **Grazing:** Do not allow livestock to graze or feed treated foliage to livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a tightly closed container in a cool, dry place.

Pesticide Disposal

Pesticide waste may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control agency, or the Hazardous Waste representative at the nearest EPA regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¹/₄ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

continued...



Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Agri-Mek®, Bravo® Weather Stik®, the ALLIANCE FRAME 
the SYNGENTA Logo, and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

Captan® is a trademark of Tomen Agro, Inc.

Viton® is a trademark of E.I. DuPont De Nemours & Co, Inc.

©2018 Syngenta

For non-emergency (e.g. current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1351A-L1K 1217
4096304