

Dosing for Infants and Children

From Your Healthcare Professional

If possible, use weight to dose; otherwise use age.

Always ask a healthcare provider which product is right for your child.

mL = milliliter
tsp = teaspoonful

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.



Infants' TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL or 1 tsp)

Use only as directed.



Children's TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL or 1 tsp)

Use only as directed.

Available in:
 Grape
 Cherry

WEIGHT	AGE			
6-11 lbs	0-3 mos	1.25 mL		—
12-17 lbs	4-11 mos	2.5 mL		—
18-23 lbs	12-23 mos	3.75 mL		—
24-35 lbs	2-3 yrs	5 mL		5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	<div style="border: 1px solid black; padding: 5px; text-align: center;"> Infants' TYLENOL® is currently not available in stores, but caregivers may have previously purchased bottles at home. </div>	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—		10 mL (2 tsp)
60-71 lbs	9-10 yrs	—		12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—		15 mL (3 tsp)

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products
- Take every 4 hours as needed. Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product
- **All Infants' and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)**



Kids' Wellness Tracker™

Free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN®. Makes it easy to track your child's height, weight, vaccines, symptoms, and medicines plus calculate dosing.



Download now at the App Store™ | App Store is a service mark of Apple Inc. Standard download rates may apply

Dosing for Infants and Children

From Your Healthcare Professional

If possible, use weight to dose; otherwise use age.

Always ask a healthcare provider which product is right for your child.

mL = milliliter
tsp = teaspoonful

DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.



Infants' MOTRIN® Concentrated Drops

Active Ingredient:
Ibuprofen 50 mg (NSAID)[†]
(in each 1.25 mL)

[†]Nonsteroidal anti-inflammatory drug
Use only as directed.

Available in:
dye-free Dye-free
Berry



Children's MOTRIN® Oral Suspension

Active Ingredient:
Ibuprofen 100 mg (NSAID)[†]
(in each 5 mL or 1 tsp)

[†]Nonsteroidal anti-inflammatory drug
Use only as directed.

Available in:
Original Berry
dye-free Dye-free
Berry

WEIGHT	AGE		
6-11 lbs	0-5 mos	Do not use	—
12-17 lbs	6-11 mos	1.25 mL	
18-23 lbs	12-23 mos	1.875 mL	
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID[†] (prescription or non-prescription)
 - MOTRIN® contains ibuprofen
- Use only the dosing device that comes with a specific product

[†]Nonsteroidal anti-inflammatory drug



Kids' Wellness Tracker™

Free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN®. Makes it easy to track your child's height, weight, vaccines, symptoms, and medicines plus calculate dosing.



Download now at the App Store™ | App Store is a service mark of Apple Inc. Standard download rates may apply