



**NICK AVALLONE, M.D.**

[www.dravallone.com](http://www.dravallone.com)

755 Memorial Pkwy  
Suite 201  
Phillipsburg, NJ 08865

22 Walmart Plaza  
2nd Level  
Clinton, NJ 08809  
908-847-8884

## **Rehabilitation Protocol Following Distal Biceps Repair with Biceps Button**

**DISCLAIMER:** The following Physical Therapy protocol is intended to be utilized by the clinician as a guideline in the treatment of this disorder. It is based on current research and has been formulated as a collaborative effort between Physicians and Physical Therapists. It is not intended to serve as a substitute for sound clinical decision making. Every patient is a unique case, and it should be anticipated that not all patients will fit into the timelines set forth in this protocol. If the Physical Therapist has any questions regarding the course of treatment, the referring physician should be contacted for further guidance.

### **PHASE I -POST OPERATIVE PHASE: (0-4 Weeks)**

- **GOALS:**

- Protect healing repair.
- Decrease pain/Edema
- Retard muscular atrophy.
- Begin to restore motion.
- Achieve full motion by 6 weeks.

- **PRECAUTIONS:**

- Caution against lifting any weight or bearing weight through arm.
- Avoid shoulder extension.
- Use of sling/brace per physician instruction.

- **Week 0-2:**

- **Sling/Splint:**

- Use of sling for first 2 weeks or as needed when in community, or use of hinge brace if repair was done under tension per physician instruction.

- **Interventions:**

- Bulky dressing removed.
- Address edema as needed, with compression wrap -hand to above elbow or with tubigrip stockinette.
- Ice may be used if edema is significant.
  - Address wound with wound cleaning and dressing, as needed.

- **Manual Therapy:**

- Myofascial soft tissue mobilization to forearm and biceps musculature to relax biceps and other tight muscles. Avoid incision site.
- **Exercises:**
  - Wrist ROM, Forearm pronation/supination
  - Finger ROM/ tendon glides.
  - Shoulder rolls.
  - Passive flexion and supination allowed as tolerated.
- **Week 2:**
  - **Splint/Sling:**
    - Wean from use of sling unless out in community.
    - If wearing splint/brace will continue per physician's instructions.
  - **Interventions:**
    - Following suture removal begin scar mobilization techniques – scar massage, kinesiotape, etc.
    - Myofascial soft tissue mobilization /IASTM to forearm and biceps musculature to relax biceps and other tight muscles.
    - Continue edema control measures as needed.
  - **Exercises:**
    - Instruction in gentle AROM of the elbow as tolerated or AAROM or table slides as needed.
    - PROM for elbow flexion and supination.
    - Gentle shoulder ROM avoiding excessive extension.
- **Week 3:**
  - **Sling/ Splint:**
    - Continue as above.
  - **Interventions:**
    - Use of heat modalities as needed to relax tight musculature.
    - Scar mobilization and myofascial soft tissue work /IASTM to forearm musculature, biceps/triceps to relax arm into extension.
    - No forceful passive stretching.
  - **Exercises:**
    - AROM activities for elbow, wrist, forearm, light sponge gripping for hand.
    - PROM for elbow flexion and supination.
    - Gentle shoulder ROM avoiding excessive extension.
    -
- **Week 4:**
  - **Sling/ Splint:**
    - D/C use of sling in community.
    - Continue use of brace per physician's instruction
  - **Interventions:**
    - As above, as needed.
  - **Exercises:**

- Gentle passive stretches may be initiated within the patient's comfort level to increase elbow extension.
- Wrist weight exercises initiated 1-3 lbs. Not pronation supination.
- If patient has reached full elbow ROM, therapy will be put on hold until the 6 week time. Patient will work on his HEP.

## **PHASE II- INTERMEDIATE PHASE (6-12 weeks)**

### **▪ GOALS:**

- Achieve full elbow ROM including pronation and supination.
- Adherence to HEP.
- Begin to regain and improve muscular strength.

### **▪ PRECAUTIONS:**

- Avoid lifting heavy weight, except as allowed with exercise program.

### **▪ Week 6:**

- **Sling/Splint:**
  - D/C Brace if was wearing one.
- **Interventions:**
  - Manual Therapy- Myofascial soft tissue work as needed.
- **Exercises:**
  - Wrist weights,-Pronation supination progression with weight.
  - Initiate light theraband program -yellow or red band.
    - Low level shoulder program- Triceps curl, ER, IR, Protraction, gentle biceps curl
  - NuStep or UBE for ROM and progress to strengthening.

### **▪ Weeks 8-12:**

- **Exercises:**
  - Progress to light pulley weights.
  - Progress HEP therabands to green.
  - Progress to light plyometrics, including body blade, therabar and light ball tossing.
  - May initiate Thrower's Ten as appropriate.
  - No Push-ups

### **▪ Weeks 10-12:**

- **Exercises:**
  - Initiate work conditioning program with lifting activities. Advance weight slowly and only with patient comfort.
  - Progress to universal set with bilateral midrows, lower trap pulls, chest press, (10-30lbs)

### **PHASE III- Advanced Strengthening (12-16wks)**

- **GOALS:** ( 3-6 months)
  - Increase strength, power, endurance
  - Maintain full elbow ROM
  - Gradually initiate sporting activities.
  - Improve Sport/work specific balance and proprioceptive skills.
- **PRECAUTIONS:**
  - 12 weeks: General activity as tolerated is permitted.
  - 6 months: Full activity without restriction is allowed. Full Return to Sport per physician
- **Week 12:**
  - **Exercises:**
    - Continue shoulder program- Progress to Advanced Thrower's Ten Program
    - Manual resistance diagonal patterns
    - Weight program.
    - Plyometric exercise program, (2 hand plyos close to body only)
      - Chest Pass, side throw close to body
- **Week 13:**
  - **Exercises:**
    - Continue all above
    - Program plyometrics to 2 hand drills away from body- side to side throws, soccer throws
- **Week 14:**
  - **Exercises:**
    - Continue all above.
    - Progress isotonic machine strengthening ex (if desired), Bench press Seated, Lat pull down
    - Initiate golf, swimming
    - Initiate interval hitting program
    - Initiate one hand plyometric throwing (stationary throws)
    - Initiate one hand wall dribble
    - Initiate one hand baseball throws into wall.

### **PHASE IV- RETURN TO ACTIVITY PHASE: (16+ weeks)**

- **GOALS:**
  - Continue to increase flexibility, endurance and strengthening of upper extremity.
  - Gradual return to sport activities per physician.
- **Week 16:**
  - **Exercises:**
    - Initiate interval throwing program Phase 1 long toss program.

- Continue Advanced thrower's Ten Program and plyometrics
- Continue to stretch before and after throwing.
- **Week 22-24+:**
  - **Exercise:**
    - Progress to Phase 2 throwing (once successfully completed Phase 1)
- **Week 30:**
  - **Exercise:**
    - Once return to sports utilize Thrower's Ten program
    - Continue shoulder and elbow ROM & stretching program
    - Gradually Progress to competitive throwing /sports per physician timeline.

#### **References:**

Rose, D.M., Archibald, J.D., Sutter, E.G. et al. Biomechanical analysis suggests early rehabilitation is possible after single-incision EndoButton distal biceps repair with Fiberwire. *Knee Surg Sports Traumatol Arthrosc* 19,1019-1022 (2011).

Panagopoulos, A., Tatani, I., Tsoumpas, P. et al. Clinical Outcomes and Complications of Cortical Button Distal Biceps Repair: A Systematic Review of the Literature. *Journal of Sports Medicine* Vol 2016 Article ID 3498403 (2016)

Spencer, E.E., Tisdale, A., Kostka, K., et al. Is Therapy Necessary After Distal Biceps Tendon Repair? *Hand* 2008 Dec; 3(4) 316-319.