

# CANCER DISCOVERY



## KDM4A Promotes Site-Specific Copy Number Gain

*Cancer Discovery* 2013;3:965. Published OnlineFirst August 1, 2013.

**Updated version** Access the most recent version of this article at:  
doi:[10.1158/2159-8290.CD-RW2013-162](https://doi.org/10.1158/2159-8290.CD-RW2013-162)

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, contact the AACR Publications Department at [permissions@aacr.org](mailto:permissions@aacr.org).

## Signaling

**Major finding:** Nuclear I $\kappa$ B $\alpha$  binds to histones H2A and H4 and facilitates Polycomb-dependent HOX gene silencing.

**Clinical relevance:** Nuclear I $\kappa$ B $\alpha$  accumulation decreases with squamous cell carcinoma progression.

**Impact:** I $\kappa$ B $\alpha$  has a noncanonical nuclear function that is critical to skin homeostasis and development.

### CHROMATIN-BOUND I $\kappa$ B $\alpha$ MEDIATES POLYCOMB ACTIVITY

The only known activity of I $\kappa$ B proteins is the cytoplasmic sequestration and inhibition of NF- $\kappa$ B. I $\kappa$ B proteins have been found in the nucleus, but their functional significance is unclear. Mulero and colleagues evaluated I $\kappa$ B $\alpha$  localization in human tissues and observed nuclear I $\kappa$ B $\alpha$  staining in the skin, particularly in keratinocytes. Both cytoplasmic and nuclear I $\kappa$ B $\alpha$  could be detected, but phosphorylation and SUMOylation of I $\kappa$ B $\alpha$  were specifically required for its nuclear accumulation. The phosphorylated, SUMOylated nuclear form of I $\kappa$ B $\alpha$  specifically interacted with histones H2A and H4, but not NF- $\kappa$ B, indicating that I $\kappa$ B $\alpha$  can directly bind chromatin in an NF- $\kappa$ B-independent manner. Chromatin immunoprecipitation sequencing in primary human keratinocytes revealed that I $\kappa$ B $\alpha$  binds near promoter and enhancer regions of genes associated with embryonic development and differentiation, including homeobox (HOX) genes. I $\kappa$ B $\alpha$ -occupied loci were also enriched for high levels of the repressive histone H3 lysine 27 trimethylation mark, and overlapped with Polycomb repressive complex (PRC) binding sites. Consistent with these findings pointing to a noncanonical role of I $\kappa$ B $\alpha$  in Polycomb-depend-

ent gene silencing, nuclear I $\kappa$ B $\alpha$  physically interacted with the PRC2 subunit SUZ12, the simultaneous binding of both proteins to I $\kappa$ B $\alpha$  targets was associated with decreased gene expression, and mutation of the *Drosophila* I $\kappa$ B homolog enhanced the Hox-related extra sex combs phenotype of Polycomb mutant flies independently of the *Drosophila* NF- $\kappa$ B homolog. Given that deregulated HOX expression occurs in squamous cell carcinomas, which develop from keratinocytes, the authors assessed I $\kappa$ B $\alpha$  localization in transformed keratinocytes and found that nuclear I $\kappa$ B $\alpha$  accumulation was lost in association with decreased PRC2 binding and massive upregulation of I $\kappa$ B $\alpha$  target genes. Moreover, depletion of nuclear I $\kappa$ B $\alpha$  correlated with squamous cell carcinoma progression. These findings point to a role of nuclear I $\kappa$ B $\alpha$  in developmental gene regulation and suggest that loss of nuclear I $\kappa$ B $\alpha$  may be oncogenic. ■

Mulero MC, Ferres-Marco D, Islam A, Margalef P, Pecoraro M, Toll A, et al. Chromatin-bound I $\kappa$ B $\alpha$  regulates a subset of Polycomb target genes in differentiation and cancer. *Cancer Cell* 2013;24:151–66.

## Genomic Instability

**Major finding:** Overexpression of the H3K9/36me3 demethylase KDM4A induces gain of 1q12, 1q21, and Xq13.1.

**Mechanism:** KDM4A promotes DNA rereplication by displacing HP1 $\gamma$  and recruiting the replication machinery.

**Impact:** The local chromatin state can determine the susceptibility of certain genomic loci to copy gains.

### KDM4A PROMOTES SITE-SPECIFIC COPY NUMBER GAIN

Recurring amplifications of specific genomic loci are observed in human cancers, but the factors that predispose to copy number gains are poorly understood. Given previous findings showing that focal amplification events occur during S phase and that lysine (K)-specific demethylase 4A (KDM4A), an H3K9/36me3 demethylase, regulates DNA accessibility and replication timing, Black and colleagues hypothesized that KDM4A could provide a link between local chromatin structure and copy number variation. Analysis of KDM4A copy number and expression level in The Cancer Genome Atlas revealed that amplification and overexpression of KDM4A occurred in 18.9% of tumors overall and in 46% of ovarian cancers. Stable overexpression of KDM4A did not induce large-scale genomic instability but instead resulted in a specific increase in KDM4A occupancy at chromosome 1q12 and focal amplification of this locus in 14% of cells. Gain of 1q12 was specifically dependent on KDM4A, required KDM4A enzymatic activity, and could be suppressed by overexpression of the H3K9me3 methyltransferase SUV39H1 or the H3K9me3-binding protein heterochromatin protein 1 $\gamma$  (HP1 $\gamma$ ). Interestingly, 1q12 copy



gain was not stably inherited but was generated during S phase of each cell cycle before being eliminated by the end of G2. KDM4A directly bound and recruited the MCM complex and DNA polymerase to 1q12 and induced rereplication of this locus in association with decreased H3K9me3 and HP1 $\gamma$  occupancy, suggesting that KDM4A can induce site-specific copy number gain by altering local chromatin structure and recruiting the DNA replication machinery. In primary tumors, KDM4A amplification was not only correlated with gain of 1q12 but was also correlated with gain of 1q21 and Xq13.1, which were likewise found to be rereplicated in KDM4A-overexpressing cells. Although the reason why these particular loci are amplified upon KDM4A overexpression remains unclear, these findings establish that deregulation of a chromatin modulator can result in site-specific copy number abnormalities. ■

Black JC, Manning AL, Van Rechem C, Kim J, Ladd B, Cho J, et al. KDM4A lysine demethylase induces site-specific copy gain and rereplication of regions amplified in tumors. *Cell* 2013;154:541–55.