

---

**SUPPLEMENTAL LABELING**


---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1278A-S3 0410**

ACCEPTED  
FOR REGISTRATION

Mar 29, 2011
--------------

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

**Revus Top®**

**Fungicide**

For control of certain diseases

Active Ingredients:	
Mandipropamid (CAS No. 374726-62-2).....	21.9%
Difenoconazole (CAS No.119446-68-3).....	21.9%
<hr/>	
Other Ingredients:	56.2%
<hr/>	
Total:	100.0%

Contains 2.08 pounds of mandipropamid active ingredient and 2.08 pounds of difenoconazole active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN.**

## **CAUTION**

**EPA Reg. 100-1278**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Revus Top as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

Crop	Target Disease	Use Rate fl oz Product per Acre	Remarks
<b>Cucurbit Vegetables</b>  Cantaloupe Cucumber Honeydew Muskmelon Watermelon Pumpkin Squash Zucchini  Including cultivars and/or hybrids of these.  See additional cucurbit crops below.	Gummy stem blight <i>(Didymella bryoniae)</i>  For Suppression of: Downy mildew <i>(Pseudoperonospora cubensis)</i>	7.0	Begin applications prior to disease onset when conditions are conducive for disease. Apply Revus Top on a 7-10 day schedule making no more than 1 application before alternating to another fungicide with a different mode of action.  For downy mildew control, Revus Top must be tank mixed with another fungicide labeled for downy mildew that has a different mode of action.  If disease pressure is high, use the shortest interval.  The addition of a spreading/penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is recommended when applying by ground or air.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Revus Top can be applied by either ground, chemigation, or aerial application. A minimum of 15 gal/A for ground applications is recommended (20 for gummy stem blight). For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

**Complete list of cucurbit vegetables:** Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.

Revus Top® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta

REV TOP 1278A-S3 0410 – df – 1/20/11