



From the office of:

**Andrew J Blackman, MD**

121 St Luke's Center Drive, Suite 302  
Chesterfield, MO 63017 314-523-2595

[AndrewBlackmanMD.com](http://AndrewBlackmanMD.com)



**POST-OPERATIVE INSTRUCTIONS  
PROCEDURE: QUADRICEPS TENDON REPAIR**

**Diet:** Begin with liquids and light foods (crackers, soup, etc.). Progress to your normal diet if you are not nauseated. You have been given a prescription for an anti-nausea medication to be taken if needed.

**Pain Medication:** You have been given a prescription for narcotic pain medicine. These types of medications can cause side effects including nausea, constipation, sedation and confusion. We recommend these only be used for 1 to 2 weeks after surgery. Do not drive while taking narcotic pain medication. An over-the-counter stool softener or laxative, such as Colace, Dulcolax, or Senokot, is recommended to prevent constipation. At any time, acetaminophen (Tylenol) may be substituted instead of your narcotic medicine and used for pain control. Do not exceed 4000 mg of acetaminophen in a 24 hour period. Do not combine with alcohol. **DO NOT** use any Non-Steroidal Anti-Inflammatory Drugs (NSAID's) such as aspirin, ibuprofen, Motrin, naproxen, or Aleve for 6 weeks after surgery.

**Response to Surgery:** It is normal to have pain and swelling in your leg after surgery. It will take several weeks for this to go away. Keep your leg elevated as much as possible to help with swelling control. It is also common to notice bruising around the leg.

**Ice:** Keep ice on your surgical site for 30 minutes at a time, then 30 minutes off. We recommend having two ice packs. Keep the unused pack in your freezer, and rotate the packs on and off the surgical site as often as needed. Continue to use the ice as often as possible for the first 5-7 days, then as needed for pain relief.

**Wound Care:** You may remove your surgical dressing after one week. Replace with a clean gauze dressing and a compressive wrap. You may have some water or blood spotting on the dressing, which is normal. If this seems excessive, call our office for further instructions.

**Brace:** Wear your brace at all times, including when you are sleeping. It will prevent your knee from bending.

**Weight-Bearing:** You may put as much weight on the operative leg as you feel comfortable; however, the brace must be in place and locked in full extension. You will need to use crutches or a walker to assist in walking.

**Showering:** When you feel up to it, you may shower. The surgical site must stay dry, though. You can use a trash bag with tape/rubber bands or purchase a waterproof cast cover. You may remove the brace when showering, but do not bear weight on the operated leg or allow the knee to bend excessively. We recommend purchasing a shower chair to facilitate safe showering.

**Exercises:** Beginning post-operative day #1, perform the following exercises.

- Ankle Pumps: While sitting, pump your ankles up and down 10 times.
- Quad Sets: While sitting or lying, tighten your quadriceps muscle; hold for 10 sec x 10
- Do this at least once an hour.

**Driving:** You will be unable to drive until you are off narcotic medications and you have been allowed to discontinue your crutches. This will be approximately 6 weeks after surgery. It is important that you feel very confident in your ability to respond quickly to changing conditions before attempting to drive. Make sure you can safely get in and out of your car, as well.

**Appointment:** A follow-up appointment should be scheduled for you. If you didn't receive an appointment, or if you need to change your appointment time, please call our office at 314-523-2595.

**Problems:** If you develop severe pain, a fever greater than 101.5°, redness around the incision, thick yellow drainage from the incision, excessive bleeding, or persistent nausea and vomiting please call our office at **314-523-2595**.

Monday - Friday, 7 a.m. to 5 p.m.

Our staff will be available to answer questions

Weekends & Evenings

Our after-hours emergency exchange will be able to contact Dr. Blackman

For more information, please visit [andrewblackmanmd.com](http://andrewblackmanmd.com)

Facebook.com/STLOrtho

Twitter - @ STLOrtho