

# Brooks Island Regional Preserve

## Checklist of Wild Plants

Column Heading	Description
<b>Scientific Name</b>	According to The Jepson Manual, Higher Plants of California, Second Edition (JM2) and eFlora (ucjeps.berkeley.edu/IJM.html)
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of The Jepson Manual (JM93) is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to JM2 and other references (not standardized)
<b>Family</b>	Scientific family name according to JM2, abbreviated by replacing the "aceae" ending with "-". (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any) as of April 2016, listed in 3 categories, divided by vertical bars ( ): Federal:California (Fed.:Calif.)   California Rare Plant Ranks (CRPR from CNPS)   East Bay chapter of the CNPS (EBCNPS) Fed.:Calif.: FE = Fed. Endangered, FT = Fed. Threatened; SC = State Candidate, SE = State Endangered, SR = State Rare CRPR (CNPS): 1B = Rare, threatened or endangered in Calif, 2B = like 1B but more common elsewhere, 3 = Review List, 4 = Watch List; suffix: 0.1 = Seriously threatened in Calif., 0.2 = Fairly threatened in Calif., 0.3 = Not very threatened in Calif. EBCNPS: * = Statewide listed rare, threatened, or endangered; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range if available) according to JM2 and other sources

Fern-like		Grouped by Growth Form					Fern-like	
<i>Polypodium</i>		Sorted Alphabetically by Scientific Name					<i>Polypodium</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)		
<b>Fern-like</b>								
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-:-   -   -	-	Np	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -		
<b>Grass-like</b>								
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass / Seashore Bentgrass	Po-	-:-   -   -	-	Np	Common. Open meadows, woodland, forest, subalpine (Jun–Aug)		
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	-:-   -   -	-	Za	Sandy soils, open or disturbed sites (Apr–Jun)		
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-:-   -   -	M	Za	Disturbed sites (Mar–Jun)		
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	-:-   -   -	-	Za	Shaded or moist, open sites (Apr–Jul)		
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-:-   -   -	L	Za	Fields, disturbed areas (Apr–Jul)		

Grass-like <i>Bromus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name						Grass-like <i>Bromus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	-:-   -   -	H	Za	Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Carex praegracilis</i>	Creeper / Freeway Sedge	Cyper-	-:-   -   C	-	Np	Common. Often alkaline, ± moist places (Apr–Aug)	
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Po-	-:-   -   -	H	Zp	Disturbed sites, many habitats, esp coastal (Sep–Feb)	
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	-:-   -   -	-	Np	Salt marshes, coastal dunes, moist, alkaline areas (Apr–Sep)	
<input type="radio"/> <i>Eleocharis</i> sp.	Spikerush	Cyper-	-:-   -   -	-	Np	--	
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western / Blue Wild-rye	Po-	-:-   -   -	-	Np	Open areas, chaparral, woodland, forest (Jun–Aug)	
<input type="radio"/> <i>Elymus gouldii</i> (JM93: <i>Leymus multiflorus</i> )	Many-flowered Wild-rye	Po-	-:-   -   -	-	Np	Open areas, often saline soils (May–Jul)	
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i> )	Beardless Wild Rye	Po-	-:-   -   -	-	Np	Dry to moist, often saline, meadows (Jun–Jul)	
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i> )	Brome Fescue	Po-	-:-   -   -	-	Za	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)	
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	-:-   -   C	-	Np	Dry, open or shady places (Jul–Sep)	
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	-:-   -   B	-	Np	Sand dunes, grassland, subalpine forest (May–Jul)	
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	-:-   -   -	-	Np	Meadows, pastures, streambanks (May–Aug)	
<input type="radio"/> <i>Juncus</i> sp.	Rush	Junc-	-:-   -   -	-	ap	--	
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-:-   -   C	-	Np	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)	
<input type="radio"/> <i>Phragmites australis</i>	Common Reed	Po-	-:-   -   B	N	Np	Pond & lake margins, sloughs, marshes (Jul–Nov)	
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-:-   -   C	-	Np	Oak woodland, chaparral, grassland (Mar–Jun)	
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	-:-   -   -	-	Np	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)	

### Herbaceous

<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-:-   -   -	-	Np	Many habitats (Apr–Sep)
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus / Common Trefoil	Fab-	-:-   -   -	-	Na	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Amaryllis belladonna</i>	Naked Ladies	Amaryllid-	-:-   -   -	-	Zp	Disturbed sites, often around abandoned home sites (Jul–Sep)
<input type="radio"/> <i>Ambrosia chamissonis</i>	Beach Bur-sage / Beach-bur	Aster-	-:-   -   A2	-	Np	Beaches, dunes (Jul–Oct)
<input type="radio"/> <i>Ambrosia psilostachya</i>	Western Ragweed	Aster-	-:-   -   -	-	Np	Common. Roadsides, dry fields (Jun–Nov)
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-:-   -   -	-	Na	Abundant. Open, generally disturbed places (Mar–Jun)

Herbaceous <i>Anagallis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Anagallis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-:-   -   -	- Za	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-:-   -   -	- Na	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe / Pipevine	Aristolochi-	-:-   -   C	- Np	Streamsides, forest, chaparral (Jan–Apr)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-:-   -   -	- Np	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Atriplex leucophylla</i>	Beach Saltbush	Chenopodi-	-:-   -   A2	- Np	Sandy soils, dunes (Apr–Oct)	
<input type="radio"/> <i>Atriplex prostrata</i> (JM93: <i>A. triangularis</i> )	Fat-hen / Spearscale	Chenopodi-	-:-   -   -	- Na	Wet places, marshes (Apr–Oct)	
<input type="radio"/> <i>Baccharis glutinosa</i> (JM93: <i>B. douglasii</i> )	Marsh Baccharis	Aster-	-:-   -   -	- Np	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)	
<input type="radio"/> <i>Bidens frondosa</i>	Sticktight	Aster-	-:-   -   B	- Na	Uncommon. Damp soil, esp disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-:-   -   -	M Za	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	-:-   -   -	L Za	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Cakile maritima</i>	Horned Sea Rocket	Brassic-	-:-   -   -	L Za	Beach dunes (May–Nov)	
<input type="radio"/> <i>Calystegia sp.</i>	Morning-glory	Convolvul-	-:-   -   -	- Np	--	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-:-   -   -	- Np	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	-:-   -   -	- Np	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	-:-   -   -	M Za	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	-:-   -   -	- Np	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-:-   -   -	- Za	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Chlorogalum pomeridianum var. pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-:-   -   -	- Np	Common. Open grassland, chaparral, woodland (May–Aug)	
<input type="radio"/> <i>Cirsium quercetorum</i>	Brownie Thistle	Aster-	-:-   -   A2	- Np	Open places, grassland, woodland (Apr–Aug)	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-:-   -   -	M Zb	Common. Disturbed areas (May–Oct)	
<input type="radio"/> <i>Clarkia purpurea subsp. ?</i>	Winecup Clarkia	Onagr-	-:-   -   -	- Na	--	
<input type="radio"/> <i>Clarkia purpurea subsp. purpurea</i>	Purple Winecup Clarkia	Onagr-	-:-   -   A2	- Na	Uncommon. Grassland (Apr–Jun)	
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	-:-   -   -	- Na	Openings in woodland, forest, chaparral near coast (May–Aug)	
<input type="radio"/> <i>Claytonia perfoliata subsp. perfoliata</i>	Common Miner's Lettuce	Monti-	-:-   -   -	- Na	Vernally moist, often shady or disturbed sites (Jan–May)	
<input type="radio"/> <i>Cotula australis</i>	Australian Brass Buttons	Aster-	-:-   -   -	- Zp	Common. Disturbed areas (Jan–May)	

Herbaceous	Grouped by Growth Form						Herbaceous
<i>Cotula</i>	Sorted Alphabetically by Scientific Name						<i>Cotula</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	-:   -   -	L	Zp	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Crassula connata</i>	Pygmy Weed	Crassul-	-:   -   -	-	Na	Open areas (Feb–May)	
<input type="radio"/> <i>Cuscuta pacifica</i> var. <i>pacifica</i> (JM93: <i>C. salina</i> var. <i>major</i> )	Goldenthread	Convolvul-	-:   -   B	-	Na	Gen on Salicornia, Jaumea in coastal salt marshes, tidal flats (Jul–Oct)	
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	-:   -   -	M	Zp	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	-:   -   -	-	Na	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Deinandra corymbosa</i> (JM93: <i>Hemizonia</i> , subsp. <i>corymbosa</i> )	Coast Spikeweed	Aster-	-:   -   A2	-	Na	Grassland, openings in scrub or woodland, dunes, disturbed sites (Mar–Nov)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	-:   -   -	-	Np	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma</i> sp.	Blue Dicks/ookow	Themid-	-:   -   -	-	Np	--	
<input type="radio"/> <i>Dudleya cymosa</i> subsp. <i>paniculata</i>	Common / Hot Rock Dudleya	Crassul-	-:   -   B	-	Np	Uncommon. Rocky outcrops, canyons (May–Jun)	
<input type="radio"/> <i>Dudleya farinosa</i>	Powdery Dudleya	Crassul-	-:   -   A1	-	Np	Common: Coastal soils and cliffs (Jun–Aug)	
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	-:   -   -	-	Np	--	
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-:   -   -	L	Za	Open, disturbed sites, grassland, scrub (Feb–Sep)	
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem / White-stem Filaree	Gerani-	-:   -   -	-	Za	Open, disturbed sites (Feb–Sep)	
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	-:   -   -	-	Np	Grassy, open areas (Feb–Sep)	
<input type="radio"/> <i>Euphorbia peplis</i>	Petty Spurge	Euphorbi-	-:   -   -	-	Za	Common. Disturbed areas (Feb–Aug)	
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	-:   -   -	H	Zp	Roadsides, disturbed sites (May–Sep)	
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	-:   -   -	-	Na	Grassy, ± shady places (Apr–Jun)	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-:   -   -	M	Za	Open, disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-:   -   -	-	Za	Open to shaded sites, disturbed ground (Feb–Aug)	
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i> )	Hairy Gumplant	Aster-	-:   -   C	-	Np	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)	
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-:   -   -	L	Zab	Common. Disturbed areas (All year)	
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-:   -   -	M	Zbp	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Horkelia californica</i> var. <i>californica</i> (JM93: subsp. <i>californica</i> )	Few-lobed Calif. Horkelia	Ros-	-:   -   A1	-	Np	Grassy openings, edges of coastal scrub, especially n slopes (Mar–Sep)	
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-:   -   -	M	Zp	Disturbed areas, grassland, open woodland (Apr–Jul)	
<input type="radio"/> <i>Iva axillaris</i> (JM93: subsp. <i>robustior</i> )	Poverty Weed	Aster-	-:   -   -	-	Np	Seasonally wet, saline habitats, roadsides (Apr–Oct)	
<input type="radio"/> <i>Jaumea carnosa</i>	Fleshy Jaumea	Aster-	-:   -   C	-	Np	Coastal salt marshes, bases of sea cliffs (Apr–Dec)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Lactuca</i>	Sorted Alphabetically by Scientific Name					<i>Lactuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-:   -   -	- Zb	Disturbed, shrubby and wooded slopes (Jun–Aug)	
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pacific Pea	Fab-	-:   -   -	- Np	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Lepidium latifolium</i>	Perennial Peppergrass	Brassic-	-:   -   -	H Zp	Pastures, disturbed areas, fields, grassland, saline meadows, streambanks, sagebrush scrub, edge of marshes (Jun–Sep)	
<input type="radio"/> <i>Limonium californicum</i>	Western Marsh-rosemary	Plumbagin-	-:   -   B	- Np	Common; coastal dunes, salt marshes (Jul–Dec)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-:   -   -	- Za	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed / Chu-chu-pate	Api-	-:   -   B	- Np	Woodland, brushy slopes (Apr–Jun)	
<input type="radio"/> <i>Lomatium caruifolium</i> var. <i>caruifolium</i>	Alkali Desertparsley	Api-	-:   -   B	- Np	Wet, clay depressions, open grassland (Mar–May)	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-:   -   -	- Na	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Madia gracilis</i>	Slender Tarweed	Aster-	-:   -   -	- Na	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	-:   -   -	- Za	Disturbed places (Mar–Jun)	
<input type="radio"/> <i>Malva pseudolavatera</i> (JM93: <i>Lavatera cretica</i> )	Cretan Mallow	Malv-	-:   -   -	- Zab	Uncommon. Disturbed places on coastal bluffs, dunes, occ inland (Apr–Jun)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-:   -   -	- Np	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-:   -   -	L Za	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i> )	Sourclover	Fab-	-:   -   -	- Zab	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>M. globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i> )	Coyote-mint	Lami-	-:   -   -	- Np	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Oenothera elata</i> subsp. <i>hookeri</i>	Hooker Evening-primrose	Onagr-	-:   -   B	- Nb	Moist, coastal, slightly inland, sandy bluffs (Jun–Sep)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	-:   -   -	M Zp	Disturbed areas, roadsides, grassland, dunes (Jan–May)	
<input type="radio"/> <i>Perideridia gairdneri</i> subsp. <i>gairdneri</i>	Gairdner's Yampah	Api-	-:   4.2   *A1	- Np	Coastal flats, grassland, pine forest (Jun–Jul)	
<input type="radio"/> <i>Perideridia</i> sp.	Yampah	Api-	-:   -   -	- Np	- -	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-:   -   -	- Np	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia distans</i>	Distant Phacelia	Boragin-	-:   -   -	- Na	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-:   -   -	- Na	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-:   -   -	L Za	Common. Disturbed areas (Apr–Aug)	

Herbaceous <i>Polygonum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Polygonum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i> )	Prostrate Knotweed	Polygon-	-:   -   -	- Za	Disturbed places (May–Nov)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-:   -   -	- Nb	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i> )	Pink Everlasting	Aster-	-:   -   -	- Nb	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)	
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i> )	Cotton-batting Plant	Aster-	-:   -   -	- Na	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-:   -   -	M Zp	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-:   -   -	L Zp	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-:   -   -	- Zp	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i> )	Willow Dock	Polygon-	-:   -   B	- Np	Wet places, margins, rocky slopes (May–Jul)	
<input type="radio"/> <i>Salicornia pacifica</i> (JM93: <i>S. virginica</i> )	Pacific Pickleweed	Chenopodi-	-:   -   -	- Np	Salt marshes, alkaline flats (Jul–Nov)	
<input type="radio"/> <i>Salsola soda</i>	Opposite-leaved Russian Thistle	Chenopodi-	-:   -   -	M Za	Uppermost intertidal zone, saline or muddy flats, open areas in salt marshes (Jul–Oct)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-:   -   -	- Np	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Blacksnakeroot / Sanicle	Api-	-:   -   -	- Np	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica, floribunda</i> )	California Figwort	Scrophulari-	-:   -   -	- Np	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-:   -   -	- Za	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>laciniata</i>	Geranium-leaved Checkerbloom	Malv-	-:   -   C	- Np	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-:   -   -	- Za	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-:   -   -	- Np	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solanum nigrum</i>	Black Nightshade	Solan-	-:   -   -	- Za	Disturbed places (Mar–Oct)	
<input type="radio"/> <i>Solanum</i> sp.	Nightshade	Solan-	-:   -   -	- ap	- -	
<input type="radio"/> <i>Soliva sessilis</i>	Field Burrweed	Aster-	-:   -   -	- Za	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	-:   -   -	- Za	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-:   -   -	- Za	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Spergula arvensis</i> (JM93: subsp. <i>arvensis</i> )	Stickwort	Caryophyll-	-:   -   -	- Za	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Spergularia</i> sp.	Sand-spurry	Caryophyll-	-:   -   -	- ap	- -	



Herbaceous	Grouped by Growth Form					Herbaceous
<b>Stachys</b>	Sorted Alphabetically by Scientific Name					<b>Stachys</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Stachys rigida</i> var. ?	Hedge-nettle	Lami-	-:- - -	- Np	--	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-:- - -	- Za	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Symphotrichum chilense</i> (JM93: <i>Aster chilensis</i> )	Pacific Aster	Aster-	-:- - -	- Np	Grassland, salt marshes, disturbed places (Jun–Oct)	
<input type="radio"/> <i>Tetragonia tetragonioides</i>	New Zealand Spinach	Aizo-	-:- - -	L Za	Common. Sand dunes, bluffs, margins of coastal wetlands (Apr–Sep)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-:- - -	- Na	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	-:- - -	- Na	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-:- - -	- Na	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-:- - -	- Na	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-:- - -	- Np	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Vicia americana</i> subsp. <i>americana</i> (JM93: var. <i>americana</i> )	American Vetch	Fab-	-:- - -	- Np	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>nigra</i>	Narrow-leaved Vetch	Fab-	-:- - -	- Za	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>sativa</i>	Spring Vetch	Fab-	-:- - -	- Za	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	-:- - C	- Np	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-:- - -	- Np	Grassland (Apr–Aug)	
<b>Woody</b>						
<input type="radio"/> <i>Acacia</i> sp.	Acacia	Fab-	-:- - -	- Zp	--	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-:- - -	- Np	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Argyranthemum foeniculaceum</i>	Canary Islands Marguerite	Aster-	-:- - -	- Zp	Escape from cult in disturbed coastal areas (Mar–Aug)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-:- - -	- Np	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	-:- - -	- Np	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Carpobrotus chilensis</i>	Sea Fig	Aizo-	-:- - -	M Zp	Common on coastal sandy shores (All year)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Carpobrotus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Carpobrotus</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Carpobrotus edulis</i>	Freeway Iceplant	Aizo-	-:-   -   -	H	Zp	Common. Many coastal habitats, esp. sand (All year)
<input type="radio"/> <i>Eriophyllum staechadifolium</i>	Seaside Woolly Sunflower	Aster-	-:-   -   A2	-	Np	Dunes, sea bluffs, coastal scrub, canyons (Apr–Sep)
<input type="radio"/> <i>Grindelia stricta</i> var. <i>angustifolia</i>	Marsh Gumplant	Aster-	-:-   -   C	-	Np	Tidal wetlands (May–Dec)
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i> )	Monterey Cypress	Cupress-	-:-   -   -	N	Np	Closed-cone-pine/cypress forests -
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-:-   -   -	-	Np	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)
<input type="radio"/> <i>Lupinus arboreus</i>	Yellow Bush Lupine	Fab-	-:-   -   A2	N	Np	Coastal bluffs, dunes, or more inland (Apr–Jul)
<input type="radio"/> <i>Phoenix canariensis</i>	Canary Island Date Palm	Arec-	-:-   -   -	L	Zp	Uncommon. Near development, disturbed areas (Oct–Apr)
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	-:-   -   -	-	Np	Closed-cone-pine forest, oak woodland -
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-:-   -   -	-	Np	Gen ± moist areas in sun, esp streambanks (Feb–Nov)
<input type="radio"/> <i>Rosa</i> sp.	Rose	Ros-	-:-   -   -	-	Np	- -
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-:-   -   -	-	Np	Open, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Salic-	-:-   -   -	-	Np	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)
<input type="radio"/> <i>Salix</i> sp.	Willow	Salic-	-:-   -   -	-	p	- -
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i> )	Blue Elderberry	Adox-	-:-   -   -	-	Np	Common. Streambanks, open places in forest (Mar–Sep)
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-:-   -   -	-	Np	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)
<input type="radio"/> <i>Washingtonia filifera</i>	California Fan Palm	Arec-	-:-   -   -	-	Np	Groves, moist places, seeps, springs, streambanks (Feb–Jun)