5

450 Broadway St. MC 6120 Redwood City, CA 94063 Ph: 650-723-5643 Fax: 650-723-3429 3801 Miranda Ave. MC Ortho 112 Palo Alto, CA 94304 Ph: 650-493-5000 x66101 Fax: 650-849-1265

ACL AUTOGRAFT WITH FEMORAL MICROFRACTURE PROTOCOL

Name:
Diagnosis:
Date of Surgery:
Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks
Weeks 0-2:
 Strict toe touch weight bearing with crutches CPM machine 6-8 hours per day WITHOUT BRACE; start 30-50 degrees and advance degrees per direction per day as tolerated up to 0-90 degrees Brace locked at full extension while ambulating and sleeping; may set 0-90 when sitting Quad sets, patellar mobs, ankle pumps
Weeks 2-6:
 Strict toe touch weight bearing with crutches as before May discontinue brace when patient comfortable and swelling has subsided CPM machine as above (6-8h/day) but increase up to max flex/extension as tolerated Focus on regaining full extension; goal ROM 0-125 degrees in PROM/AAROM
Weeks 6-8:
 Advance to full WBAT over 1-2 weeks and d/c crutches when gait normalizes Begin AROM without restriction; straight leg raises, closed chain quad exercises Normalize gait pattern; bike exercises
Weeks 8-12
 Mini-squats, weight shifts, initiate step up/down program Advance closed chain strengthening, weight bearing gastroc/soleus stretch Proprioceptive training, progressive squat program Leg press and lunges (start initially with body weight only)
Months 3-4:
 Begin forward running in straight line (no cutting/pivoting) when 8" stepdown adequate Progress strengthening and flexibility exercises
Months 4-6:
Start plyometric exercises; start sports specific agility program
Signature Date:
orginature Date