



From the office of:

Andrew J Blackman, MD

121 St Luke's Center Drive, Suite 302
Chesterfield, MO 63017 314-523-2595
AndrewBlackmanMD.com



POST-OPERATIVE INSTRUCTIONS PROCEDURE: OCD REPAIR (KNEE)

Diet: Begin with liquids and light foods (crackers, soup, etc.). Progress to your normal diet if you are not nauseated. You have been given a prescription for an anti-nausea medication to be taken if needed.

Pain Medication: You have been given a prescription for narcotic pain medicine. These types of medications can cause side effects including nausea, constipation, sedation and confusion. We recommend these only be used for 1 to 2 weeks after surgery. Do not drive while taking narcotic pain medication. An over-the-counter stool softener or laxative, such as Colace, Dulcolax, or Senokot, is recommended to prevent constipation. At any time, acetaminophen (Tylenol) may be substituted instead of your narcotic medicine and used for pain control. Do not exceed 4000 mg of acetaminophen in a 24 hour period. Do not combine with alcohol. Avoid using Non-Steroidal Anti-Inflammatory Drugs (NSAID's) such as aspirin, ibuprofen, Motrin, naproxen, or Aleve for 6 weeks after surgery.

Response to Surgery: It is normal to have pain and swelling in your leg after surgery. It will take several weeks for this to go away. Keep your leg elevated as much as possible to help with swelling control. It is also common to notice bruising around the leg.

Ice: Keep ice on your surgical site for 30 minutes at a time, then 30 minutes off. We recommend having two ice packs. Keep the unused pack in your freezer, and rotate the packs on and off the surgical site as often as needed. Continue to use the ice as often as possible for the first 5-7 days, then as needed for pain relief.

Wound Care: You may remove your surgical dressing on post-operative day #3 and replace with a clean, dry gauze dressing. You may have some water or blood spotting on the dressing, which is normal. If this seems excessive, call our office for further instructions.

Weight-Bearing: Your weightbearing status is "toe touch" which means you are allowed place the foot of the operated leg on the ground for balance, but do not put any body weight onto it. This restriction will be lifted 4-6 weeks after surgery. You will need to use crutches until then.

Showering: After post-operative day #3, you may shower. If the surgical incisions are not draining, you may allow soapy water to run over them, but do not soak or scrub the incision sites. We recommend purchasing a shower chair to facilitate safe showering.

Exercises: Beginning post-operative day #1, perform the following exercises.

- Ankle Pumps: While sitting, pump your ankles up and down 10 times.
- Quad Sets: While sitting or lying, tighten your quadriceps muscle; hold for 10 sec x 10
- Do this at least once an hour.

Driving: You will be unable to drive until you are off narcotic medications and you have been allowed to discontinue your crutches. This may be up to 6 weeks after surgery. It is important that you feel very confident in your ability to respond quickly to changing conditions before attempting to drive. Make sure you can safely get in and out of your car, as well.

Appointment: A follow-up appointment should be scheduled for you. If you didn't receive an appointment, or if you need to change your appointment time, please call our office at 314-523-2595.

Problems: If you develop severe pain, a fever greater than 101.5°, redness around the incision, thick yellow drainage from the incision, excessive bleeding, or persistent nausea and vomiting please call our office at **314-523-2595**.

Monday - Friday, 7 a.m. to 5 p.m.

Our staff will be available to answer questions

Weekends & Evenings

Our after-hours emergency exchange will be able to contact Dr. Blackman

For more information, please visit andrewblackmanmd.com

Facebook.com/STLOrtho

Twitter - @ STLOrtho