



## **The CCB Science 2 Service Distance Learning Program<sup>®</sup>**

*Please complete all required information and fax to 203.284.9500*

*For questions or assistance with the program, please email Jeff at JQuamme@ctcertboard.org*

### **S2S Course 2042 TIP 49 Incorporating Alcohol Pharmacotherapies into Medical Practice Module 1 PRE-TEST**

1. Annual economic costs of AUDs in the United States has been estimated at:
  - a) \$185 million
  - b) \$200 million
  - c) \$300 million
  - d) \$1 billion
  
2. The most common and persistent side effect of acamprosate is \_\_\_\_\_.
  - a) sexual dysfunction
  - b) diarrhea
  - c) nausea
  - d) vomiting
  
3. The disulfiram-alcohol reaction usually begins \_\_\_\_\_ after alcohol is ingested.
  - a) immediately
  - b) 10-30 minutes
  - c) 45 min to 1 hour
  - d) 1-1 1/2 hours
  
- 4) Alcohol ingestion may produce unpleasant symptoms for up to \_\_\_\_\_ after the last dose of disulfiram is taken.
  - a) 3 days
  - b) 1 week
  - c) 10 days
  - d) 2 weeks
  
5. The capacity to arrange for ongoing supervision if disulfiram ingestion may be limited in a/an \_\_\_\_\_ setting.
  - a) inpatient
  - b) intermediate care
  - c) primary care
  - d) residential
  
6. Disulfiram appears to have \_\_\_\_\_ clinical efficacy in maintaining abstinence in individuals with AUDs.
  - a) modest
  - b) well documented
  - c) no
  - d) statistically significant