

Solomon Press
Core Mathematics C2
Paper C
(Mark Scheme)

**All exam papers are issued free to students for education purpose only.
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations
Advanced Subsidiary

Core Mathematics C2

Paper C

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

C2 Paper C – Marking Guide

1. $(1-x)^6 = 1 + 6(-x) + \binom{6}{2}(-x)^2 + \dots = 1 - 6x + 15x^2$ M1 A1
 $(1+x)(1-x)^6 = (1+x)(1 - 6x + 15x^2 + \dots)$
 coeff. of $x^2 = 15 - 6 = 9$ M1 A1 (4)

2. (a) $\frac{a[1-(\frac{1}{3})^4]}{1-\frac{1}{3}} = 200$ M1 A1
 $a = 200 \times \frac{27}{40} = 135$ A1


(b) $= \frac{135}{1-\frac{1}{3}} = 202\frac{1}{2}$ M1 A1 (5)

3. (a) $(-4, 0) \therefore 0 = 4 - 20 + 16k + 128$ M1
 $16k = -112, k = -7$ A1

(b) $4 + 5x - 7x^2 - 2x^3 = 0$
 $x = -4$ is a solution $\therefore (x + 4)$ is a factor B1

$$\begin{array}{r}
 -2x^2 + x + 1 \\
 x + 4 \overline{) -2x^3 - 7x^2 + 5x + 4} \\
 \underline{-2x^3 - 8x^2} \\
 x^2 + 5x \\
 \underline{x^2 + 4x} \\
 x + 4 \\
 \underline{x + 4} \\
 0
 \end{array}$$

$\therefore (x + 4)(1 + x - 2x^2) = 0$ M1 A1
 $(x + 4)(1 + 2x)(1 - x) = 0$ M1
 $x = -4$ (at A), $-\frac{1}{2}, 1$
 $\therefore (-\frac{1}{2}, 0), (1, 0)$ A1 (7)



5. (a) $= 3 - \log_8 8^{\frac{2}{3}}$ B1 M1 A1
 $= 3 - \frac{2}{3} = \frac{7}{3}$ A1

(b) $(2^2)^x - 3(2 \times 2^x) = 0$ M1
 $(2^x)^2 - 6(2^x) = 0$
 $2^x(2^x - 6) = 0$ M1
 $2^x = 0$ (no solutions) or 6 A1
 $x = \frac{\lg 6}{\lg 2} = 2.58$ (3sf) M1 A1 (9)

6. (a) $f'(x) = -1 + 2x^{-\frac{1}{3}}$ M1 A1
 $f''(x) = -\frac{2}{3}x^{-\frac{4}{3}}$ A1
- (b) for TP, $-1 + 2x^{-\frac{1}{3}} = 0$ M1
 $x^{\frac{1}{3}} = 2$ M1
 $x = 8$ A1
 $\therefore (8, 6)$ A1
- (c) $f''(8) = -\frac{1}{24}$, $f''(x) < 0 \therefore$ maximum M1 A1 (9)

7. (a) $\text{grad } PQ = \frac{8-2}{-3-(-5)} = 3$, $\text{grad } QR = \frac{4-8}{9-(-3)} = -\frac{1}{3}$ M1 A1
 $\text{grad } PQ \times \text{grad } QR = 3 \times (-\frac{1}{3}) = -1$ M1
 $\therefore PQ$ perp. to QR , $\therefore \angle PQR = 90^\circ$ A1
- (b) $\angle PQR = 90^\circ \therefore PR$ is a diameter M1
 \therefore centre = mid-point of $PR = (\frac{-5+9}{2}, \frac{2+4}{2}) = (2, 3)$ M1 A1
- (c) radius = dist. $(-5, 2)$ to $(2, 3) = \sqrt{49+1} = \sqrt{50}$ B1
 $\therefore (x-2)^2 + (y-3)^2 = (\sqrt{50})^2$ M1
 $x^2 - 4x + 4 + y^2 - 6y + 9 = 50$
 $x^2 + y^2 - 4x - 6y = 37$ [$k = 37$] A1 (10)

8. (a) $= 12 \times (2\pi - \frac{2\pi}{3}) = 16\pi$ cm M1 A1
- (b) chord $= 2 \times 12 \sin \frac{\pi}{3} = 24 \times \frac{\sqrt{3}}{2} = 12\sqrt{3}$ M1 A1
 $P = (12 \times \frac{2\pi}{3}) + 12\sqrt{3}$ M1
 $= 8\pi + 12\sqrt{3} = 4(2\pi + 3\sqrt{3})$ cm [$k = 4$] A1
- (c) area of segment $= (\frac{1}{2} \times 12^2 \times \frac{2\pi}{3}) - (\frac{1}{2} \times 12^2 \times \sin \frac{2\pi}{3})$ M2
 $= 72(\frac{2\pi}{3} - \frac{\sqrt{3}}{2}) = 88.443$
as % of area of circle $= \frac{88.443}{\pi \times 12^2} \times 100\% = 19.6\%$ (1dp) M1 A1 (10)

9. (a)
- | | | | | | | |
|--|---|-------|---|-------|-------|----------|
| | x | 2 | 4 | 6 | 8 | |
| | $1 + 3\sqrt{x}$ | 5.243 | 7 | 8.348 | 9.485 | M1 A1 |
| | area $\approx \frac{1}{2} \times 2 \times [5.243 + 9.485 + 2(7 + 8.348)]$ | | | | | B1 M1 A1 |
| | $= 45.4$ (3sf) | | | | | A1 |
- (b) $= \int_2^8 (1 + 3\sqrt{x}) \, dx$
 $= [x + 2x^{\frac{3}{2}}]_2^8$ M1 A1
 $= [8 + 2(2\sqrt{2})^3] - [2 + 2(2\sqrt{2})]$ M1
 $= (8 + 32\sqrt{2}) - (2 + 4\sqrt{2})$ M1
 $= 6 + 28\sqrt{2}$ A1
- (c) $= \frac{(6 + 28\sqrt{2}) - 45.4}{6 + 28\sqrt{2}} \times 100\% = 0.43\%$ M1 A1 (13)

Total (75)