

**Solomon Press**  
**Statistics S1**  
**Paper H**  
**(Mark Scheme)**

**All exam papers are issued free to students for education purpose only.  
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)  
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations  
Advanced Subsidiary / Advanced Level

**Statistics**  
**Module S1**

Paper H

**MARKING GUIDE**

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



*Written by Shaun Armstrong & Chris Huffer*

© *Solomon Press*

*These sheets may be copied for use solely by the purchaser's institute.*

## S1 Paper H – Marking Guide

1. (a)  $\sum xP(x) = \frac{1}{8}k + \frac{3}{8}(k+4) + \frac{1}{2}(2k) = \frac{3}{2}(k+1)$  M2 A1  
 (b)  $\frac{3}{2}(k+1) = 9; k = 5$  M1 A1 **(5)**

2. (a) e.g. using a distribution or other simplified way of representing a real situation that allows predictions to be made about it B2  
 (b) (i) not suitable e.g. discrete etc. / +ve skew B2  
 (ii) suitable e.g. likely to be similar time most days, sometimes fair bit more, sometimes fair bit less B2  
 (iii) not suitable e.g. very different values in winter / summer B2 **(8)**

3. (a)  $1 - 0.22 = 0.78$  M1 A1  
 (b)  $0.78 - 0.35 = 0.43$  M1 A1  
 (c)  $\frac{P(A \cap B)}{P(A)} = \frac{0.7 - 0.43}{0.7} = 0.386$  (3sf) M2 A1  
 (d) not independent as e.g.  $P(B|A) \neq P(B)$  B2 **(9)**


4. (a)  $P(Z < \frac{127 - 122.3}{2.6}) = P(Z < 1.81) = 0.9649$  M2 A1  
 (b)  $P(Z < \frac{121.5 - 122.3}{2.6}) = P(Z < -0.31) = 0.3783$  M2 A1  
 (c)  $P(Z < \frac{454 - \mu}{1.6}) = 0.05$  M1  
 $\frac{454 - \mu}{1.6} = -1.6449; \mu = 456.6$  (4sf) M1 A2 **(10)**

5. (a) 5 vowels, 7 consonants  
 $P(V = 1) = 3 \times \frac{5}{12} \times \frac{7}{11} \times \frac{6}{10} = \frac{21}{44}$  M2 A1  
 (b)  $P(V = 0) = \frac{7}{12} \times \frac{6}{11} \times \frac{5}{10} = \frac{7}{44}$   
 $P(V = 2) = 3 \times \frac{5}{12} \times \frac{4}{11} \times \frac{7}{10} = \frac{7}{22}$   
 $P(V = 3) = \frac{5}{12} \times \frac{4}{11} \times \frac{3}{10} = \frac{1}{22}$   

$v$	0	1	2	3	
$P(V = v)$	$\frac{7}{44}$	$\frac{21}{44}$	$\frac{7}{22}$	$\frac{1}{22}$	M2 A2

 (c)  $E(V) = \sum vP(v) = 0 + \frac{21}{44} + \frac{14}{22} + \frac{3}{22} = \frac{5}{4}$  M1 A1  
 $E(V^2) = \sum v^2P(v) = 0 + \frac{21}{44} + \frac{28}{22} + \frac{9}{22} = \frac{95}{44}$  M1 A1  
 $\text{Var}(V) = \frac{95}{44} - (\frac{5}{4})^2 = \frac{105}{176}$  or 0.597 (3sf) M1 A1 **(13)**

6. (a) freq. dens. = 0.9, 1, 1.65, 1.2, 0.72, 0.78, 0.52 M1 A1




B2

- (b) cum. freqs: 36, 56, 89, 113, 149, 188, 240  
 $Q_1 = 60^{\text{th}} = 60.5 + 20\left(\frac{4}{33}\right) = 62.9$  [60.25<sup>th</sup> → 63.1]  
 $Q_2 = 120^{\text{th}} = 100.5 + 50\left(\frac{7}{36}\right) = 110.2$  [120.5<sup>th</sup> → 110.9]  
 $Q_3 = 180^{\text{th}} = 150.5 + 50\left(\frac{31}{39}\right) = 190.2$  [180.75<sup>th</sup> → 191.2] M1  
M2 A3
- (c)  $Q_3 - Q_2 = 80.0$ ,  $Q_2 - Q_1 = 47.3$ ;  $Q_3 - Q_2 > Q_2 - Q_1 \therefore$  +ve skew M2 A1 (13)

7. (a) 

$x$	1	0.5	0.333	0.25	0.2	0.167
$d$	102	69	61	58	52	48

M1 A1



B3

- (b) the points lie roughly on a straight line B1
- (c)  $S_{xd} = 189.733 - \frac{2.45 \times 390}{6} = 30.483$  M1  
 $S_{xx} = 1.491 - \frac{2.45^2}{6} = 0.490583$  M1  
 $b = \frac{30.483}{0.490583} = 62.136$  M1 A1  
 $a = \frac{390}{6} - (62.136 \times \frac{2.45}{6}) = 39.628$  M1 A1  
 $d = 39.6 + 62.1x$  A1
- (d)  $m = 13$ ,  $x = \frac{1}{13}$ ;  $d = 39.6 + (62.1 \times \frac{1}{13}) = 44.4$ , so 44 cases M2 A1
- (e) not very reliable as it requires extrapolation well outside the data B1 (17)

Total (75)