

PDRX Clinic  
727 N Ann Arbor  
Oklahoma City, OK 73127  
405-942-3040

NDC: 55289-0707-20  
Lot: f11d40  
Refills: 0

Bellatrix Lestrangle  
DEA: AJ3247194

**TAKE WITH FOOD IF UPSET  
STOMACH OCCURS**

PDRX Clinic  
727 N Ann Arbor  
Oklahoma City, OK 73127

Bellatrix Lestrangle  
DEA: AJ3247194  
Ph # 405-942-3040

**Rx # 1600** 10/30/2015  
**JOHN SMITHE**  
**USE AS DIRECTED**  
**AMOXICILLIN 875MG 20 TABLETS**  
**Manufacturer: HIKMA**

AMOXICILLIN 875MG 20 TABLETS  
USE AS DIRECTED

JOHN SMITHE MRID: JS8  
Rx # 1600 Refills: 0  
" Lot: f11d40  
NDC: 55289-0707-20

Do you require additional counseling for this medication? YES / NO

Signature: \_\_\_\_\_ 10/30/2015

.. Exp/Discard: 3/31/2013

JS8 JOHN SMITHE  
AMOXICILLIN 875MG 20 TABLETS  
USE AS DIRECTED

DOB: 10/01/2015 JS8



JS8 JOHN SMITHE 10/30/2015  
AMOXICILLIN 875MG 20 TABLETS  
USE AS DIRECTED

Bellatrix Lestrangle 55289-0707-20  
Chart 10/30/2015 NB

Rx # 1600  
\$6.65 Folder Tab

Bellatrix Lestrangle 55289-0707-20 Rx # 1600  
Billing NB AWP / Qty : \$87.20 / 100 \$6.65

JS8 JOHN SMITHE  
Rx # 1600  
Bellatrix Lestrangle

DOB: 10/01/2015  
Refills: 0  
Lot Number: f11d40

PDRX Clinic 10/30/2015  
727 N Ann Arbor  
Oklahoma City, OK 73127 405-942-3040

AMOXICILLIN 875MG 20 TABLETS  
USE AS DIRECTED

AMOXICILLIN 875MG 20 TABLETS  
USE AS DIRECTED

10/30/2015 \$6.65

Bellatrix Lestrangle  
Rx # 1600  
Refills: 0

**AMOXICILLIN ORAL (AMOXIL)**

**WHY IS IT PRESCRIBED?** Used to eliminate bacteria that causes infections. **BEFORE USING THIS MEDICINE: Tell your doctor if you are having any of the following conditions:** KIDNEY DISEASE.  
**HOW MEDICINE IS ADMINISTERED?** Use this medicine exactly as directed on the label, unless instructed differently by your doctor \* Use this medicine for the full prescribed treatment period. \* This medicine can be taken with or without food, REGARDLESS of meal times. If stomach upset occurs, take with food. **WHAT TO DO IF YOU MISS A DOSE?** Take or Use when remembered unless time for next dose. No double doses. **HOW THIS MEDICINE SHOULD BE STORED?** Keep in original closed container in a dark, cool, and dry place. Refrigerate liquid forms and discard unused portion.  
**WARNINGS/PRECAUTIONS:** \* STOP TAKING MEDICINE & GET EMERGENCY HELP IMMEDIATELY IF A HYPERSENSITIVITY OR ANAPHYLACTIC REACTION occurs: (see symptoms below):- \* PEDIATRIC: Dosage alert for children LESS than 3 months of age. \* shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching. \* There are no adequate or well-controlled safety studies in pregnant women. NOTIFY YOUR DOCTOR IF PREGNANT or intend to become pregnant. \* TELL YOUR DOCTOR if you have ever had any unusual or ALLERGIC reaction to any of the CEPHALOSPORINS, PENICILLINS, or penicillin-like medicines. \* The effect of BIRTH CONTROL pills may be decreased while taking this medicine. To prevent pregnancy, use an additional form of birth control. \* To prevent drug resistance, this medication should be used to treat BACTERIAL INFECTIONS ONLY. It is not effective against viral infections. \* This medicine may cause false test results with some urine tests. If you are DIABETIC, check with your doctor before changing diabetes regimen. \* If DIARRHEA develops/persists and is severe or lasts more than 2 or 3 days, call your doctor. **ADVERSE REACTIONS: Stop taking this medicine and get emergency help IMMEDIATELY if you experience:** Shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching. **Stop taking this medicine and notify your doctor AS SOON AS POSSIBLE if you experience:** Skin rash; Unusual bleeding or bruising. **Other Common Side Effects:** Severe allergic reaction; Headache; Diarrhea; Nausea. If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **Taking this medication may cause the following mineral depletion:** BIOTIN FORTE; INOSITOL ORAL; LACTOBACILLUS HELVETICUS ORAL (LACTINEX); VITAMIN B1 ORAL (THIAMINE); VITAMIN B2 ORAL (RIBOFLAVIN); VITAMIN B3 ORAL (NICOTINAMIDE); VITAMIN B6 ORAL (PYRIDOXINE); VITAMIN B12 ORAL; PHYTONADIONE-VIT K1 ORAL (MEPHYTON); VITAMIN K. **DISCLAIMER:** This leaflet has been developed by CliniDATA Source, Inc. based primarily on labeling information provided by the manufacturer. The information does not cover all possible uses, actions, precautions, side effects, or interactions of this medicine. It is not intended as medical advice for individual patients. Copyright CliniDATA Source, Inc. 09/15. All rights reserved. Copyright PD-Rx Pharmaceuticals, Inc. 2006,2007