

**Solomon Press**  
**Core Mathematics C2**  
**Paper A**  
**(Mark Scheme)**

**All exam papers are issued free to students for education purpose only.  
Mr.S.V.Swarnaraja (Marking Examiner, Team Leader & Author)  
www.swanash.com, Mobile: +94777304755 , email: swa@swanash.com**

GCE Examinations  
Advanced Subsidiary

## Core Mathematics C2

Paper A

### MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



*Written by Shaun Armstrong*

© *Solomon Press*

*These sheets may be copied for use solely by the purchaser's institute.*

## C2 Paper A – Marking Guide

- 
1. (a)  $f(-2) = -35 \quad \therefore -24 - 8 - 2k + 9 = -35$  M1  
 $k = 6$  A1
- (b)  $= f\left(\frac{2}{3}\right)$  B1  
 $= 3\left(\frac{8}{27}\right) - 2\left(\frac{4}{9}\right) + 6\left(\frac{2}{3}\right) + 9 = \frac{8}{9} - \frac{8}{9} + 4 + 9 = 13$  M1 A1 (5)
- 
2.  $x$      $-2$      $-1$      $0$      $1$      $2$   
 $2^x$      $\frac{1}{4}$      $\frac{1}{2}$      $1$      $2$      $4$  B1  
 area  $\approx \frac{1}{2} \times 1 \times \left[\frac{1}{4} + 4 + 2\left(\frac{1}{2} + 1 + 2\right)\right]$  B1 M1 A1  
 $= 5\frac{5}{8}$  or 5.63 (3sf) A1 (5)
- 
3.  $\tan^2 \theta = \frac{1}{3}$  M1  
 $\tan \theta = \pm \frac{1}{\sqrt{3}}$  A1  
 $\theta = \frac{\pi}{6}, \frac{\pi}{6} - \pi$  or  $\pi - \frac{\pi}{6}, -\frac{\pi}{6}$  B1 M1  
 $\theta = -\frac{5\pi}{6}, -\frac{\pi}{6}, \frac{\pi}{6}, \frac{5\pi}{6}$  A2 (6)
- 
4. (a)  $= 1 + 8(3x) + \binom{8}{2}(3x)^2 + \binom{8}{3}(3x)^3 + \dots$  M1 A1  
 $= 1 + 24x + 252x^2 + 1512x^3 + \dots$  M1 A1
- (b)  $x = 0.001$  B1  
 $(1.003)^8 \approx 1 + 0.024 + 0.000\ 252 + 0.000\ 001\ 512$  M1  
 $= 1.024\ 253\ 5$  (8sf) A1 (7)
- 
5. (a) (i)  $= 2 \log_3 x = 2t$  M1 A1  
 (ii)  $= \frac{\log_3 x}{\log_3 9} = \frac{\log_3 x}{2} = \frac{1}{2}t$  M1 A1
- (b)  $2t - \frac{1}{2}t = 4$   
 $t = \frac{8}{3}$  M1  
 $\log_3 x = \frac{8}{3}, \quad x = 3^{\frac{8}{3}} = 18.7$  M1 A1 (7)
- 
6. (a) radius  $= \sqrt{25+1} = \sqrt{26}$  M1 A1  
 $\therefore (x+3)^2 + (y-2)^2 = (\sqrt{26})^2$  M1  
 $(x+3)^2 + (y-2)^2 = 26$  A1
- (b)  $(-4, 7)$ , LHS  $= (-4+3)^2 + (7-2)^2 = 1 + 25 = 26 \quad \therefore$  lies on circle B1
- (c) grad of radius  $= \frac{7-2}{-4-(-3)} = -5$  M1  
 $\therefore$  grad of tangent  $= \frac{-1}{-5} = \frac{1}{5}$  M1 A1  
 $\therefore y - 7 = \frac{1}{5}(x + 4)$  M1  
 $5y - 35 = x + 4$   
 $x - 5y + 39 = 0$  A1 (10)
-

7.	(a)	$2x^2 + 6x + 7 = 2x + 13$ $x^2 + 2x - 3 = 0$ $(x + 3)(x - 1) = 0$ $x = -3, 1$ $\therefore (-3, 7), (1, 15)$	M1 M1 A1 A1
	(b)	area under curve = $\int_{-3}^1 (2x^2 + 6x + 7) dx$ $= [\frac{2}{3}x^3 + 3x^2 + 7x]_{-3}^1$ $= (\frac{2}{3} + 3 + 7) - (-18 + 27 - 21) = 22\frac{2}{3}$ area of trapezium = $\frac{1}{2} \times (7 + 15) \times 4 = 44$ shaded area = $44 - 22\frac{2}{3} = 21\frac{1}{3}$	M1 A2 M1 B1 M1 A1 (11)

8.	(a)	$\frac{a(r^4 - 1)}{r - 1} = 10 \times \frac{a(r^2 - 1)}{r - 1}$ $r^4 - 1 = 10(r^2 - 1)$ $r^4 - 10r^2 + 9 = 0$ $(r^2 - 1)(r^2 - 9) = 0$ $r^2 = 1, 9$ $r = \pm 1, \pm 3$ $r > 1 \therefore r = 3$	B1 M1  A1 M1 M1 A1
	(b)	$\frac{a(3^3 - 1)}{3 - 1} = 26$ $a = \frac{26}{13} = 2$	M1 A1 A1
	(c)	$S_6 = \frac{2(3^6 - 1)}{3 - 1} = 728$	M1 A1 (11)

9.	(a)	area = $2xy + (\frac{1}{2} \times x^2 \times 0.5) = 2xy + \frac{1}{4}x^2 = 50$ $\therefore y = \frac{50 - \frac{1}{4}x^2}{2x} = \frac{25}{x} - \frac{1}{8}x$ $P = 2x + 4y + (x \times 0.5) = \frac{5}{2}x + 4y$ $= \frac{5}{2}x + 4(\frac{25}{x} - \frac{1}{8}x)$ $= \frac{5}{2}x + \frac{100}{x} - \frac{1}{2}x = 2x + \frac{100}{x}$	M1 A1 M1 M1 A1
	(b)	$\frac{dP}{dx} = 2 - 100x^{-2}$ for minimum, $2 - 100x^{-2} = 0$ $x^2 = 50$ $x = \sqrt{50}$ or $5\sqrt{2}$	M1 A1 M1 A1
	(c)	$\frac{d^2P}{dx^2} = 200x^{-3}$ when $x = 5\sqrt{2}$ , $\frac{d^2P}{dx^2} = \frac{2}{5}\sqrt{2}$ , $\frac{d^2P}{dx^2} > 0 \therefore$ minimum	M1 A1
	(d)	$= 2(5\sqrt{2}) + \frac{100}{5\sqrt{2}} = 10\sqrt{2} + 10\sqrt{2} = 20\sqrt{2}$	M1 A1 (13)

Total (75)