

Cervical Collar

You have been fit with a Cervical Collar. Unless otherwise instructed, please use a gradual break in schedule to allow your body to adjust to the device. Start with one hour today and increase one hour per day as tolerated. Following use, remove your orthosis and inspect your skin. There may be some skin redness from pressure areas. Redness should not persist for more than minutes after removal. Redness that does not disappear can be a sign of excess pressure and should be addressed. Discontinue use if any sign of numbness, tingling, color change, excessive itching, swelling or pain develops. Please contact our office for adjustments and do not reapply orthosis. Failure to do so could lead to blisters or open sores.



15

*Your orthosis may
look different.*

The collar is designed to support your neck and limit the motion of your spine. You should wear it 24 hours each day unless otherwise instructed by your doctor. Washing of the orthosis should be done only with your doctor's authorization. If it is approved, your neck must be immobilized during the procedure. Your orthosis or lining can be cleaned with a wash cloth and mild soapy water. Make certain all soap residue is removed and the orthosis is completely dry before re-applying. Make certain the orthosis is reapplied properly, with all straps secured snugly.

Make sure you are comfortable with how to properly put on and take off your orthosis. Never attempt to modify your orthosis yourself.

Please inspect your orthosis daily for any signs of wear including cracking, loose parts, or decreased effectiveness of the device and call our office as needed. A check-up every 3-6 months is recommended.

If you have any questions or concerns, please contact our office.

Arise Orthotics & Prosthetics
Blaine Medical Center
11855 Ulysses St NE, Suite 230
Blaine, MN 55434
763-755-9500 Phone
763-755-9510 Fax