

**Mouse Anti-CD138 [B-A38]: MC0510**

**Intended Use:** For Research Use Only

**Description:** CD138, also known as Syndecan-1, is a member of the transmembrane heparan sulfate proteoglycan family, acts as an extracellular matrix receptor and is involved in many cellular functions, including cell-cell adhesion and cell-matrix adhesion. CD138 expression is found in both hematopoietic and non-hematopoietic cells. In the hematopoietic system, CD138 labels plasma cells. It is an excellent marker for plasmacytic differentiation within the spectrum of hematologic malignancy. Among non-hematolymphoid cells, CD138 reactivity is observed in many types of epithelial cells and stoma cells in both normal and tumor tissues.

**Specifications:**

Clone: B-A38  
Source: Mouse  
Isotype: IgG2a/k  
Reactivity: Human  
Localization: Membrane  
Formulation: Purified ascites in PBS pH7.4, containing BSA, glycerol, and ≤0.09% sodium azide (NaN<sub>3</sub>).  
Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles  
Applications: IHC  
Package:

Description	Catalog No.	Size
CD138 Concentrated	MC0510	1 ml

**IHC Procedure\*:**

Positive Control Tissue: Tonsil, plasmacytoma  
Concentrated Dilution: 20-100  
Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C  
Incubation Time and Temp: 30-60 minutes @ RT  
Detection: Refer to the detection system manual

\* Result should be confirmed by an established diagnostic procedure.