DOCUMENT SCAN CONTROL SHEET

Document ID	505355	

EPA# 100-1098

EPA# For SLN

Product QUADRIS FLOWABLE FUNGICIDE

<u>Name</u>

<u>Accepted</u>

Date 1/27/2006

<u>Document</u>

Type 2EELAB

Registration ID 106907

PM: SJJ



MORRA

Section 2 (ee) Recommendation

Date:

November 22, 2005

Products:

Quadris® Flowable Fungicide

EPA Reg. No. 100-1098

Use:

Reduced rates of Quadris Flowable Fungicide on soybean for foliar

disease control

State:

New York

This recommendation is made under authority of FIFRA Section 2(ee) and recommends the use of Quadris Flowable Fungicide on soybean at reduced rates for foliar disease control.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Target Diseases	Use Rate Fl. Oz. Product/A (lb. a.i./A)
Aerial blight (Rhizoctonia solani) Rust (Phakopsora spp.) Anthracnose (Colletotrichum truncatum) Alternaria leaf spot (Alternaria spp) Brown spot (Septoria glycines)	6 - 15 (0.1 - 0.25)
spot	DOC ID 505355
(Cercospora kikuchii) Frogeye leafspot (Cercospora sojina) Pod and Stem blight (Diaporthe phaseolorum)	ACCEPTED
	Aerial blight (Rhizoctonia solani) Rust (Phakopsora spp.) Anthracnose (Colletotrichum truncatum) Alternaria leaf spot (Alternaria spp) Brown spot (Septoria glycines) Cercospora blight and leaf spot (Cercospora kikuchii) Frogeye leafspot (Cercospora sojina) Pod and Stem blight

NEW YORK STATE DEPARAMENT OF ENVIRONMENTAL CONSENSATION DIVISION OF SOLID & HAZARDOLD MADE PESTICIDE PRODUCT SEASON

Remarks

Integrated Pest (Disease) Management: Quadris should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, plant residue management, crop rotation and proper timing and placement of irrigation.

Resistance Management: No more than two foliar applications of Quadris or other Qol fungicides should be made per growing season. Do not alternate or tank mix with fungicides to which resistance has developed in the pathogen population.

Application Directions: Quadris applications should begin prior to disease development. Use the high rates under conditions favorable for severe disease pressure, dense plant canopies, or when susceptible varieties are planted. Contact Extension personnel for local economic thresholds and timings for specific diseases in your area. Applications may be made by ground, air or chemigation. An adjuvant may be added at recommended rate. Use of a non-ionic surfactant with the lower use rate is recommended.

Specific Use Restrictions:

- Do not apply more than 2.88 quarts of product/crop/acre/season (1.5 lbs. a.i./A).
- Do not make more than one application at 15 fl. oz. product/acre or 0.25 lb. a.i./A
 to soybean forage and hay.
- Do not apply within 14 days of harvest of soybeans (bean).
- May be applied the day of harvest to soybean forage and hay.

The user must refer to the federally approved label and read and follow all directions for use, restrictions, and precautions.

The user should have this recommendation in its possession at the time of use.

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the EPA.

Syngenta Crop Protection, Inc. Greensboro, NC 27419

Quadris® trademark of a Syngenta group company.

Effective Date: November 22, 2005

Label Code: NY1098020AC1105

ACCEPTED

JAN 2 7 2006

NEW YORK STATE DEPARTMENT OF ENVIRONMENTAL CONSCIENTATION DIVISION OF SOLID & HAZARDOUS MAN ROPESTICIDE PRODUCT REGISSELVEN ROPESTICIDE PRODUCT ROPESTICIDE PR