# **Kittitas County Prehospital Care Protocols**

## SUBJECT: ASPIRIN (per off-line Medical Control)

## INDICATIONS FOR USE IN AN ACUTE CORONARY EVENT:

- 1) Patient exhibits any of the following signs or symptoms:
  - a) Uncomfortable pressure, fullness, squeezing, or pain in the center of the chest that lasts more than a few minutes, or goes away and comes back.
  - b) Pain that spreads to the shoulders, neck, or arms.
  - c) Chest discomfort with light headedness, fainting, sweating, nausea, or shortness of breath.

-OR-

- 2) Patient exhibits any **two** of the following signs or symptoms, and you think it is of cardiac origin:
  - a) Atypical chest pain, stomach, or abdominal pain. This may include discomfort that can be localized to a point, that is "sharp" in nature, that is reproducible by palpation, or that is in the "wrong" location (such as the upper abdomen).
  - b) Unexplained nausea (without vomiting) or lightheadedness (not vertigo) without chest pain.
  - c) Shortness of breath and difficulty breathing (with or without chest pain).
  - d) Unexplained anxiety, weakness, or fatigue.
  - e) Palpitations, cold sweat or paleness.

### **CONTRAINDICATIONS FOR USE:**

- 1) Patient is allergic to Aspirin or Ibuprofen (Motrin®, Advil®).
- 2) If the patient has just taken the recommended dose of aspirin for this event, do not administer Aspirin.
- 3) Radial pulses are not equal strength in both arms and systolic BP varies by at least 20 mm/Hg.
- 4) Head injury suspected (includes stroke) Check w/ medical control if uncertain

### PROCEDURE:

- 1) ALS/ILS upgrade and evaluation required unless ALS/ILS is unavailable.
- 2) Be sure that the patient is alert and responsive with ability to swallow.
- 3) If the patient has his/her own nitroglycerin and meets the criteria for administration, do not delay assisting in the administration per protocol.
- 4) Check radial pulses in both arms. If both are present and equal, have the patient chew four 81 mg aspirin or equivalent (324 mg). "Chewable" Aspirin are recommended.
- 5) Record all treatment, including the dosage and the time of administration.