

Main Office:

Scottsdale Vein & Proctology Center
8752 E Via De Commercio #2
Scottsdale, Arizona 85258
Office: (602) 492-9919 | Mobile: (602) 920-1023

RICK A SHACKET
DO, MD(H)



Name: _____ DOB: _____ Date: _____

DISPENSE ONE OF THE FOLLOWING PRESCRIPTIONS:

IMIQUIMOD

(\$28.16 with GoodRx coupon)

Aldara 5% Imiquimod topical cream:

- 5 % (12 single-use pkts, 0.25 g):
- condyloma acuminata, external [apply 3x/wk at bedtime] Max: x16wk; Info: apply until total clearance or max duration
- Substitution OK
- Refill 3 times

*genital HSV, antiviral-resistant [apply qhs x5 days] Info: for immunocompromised pts

(\$1,223.99 w/o manufacturers coupon)

Zyclara 3.75% imiquimod topical cream:

- 3.75 % (1 carton, 28 single-use pkts):
- condyloma acuminata, external [apply 3.75% qhs] Max: 8wk; Info: apply until total clearance or max duration
- Substitution OK
- Refill 1 time

Rick A. Shacket, DO, MD (H), BS9262611
Diplomate American Osteopathic Board of Proctology

LOCATIONS

Scottsdale Vein Center 8752 E Via De Commercio, Suite 2, Scottsdale, Arizona 85258, 602.492.9919

Dr. Rick Shacket PLLC 3543 N. 7th Street, Phoenix AZ 85014, 602.492.9919

Rick Shacket, DO, MD(H) 81 W. Guadalupe Road, Suite 111, Gilbert, AZ 85233, 602.492.9919

What is the most important information I should know about imiquimod topical?

Using too much of this medicine or using it for too long can increase your risk of severe skin reactions. Follow your doctor's instructions.



Do not use imiquimod topical on areas of broken, wounded, or burned skin. Wait until these conditions have healed before using imiquimod topical.

Before using this medication, tell your doctor if you have a weak immune system, an autoimmune disorder, graft-versus-host disease, or if you have recently received a bone marrow transplant or cord blood transplant.

When treating genital warts around the vagina, avoid getting the cream on the more sensitive inner layers of vaginal tissue. This could result in vaginal swelling or irritation and painful urination.



Avoid exposure to sunlight or tanning beds. Imiquimod topical can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Imiquimod topical is not a cure for genital warts and it may not keep you from spreading this condition to others through vaginal, anal, or oral sex. You may develop new lesions during treatment with imiquimod topical. For best results, keep using the medicine for the entire length of time prescribed by your doctor.

Imiquimod will not protect against sexually transmitted diseases such as chlamydia, gonorrhea, herpes, HIV, syphilis, and trichomoniasis.



If you are treating the genital or rectal area with imiquimod topical, avoid sexual activity while the medicine is on your skin. Imiquimod topical can weaken the rubber that condoms or diaphragms are made out of. If you use a condom or diaphragm, these items could break if the rubber weakens, and an unplanned pregnancy could result.




Before applying imiquimod topical, wash your hands and wash the skin area to be treated. Allow the skin to dry for at least 10 minutes before applying the medicine. Always wash your hands after applying the medicine also.



Stop using imiquimod topical and call your doctor at once if you have a serious side effect such as flu symptoms such as fever, chills, body aches, tired feeling, swollen glands.

Log onto <http://www.orthorxaccess.com/> to activate your savings:

This offer is **only** valid for patients with commercial insurance



MOST ELIGIBLE COMMERCIALLY INSURED PATIENTS PAY NO MORE THAN

\$40 Copay for ZYCLARA

WHEN DEDUCTIBLE IS MET