

# Sharpen<sup>®</sup>

Powered by **Kixor<sup>®</sup>** Herbicide

**Group 14 Herbicide**

Doc id: 541790

ACCEPTED  
VIA NOTIFICATION  
LABEL NOT REVIEWED

Classified for  
"RESTRICTED USE"  
in New York State  
under 6NYCRR Part 326

April 14, 2015

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

**For use in selected agricultural crops**

**Active Ingredient:**

saflufenacil: N'-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3,6-dihydro-1(2H)-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide..... 29.74%

**Other Ingredients:** ..... 70.26%

**Total:** ..... 100.00%

Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate

**EPA Reg. No. 7969-278**

**EPA Est. No. 241-PR-002**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**Net Contents: 1 gallon**

67611782 NVA 2015-05-322-0042  
Product of U.S.A.

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

**BASF**  
We create chemistry

## FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give any liquid to the person.</li><li>• <b>DO NOT</b> give anything to an unconscious person.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li><li>• Call a poison control center for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information at 1-800-832-HELP (4357).

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (such as natural rubber)
- Protective eyewear such as face shield, goggles, or safety glasses

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Dis-

card clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for **applicators and other handlers** and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## Environmental Hazards

For terrestrial uses, **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

**Groundwater Advisory.** Saflufenacil has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory.** Saflufenacil may impact surface water due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks after application. A level, well-maintained buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall is forecast to occur within 48 hours.

## Endangered Species Protection Requirements

This product may have effects on federally listed threatened or endangered plant species or their critical habitat. When using this product, you must fol-

low the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

## Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at time of herbicide application.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all precautions and limitations in this label and the labels of products used in combination with **Sharpen® herbicide**. The use of **Sharpen** not consistent with this label can result in injury to crops, animals or persons. Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application in crops.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

**EXCEPTION:** If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as natural rubber  $\geq$  14 mils
- Shoes plus socks
- Protective eyewear

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

### Pesticide Storage

**DO NOT** use or store near heat or open flame. Store in original container in a well-ventilated area separately from fertilizer, feed, or foodstuffs. Avoid cross-contamination with other pesticides.

*(continued)*

## STORAGE AND DISPOSAL

*(continued)*

### Pesticide Disposal

Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### Container Handling

**Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### Triple rinse containers small enough to shake (capacity $\leq$ 5 gallons) as follows:

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

### Steps to take if material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

---

## Product Information

**Sharpen® herbicide** provides both contact burn-down and rate-dependent residual preemergence broadleaf weed control (refer to **Table 1** and **Table 2** for lists, respectively). It can be used in select field and row crops [chickpea (garbanzo beans), corn (field, pop, seed, and silage), cotton, edible pea, field

pea, lentils, rice, small grains, sorghum, soybean], fallow and postharvest croplands, for harvest aid/desiccation, and in noncropland areas. **Sharpen** does not control grass weeds and must be used sequentially or tank mixed with a grass herbicide for a complete weed control program. Refer to **Crop-specific Information** section for recommendations on herbicide tank mixtures or sequential programs.

Make burndown applications of **Sharpen** when broadleaf weeds are small and actively growing. An adjuvant is required with **Sharpen** for optimum burndown activity (refer to **Additives** section for details). Burndown activity may be slowed or reduced under cloudy and/or foggy or cooler weather conditions, or when weeds are growing under drought or other stress conditions. When targeting dense weed populations and/or larger broadleaf weeds, use a higher application rate within an application rate range and/or higher spray volumes. Angling nozzles forward (to 45 degrees) may improve penetration of denser weed canopies.

Residual preemergence applications of **Sharpen** must be activated by at least 1/2 inch of rainfall or sprinkler irrigation before weed seedling emergence. When **Sharpen** is not activated, a labeled post-emergence herbicide or cultivation may be needed to control weed escapes.

**Table 1. Broadleaf Weeds Controlled by a Burndown Application of Sharpen® herbicide**

Common Name	Scientific Name	C = Control S = Suppression		Maximum Height or Diameter (inches)
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C		6
Bedstraw, catchweed	<i>Galium aparine</i>	C		3
Beggarticks, hairy	<i>Bidens pilosa</i>	C		6
Beggarweed, Florida	<i>Desmodium tortuosum</i>	C		6
Bindweed, field	<i>Convolvulus arvensis</i>	S <sup>1</sup>		6
Buckwheat, wild	<i>Polygonum convolvulus</i>	C		3
Canola, volunteer (rapeseed)	<i>Brassica</i> spp.	C		6
Carpetweed	<i>Mollugo verticillata</i>	C		6
Chickweed, common	<i>Stellaria media</i>	S		3

(continued)

**Table 1. Broadleaf Weeds Controlled by a Burndown Application of Sharpen® herbicide (continued)**

Common Name	Scientific Name	Maximum Height or Diameter (inches)	
		C = Control S = Suppression	
Cocklebur, common	<i>Xanthium strumarium</i>	C	6
Cotton, volunteer <sup>2</sup>	<i>Gossypium hirsutum</i>	C	≤ 12 leaves
Cowcockle	<i>Vaccaria pyramidata</i>	C	4
Dandelion	<i>Taraxacum officinale</i>	S <sup>1</sup>	6
Eveningprimrose, cutleaf	<i>Oenothera laciniata</i>	C	4
Falseflax, smallseed	<i>Camelina microcarpa</i>	C	4
Filaree, redstem	<i>Erodium cicutarium</i>	S	3
Fleabane, hairy	<i>Conyza bonariensis</i>	C	6
Flixweed	<i>Descurainia sophia</i>	C	6
Groundcherry, cutleaf	<i>Physalis angulata</i>	C	6
Groundsel, common	<i>Senecio vulgaris</i>	C	4
Hawksbeard, narrowleaf <sup>2</sup>	<i>Crepis tectorum</i>	C	6
Hemlock, poison <sup>2</sup>	<i>Conium maculatum</i>	C	6
Henbit	<i>Lamium amplexicaule</i>	S	3
Horseweed (marestail)	<i>Conyza canadensis</i>	C	6
Knotweed, prostrate	<i>Polygonum aviculare</i>	C	3
Kochia	<i>Kochia scoparia</i>	C	1 to 3 Suppression of button/puffball stage at < 1-inch tall
Ladysthumb	<i>Polygonum persicaria</i>	C	6
Lambsquarters, common	<i>Chenopodium album</i>	C	6
Lambsquarters, narrowleaf	<i>Chenopodium pratericola</i>	C	6
Lettuce, prickly	<i>Lactuca serriola</i>	C	6
Mallow, common	<i>Malva neglecta</i>	C	6
Mallow, little (cheeseweed)	<i>Malva parviflora</i>	C	6
Mallow, Venice	<i>Hibiscus trionum</i>	C	6
Marestail (horseweed)	<i>Conyza canadensis</i>	C	6
Morningglory, entireleaf	<i>Ipomoea hederacea</i> <i>var. integruscula</i>	C	6
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C	6
Morningglory, palmleaf	<i>Ipomoea wrightii</i>	C	6
Morningglory, pitted	<i>Ipomoea lacunosa</i>	C	6
Morningglory, tall	<i>Ipomoea purpurea</i>	C	6

(continued)

**Table 1. Broadleaf Weeds Controlled by a Burndown Application of Sharpen® herbicide (continued)**

<b>Common Name</b>	<b>Scientific Name</b>	<b>C = Control S = Suppression</b>	<b>Maximum Height or Diameter (inches)</b>
Mustard, black	<i>Brassica nigra</i>	C	6
Mustard, tumble	<i>Sisymbrium altissimum</i>	C	6
Mustard, wild	<i>Sinapis arvensis</i>	C	6
Needles, Spanish <sup>2</sup>	<i>Bidens pilosa</i>	C	6
Nettle, burning	<i>Urtica urens</i>	C	4
Nightshade, black	<i>Solanum nigrum</i>	C	6
Nightshade, cutleaf	<i>Solanum triflorum</i>	C	6
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	C	6
Nightshade, hairy	<i>Solanum saccharoides</i>	C	6
Parthenium	<i>Parthenium hysterophorus</i>	C	6
Pennycress, field	<i>Thlaspi arvense</i>	C	6
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C	6
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C	6
Pigweed, smooth	<i>Amaranthus hybridus</i>	C	6
Puncturevine	<i>Tribulus terrestris</i>	C	6
Purslane, common	<i>Portulaca oleracea</i>	C	3
Pusley, Florida	<i>Richardia scabra</i>	S	3
Ragweed, common <sup>3</sup>	<i>Ambrosia artemisiifolia</i>	C	6
Ragweed, giant	<i>Ambrosia trifida</i>	C	6
Rocket, London <sup>2</sup>	<i>Sisymbrium irio</i>	C	6
Sesbania, hemp	<i>Sesbania exaltata</i>	C	4
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	C	6
Sida, prickly	<i>Sida spinosa</i>	C	6
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C	6
Sowthistle, annual	<i>Sonchus oleraceus</i>	C	6
Sowthistle, spiny	<i>Sonchus asper</i>	C	6
Sunflower, common	<i>Helianthus annuus</i>	C	6
Tansymustard, pinnate	<i>Descurainia pinnata</i>	C	6
Texasweed	<i>Caperonia palustris</i>	C	6
Thistle, Canada	<i>Cirsium arvense</i>	S <sup>1</sup>	6
Thistle, Russian	<i>Salsola kali</i>	C	3
Velvetleaf	<i>Abutilon theophrasti</i>	C	6

(continued)

**Table 1. Broadleaf Weeds Controlled by a Burndown Application of Sharpen® herbicide (continued)**

Common Name	Scientific Name	C = Control S = Suppression	Maximum Height or Diameter (inches)
Waterhemp <sup>3</sup>	<i>Amaranthus tuberculatus</i>	C	4
Willowweed	<i>Epilobium adenocaulon</i>	C	3

<sup>1</sup> Control of seedling stage and suppression of perennial growth stage.

<sup>2</sup> Not controlled in California.

<sup>3</sup> Populations of noted weeds that are known to be resistant to burndown applications of **Group 14/Group E** herbicides and will not be controlled by herbicides like **Sharpen**. See the **Resistance Management** section for practices to manage and minimize the impact of resistant weeds (e.g. tank mixes or alternation with other herbicide modes of action, crop rotation, and mechanical control).

**Table 2. Broadleaf Weeds Controlled with a Residual Preemergence Application of Sharpen® herbicide**

Common Name	Scientific Name	C = Control S = Suppression <sup>1</sup>
<b>Annual Broadleaf Weeds</b>		
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	C
Buckwheat, wild	<i>Polygonum convolvulus</i>	C
Burcucumber	<i>Sicyos angulatus</i>	S
Canola, volunteer (rapeseed), all types <sup>2</sup>	<i>Brassica</i> spp.	C
Carpetweed	<i>Mollugo verticillata</i>	C
Chickweed, common	<i>Stellaria media</i>	C
Cocklebur, common	<i>Xanthium strumarium</i>	C
Copperleaf, Virginia	<i>Acalypha virginica</i>	C
Galinsoga, smallflower	<i>Galinsoga parviflora</i>	C
Groundcherry, cutleaf	<i>Physalis angulata</i>	C
Horseweed (marestail)	<i>Conyza canadensis</i>	C
Jimsonweed	<i>Datura stramonium</i>	C
Kochia	<i>Kochia scoparia</i>	C
Ladysthumb	<i>Polygonum persicaria</i>	C
Lambsquarters, common	<i>Chenopodium album</i>	C
Mallow, Venice	<i>Hibiscus trionum</i>	C
Marestail (horseweed)	<i>Conyza canadensis</i>	C
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>integriscula</i>	C
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C



**Table 2. Broadleaf Weeds Controlled with a Residual Preemergence Application of Sharpen® herbicide** (continued)

Common Name	Scientific Name	C = Control S = Suppression <sup>1</sup>
<b>Annual Broadleaf Weeds</b>		
Morningglory, pitted	<i>Ipomoea lacunosa</i>	C
Morningglory, tall	<i>Ipomoea purpurea</i>	C
Mustard, wild	<i>Sinapis arvensis</i>	C
Nightshade, black	<i>Solanum nigrum</i>	C
Pennycress, field	<i>Thlaspi arvense</i>	C
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Pigweed, tumble	<i>Amaranthus albus</i>	C
Puncturevine	<i>Tribulus terrestris</i>	S
Purslane, common	<i>Portulaca oleracea</i>	C
Pusley, Florida	<i>Richardia scabra</i>	S
Ragweed, common	<i>Ambrosia artemisiifolia</i>	C
Ragweed, giant	<i>Ambrosia trifida</i>	C
Sida, prickly	<i>Sida spinosa</i>	C
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>	C
Starbur, bristly	<i>Acanthospermum hispidum</i>	C
Sunflower, common	<i>Helianthus annuus</i>	C
Texasweed	<i>Caperonia palustris</i>	C
Thistle, Russian	<i>Salsola kali</i>	C
Velvetleaf	<i>Abutilon theophrasti</i>	C
Waterhemp	<i>Amaranthus tuberculatus</i>	C

<sup>1</sup> Use **Sharpen** in tank mixes or sequential applications with other labeled herbicides that provide additional control of noted weeds.

<sup>2</sup> Not controlled in California.

## Mode of Action

**Sharpen** is a potent inhibitor of protoporphyrinogen-oxidase belonging to herbicide mode of action **Group 14 (WSSA)/Group E (HRAC)**. **Sharpen** is rapidly absorbed by roots and foliage. Following inhibition of protoporphyrinogen-oxidase, plant death

is the result of membrane damage. Under active growing conditions, susceptible emerged weeds usually develop chlorotic and necrotic injury symptoms within hours and die within a few days. Susceptible emerging weed seedlings will usually die as they reach the soil surface or shortly after emergence.

## Resistance Management

While weed resistance to protoporphyrinogen-oxidase inhibiting herbicides is relatively infrequent, populations of resistant biotypes are known to exist. Resistance management practices include:

1. Following labeled application rate and weed growth stage recommendations
2. Avoiding repeated applications of herbicides with the same mode of action
3. Utilizing tank mixes and sequential applications with other effective herbicides possessing different modes of action
4. Using crop rotation so that crop competition, tillage or herbicides with alternative modes of action can be used to control weed escapes

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended.

## Crop Tolerance

Crops listed on this label are tolerant to **Sharpen**<sup>®</sup> herbicide when applied according to label directions as a preplant to preemergence treatment and under normal environmental conditions. Crop injury may occur under stressful growing conditions (e.g. low soil fertility, seedling disease, extreme hot or cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought).

Severe crop injury will result if **Sharpen** is applied postemergence (over the top) to any crop, except to rice.

---

## Application Instructions

**Sharpen** may only be applied before crop emergence, except for harvest aid/desiccation uses and postemergence in rice.

## Application Rates

Application rates of **Sharpen** may vary depending on soil texture and organic matter. Refer to **Table 3** for soil texture groups used in this label.

**Table 3. Soil Texture Groups**

Coarse	Medium	Fine
Sand	Silt	Sandy clay
Loamy sand	Silt loam	Silty clay
Sandy loam	Loam	Silty clay loam
	Sandy clay loam	Clay loam
		Clay

Refer to the **Crop-specific Information** section for specific application rates, timings, and the restrictions and limitations by crop and use pattern.

In California, **DO NOT** apply more than 2.0 fl ozs/A of **Sharpen** in a single application.

## Application Methods and Equipment

**Sharpen** may be applied by ground or air. Thorough spray coverage is required for optimum broadleaf weed control and can be improved with proper adjuvant, nozzle and spray volume selection.

Use and configure application equipment to provide an adequate spray volume, an accurate and uniform distribution of spray droplets over the treated area, and to avoid spray drift to nontarget areas. Equipment should be adjusted to maintain continuous agitation during spraying with good mechanical or bypass agitation. Avoid overlaps that will increase rates above the use rates specified in this label.

**Sharpen** may be applied using either water or sprayable fluid nitrogen fertilizer solutions as the spray carrier. Additionally, **Sharpen** may be impregnated on and applied with dry bulk fertilizer.

## Aerial Application Requirements

**Water Volume.** Use 3 or more gallons of water per acre for weed control applications. Use a minimum of 5 gallons of water per acre for harvest aid/desiccation applications.

**DO NOT** apply aerially in California.

The following measures must be followed to reduce the potential of spray drift to nontarget areas from aerial applications:

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the fixed wingspan or 90% of rotor blade diameter.
2. Use low-drift nozzles such as straight-stream nozzles (D-8 or larger). **DO NOT** use nozzles producing a mist droplet spray.
3. Nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.
4. Without compromising aircraft safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants.
5. **DO NOT** apply during periods of temperature inversions or stable atmospheric conditions.
6. Avoid potential adverse effects to nontarget areas by maintaining a 160-foot buffer between the point of direct application and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, shrub lands, and crop lands).

## Ground Application Requirements

**Spray Carrier Volume.** Use 5 or more gallons of water per treated acre or 20 or more gallons of sprayable fluid nitrogen fertilizer per treated acre for weed control applications. Thorough spray coverage is required for control of emerged broadleaf weeds. High populations and/or variations in size can prevent adequate spray coverage. Controlling fall-germinated weeds in the spring (e.g. horseweed/marestail) will also require thorough spray coverage. Use higher spray volumes (e.g. 15 to 20 gallons of water per acre) in these situations to increase spray coverage and optimize burndown activity. Use a minimum of 5 to 10 gallons of water per acre for harvest aid/desiccation applications.

The following measures must be followed to reduce the potential of spray drift to nontarget areas from ground applications:

1. Apply this product using nozzles which deliver **medium-to-coarse spray droplets** as defined by ASAE standard S-572 and as shown in nozzle manufacturer's catalogs. Flat-fan nozzles are rec-

- ommended for burndown applications while flood-jet type nozzles are recommended for residual soil surface applications. Nozzles that deliver coarse spray droplets may be used to reduce spray drift provided spray volume per acre (GPA) is increased to maintain coverage of target (i.e. weeds or soil surface). **DO NOT** use nozzles that produce fine (e.g. cone) spray droplets. In California, nozzles must be affixed to spray no higher than 20 inches above the spray target (e.g. top of weed foliage).
2. Apply this product only when the potential for drift to adjacent nontarget areas is minimal (e.g. when the wind is **10 MPH or less and is blowing away** from sensitive areas). **DO NOT** apply during periods of temperature inversions or stable atmospheric conditions.
  3. Avoid potential adverse effects to nontarget areas by maintaining a 75-foot buffer between the application area and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, shrub lands, and crop lands).

## Ground Application (dry bulk fertilizer)

**Sharpen® herbicide** may be impregnated or coated onto dry bulk granular fertilizer carriers for residual soil surface applications. Impregnation or coating may be conducted by either in-plant bulk or on-board systems. Perform the mixing operation in well-ventilated areas.

Addition of a drying agent may be necessary if the fertilizer and herbicide blend is too wet for uniform application due to high humidity, high urea concentration, or low fertilizer use rate. Slowly add the drying agent to the blend until a flowable mixture is obtained. Drying agents are not recommended for use with on-board impregnation systems.

Under some conditions, fertilizer impregnated with **Sharpen** may clog air tubes or deflector plates on pneumatic application systems. Mineral oil may be added to **Sharpen** before blending with fertilizer to reduce plugging. **DO NOT** use drying agents when

mineral oil is used. To avoid separation of **Sharpen® herbicide** and mineral oil mixes in cold temperatures, keep mixture heated or agitated before blending with fertilizer. Mineral oil may be used with in-plant blending stations or with on-board injection systems. Generally, fertilizer application rates of at least 200 lbs to 700 lbs per acre of herbicide and fertilizer blend will provide adequate distribution or coverage of **Sharpen** across the soil surface. Application must be made uniformly to the soil to prevent possible crop injury and offer satisfactory weed control. Impregnated fertilizer spread at half rate and overlapped to obtain a full rate will offer a more uniform distribution. A shallow (< 2 inches) incorporation is desirable for improved weed control. Deeper incorporation will dilute the herbicide layer near the soil surface and may result in unsatisfactory weed control.

Use the following formula to determine the herbicide rate when using dry bulk fertilizer applications:

$$\frac{\text{fl ozs herbicide per acre}}{\text{pounds fertilizer per acre}} \times 2000 = \frac{\text{fl ozs herbicide}}{\text{per ton of fertilizer}}$$

## Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions, followed by triple rinsing the equipment before and after applying this product.

## Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site, especially onto nontarget areas. The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The applicator should be familiar with and take into account the information covered in the following spray drift reduction advisory information.

**Controlling Droplet Size.** The most effective way to reduce drift potential is to apply the largest droplets that provide sufficient coverage and control.

**Volume.** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure. DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles.** Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Type.** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets.

**Swath Adjustment.** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

**Wind.** Drift potential is lowest between wind speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. If applying at wind speeds less than 3 mph, the applicator must determine if:

1. Conditions of temperature inversion exist, or
2. Stable atmospheric conditions exist at or below nozzle height.

**DO NOT** make applications into areas of temperature inversions or stable atmospheric conditions.

**NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Wind Erosion.** Avoid treating powdery, dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

## Additives

For optimum burndown or harvest aid/desiccation activity with **Sharpen® herbicide**, an adjuvant system must be used that includes the following:

Adjuvant	Rate
methylated seed oil <sup>1</sup> (MSO)	1 gal/100 gals <sup>2</sup> (1% v/v)
<b>PLUS</b>	<b>PLUS</b>
ammonium sulfate (AMS)	8.5 to 17 lbs/100 gals (1% to 2% w/v)
<b>or</b>	<b>or</b>
urea ammonium nitrate (UAN)	1.25 to 2.5 gals/100 gals (1.25% to 2.5% v/v)

<sup>1</sup> MSO-based adjuvant **MUST** contain at least 60% methylated seed oil. Poor performance may occur with adjuvants containing less than 60% methylated seed oil.

<sup>2</sup> **DO NOT** use less than 1 pint/A of MSO with low-volume (< 12.5 gallons per acre) aerial or ground applications.

Refer to the **Crop-specific Information** section for specific adjuvant requirements for certain crop uses. The use of AMS fertilizer is highly recommended when mixing **Sharpen** with glyphosate-based herbicides.

**DO NOT** use a nonionic surfactant (NIS) as a substitute for MSO, or poor performance on broadleaf weeds or for desiccation will occur.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant.

**DO NOT** add acidifying agents to the spray tank when applying **Sharpen**.

## Tank Mixing Information

**Sharpen** may be tank mixed with 1 or more registered herbicide products according to the specific tank mixing instructions in this label and respective product labels. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Always follow the most restrictive label use directions. Refer to **Crop-specific Information** section for tank mixing details.

Tank mixtures with contact herbicides (e.g. carfentrazone, paraquat) may reduce the burndown activity of **Sharpen**.

### Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

1. For 20 gallons per acre spray volume, use 3.3 cups (800 mL) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.
2. Add components in the sequence indicated in the mixing order using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre.
3. Always cap the jar and invert 10 cycles between component additions.
4. When the components have all been added to the jar, let the solution stand for 15 minutes.
5. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, or fine particles that precipitate to the bottom, or thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, **DO NOT** mix the ingredients in the same tank.

## Mixing Order

1. **Water** - Fill tank 1/2 to 3/4 full with clean water and start agitation.
2. **Agitation** - Maintain agitation throughout mixing.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-soluble additives** (including dry and liquid fertilizers such as ammonium sulfate or urea ammonium nitrate)
6. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
7. **Water-soluble products**
8. Emulsifiable concentrates (including methylated seed oil adjuvants)
9. **Remaining quantity of water**

Maintain agitation throughout application until spraying is completed. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

---

## Use Precautions

---

- **Maximum seasonal use rate** - Refer to **Crop-specific Information** section for maximum cropping seasonal application use rates for each crop and use pattern. A cropping season is defined as the period following harvest of the preceding crop through the harvest of the planned or current crop.
- Except for labeled harvest aid/desiccation and postemergence to rice uses, **DO NOT** apply **Sharpen® herbicide** after crop emergence or severe crop injury will occur.
- **Rainfastness** - **Sharpen** is rainfast 1 hour after application. Burndown activity may be reduced if rain or irrigation occurs within 1 hour of application.
- **DO NOT** contaminate irrigation ditches or water used for domestic purposes.
- **DO NOT** apply **Sharpen** through any type of irrigation system (e.g. chemigation).
- **Sharpen is not for sale, distribution, or use in Nassau and Suffolk counties in New York State.**

## Crop Rotation and Emergency Replanting Intervals

Use **Table 4** to determine the proper interval between **Sharpen® herbicide** application and planting of rotational crops or replanting after crop failure (because of environmental factors such as drought, frost, or hail, etc.). Be sure to determine the rotational crop interval for tank mix products and utilize the most restrictive interval of all products applied.

**Table 4. Rotational Crop Planting and Emergency Replanting Intervals by Sharpen Application Rate**

Crop	Sharpen Rate (fl ozs/A)					
	1.0	2.0	3.0	4.0	5.0	6.0
	Rotational Crop Interval (months after application) <sup>1</sup>					
Corn	0	0	0	0	0	0
Corn, sweet	0.5	1	2	3	4	4
Sorghum	0	0	0	0	1	1
Small grains <sup>2</sup>	0	0	0	0	3	3
Rice	0	0	4	4	4	4
Chickpea	0	0	2*	4	6	6
Edible pea	0	1	3	4	6	6
Field pea, dry	0	0	2	4	6	6
Soybean <sup>3</sup>	0 to 1	1 to 1.5	2 to 3	4	6	6
Lentil	0	1	3	4	6	6
Cotton <sup>3</sup>	1.5	3	4	6	6	9
Sugarbeet	4	5	6	7	8	9
Sunflower	4	5	6	7	8	9
Other Crops	4	5	6	7	8	9

<sup>1</sup> **DO NOT** include time when the soil is frozen.

<sup>2</sup> Small grains are defined in **Crop-specific Information** section of this label. For other small grains, use the rotational crop interval for **Other Crops**.

<sup>3</sup> The planting interval for these crops and rates is further defined in the respective **Crop-specific Information** section of this label. Use the longer interval within listed ranges for indicated crops grown on coarse-texture soils with organic matter less than 2.0%.

\*Interval is 0 months in Idaho, Oregon, and Washington.

## Crop-specific Information

This section provides use directions for **Sharpen® herbicide** in specific crops. Be sure to read product information, mixing, application, weeds controlled and adjuvant instructions in preceding sections of the label. Read and follow tank mix product labels for restrictions, precautions, instructions, and rotational crop restrictions.

Depending on specific crop application directions, **Sharpen** may be applied for burndown control of emerged broadleaf weeds and/or residual control of germinating broadleaf weeds (refer to **Table 1** and **Table 2** for lists of weeds controlled) before crop planting (preplant and/or preseed) or after planting but before crop emergence (preemergence).

Thorough spray coverage is required for control of emerged broadleaf weeds. High populations and/or variations in size can prevent adequate spray coverage. Controlling fall-germinated weeds in the spring (e.g. horseweed/marestail) will also require thorough spray coverage. Use higher spray volumes (e.g. 15 to 20 gallons of water per acre) in these situations to increase spray coverage and optimize burndown activity.

### Field Corn (grain, seed, silage) and Popcorn

**Sharpen** may be applied preplant surface, preplant incorporated, or preemergence to corn for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled). Corn in this label refers to field corn (grown for grain, seed, or silage) and popcorn. Before applying **Sharpen** to seed corn or popcorn, verify the selectivity of **Sharpen** on your inbred line or hybrid with your local seed company (supplier) to help avoid potential injury to sensitive inbreds or hybrids.

### Application Rates

**Sharpen** can be applied as part of a planned sequential (two-pass) weed control program.

**Sharpen** use rates applied as the residual component of a planned sequential (two-pass) program (see **Table 5**) will provide control or suppression of

listed weeds (**Table 1**) through early to mid-season. For full-season weed control, apply a labeled postemergence treatment such as **Status® herbicide** plus glyphosate as the sequential component.

**Table 5. Residual Preemergence Rates of Sharpen in a Planned Sequential Program<sup>1</sup> in Field Corn and Popcorn**

Soil Texture <sup>2</sup>	Rate by Soil Texture (fl ozs/A)
Coarse	2.0 to 2.5
Medium	2.5 to 3.0
Fine	3.0 to 3.5

<sup>1</sup> Application rates in **Table 5** will eliminate early season broadleaf weed interference until cultivation or a labeled postemergence herbicide is applied.

<sup>2</sup> Refer to **Table 3** for definition of soil texture groups.

## Application Timings

### Early Preplant Surface Application (15 to 30 days before planting)

Early preplant surface applications are not recommended on coarse soils, in areas where average annual rainfall (or rainfall plus irrigation) typically exceeds 40 inches, or for popcorn. Cultivation or a labeled postemergence herbicide application may still be required under certain conditions for complete weed control.

Early preplant surface applications may be applied as part of a split application program where applications are made as part of the application timings described in this label. However, the cumulative total of sequential application rates must not exceed the maximum labeled rate for a given soil texture.

### Preplant Surface and Preplant Incorporated Applications (up to 14 days before planting)

**Sharpen** can be applied at use rates specified in **Table 5** to the soil surface or incorporated up to 14 days before planting on all soil types. For preplant incorporated applications, apply **Sharpen** and



incorporate into the upper soil surface (1 to 2 inches). Use a harrow, rolling cultivator, field cultivator or other implement capable of providing uniform shallow incorporation. Avoid deeper incorporation or reduced weed control may result.

## Preemergence Surface Application

Apply **Sharpen**® herbicide at use rates specified in **Table 5** as a broadcast spray to the soil surface after planting and before crop emergence. **Sharpen** must be applied before crop emergence or injury will occur.

## Burndown plus Residual Weed Control

In addition to residual broadleaf weed control obtained at any of the application timings described above, **Sharpen** will also provide burndown of emerged broadleaf weeds listed in **Table 1**. An adjuvant system (refer to **Additives** section for details) is required for optimum burndown activity. Burndown control of emerged grasses and/or additional broadleaf weeds not listed on the label will require a tank mix with another herbicide (such as glyphosate).

## Burndown Weed Control Only

If limited or no residual broadleaf weed control is desired, **Sharpen** can be applied at 1.0 fl oz/A (all soil types) with an adjuvant system any time before corn emergence to provide burndown of broadleaf weeds listed in **Table 1**. A burndown application of **Sharpen** can be followed by residual rates of **Sharpen** (**Table 5**) or **Verdict**® herbicide. Separate sequential applications by at least 14 days. However, **DO NOT** exceed the cropping seasonal maximum cumulative amount per acre of saflufenacil from all product sources per cropping season.

**Enhanced burndown in seed corn.** Apply **Sharpen** preplant surface or preemergence at 1.0 to 2.0 fl ozs/A with an adjuvant system for enhanced burndown broadleaf weed control in seed corn before crop emergence.

**DO NOT** exceed 1.0 fl oz/A on coarse soils. A sequential application of **Sharpen** may be made with a minimum of 30 days between applications.

**DO NOT** apply more than a maximum cumulative amount of 4.0 fl ozs/A of **Sharpen** per cropping season in seed corn.

**State-specific Use in California.** Apply **Sharpen** early preplant through preemergence at 1.0 to 2.0 fl ozs/A with an adjuvant system for enhanced burndown broadleaf weed control before crop emergence. Separate sequential applications by at least 14 days. **DO NOT** apply greater than 2.0 fl ozs/A in a single application.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply **Sharpen** after corn emergence or severe crop injury will occur.
- **Sharpen** use may result in delayed corn emergence and stunting under certain environmental conditions including cool temperatures, excessive rainfall/irrigation, and/or persistent wet soil conditions occurring after application.
- Ensure that the corn seed row is closed. Soil conditions that cause poor seed furrow closure and coverage may result in delayed corn emergence or stunting.
- **DO NOT** apply **Sharpen** at greater than 1.0 fl oz/A where an at-planting application of an organophosphate (OP) or carbamate insecticide(s) is planned and/or has occurred because severe injury may result.

**EXCEPTION:** **Sharpen** may be applied when **Aztec**® 2.1% granular insecticide, **Aztec**® 4.67 G granular insecticide, **Fortress**® 5G granular insecticide, or **SmartChoice**™ 5G granular insecticide is applied at planting as a band, T-band, or in-furrow. **Sharpen** may be applied with all other classes of at-planting insecticides including pyrethroids, neonicotinoids, and fipronil.

- **DO NOT** apply more than a maximum cumulative amount of 6.0 fl ozs/A of **Sharpen** (0.134 lb ai/A of saflufenacil) per cropping season.
- **DO NOT** apply more than a maximum cumulative amount of 0.134 lb ai/A of saflufenacil per cropping season in corn from all product sources.

- Corn forage and silage can be harvested, fed, or grazed 80 or more days after application.
- Refer to **Table 4** for rotational crop planting intervals.

## Tank Mixtures

**Sharpen**® herbicide may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Clarity**® herbicide
- **Outlook**® herbicide
- **Prowl**® H<sub>2</sub>O herbicide
- **Status**® herbicide
- **Verdict**® herbicide
- **Zidua**® herbicide
- atrazine
- glyphosate (e.g. **Roundup**® herbicide)
- **Harness**® herbicide
- **Harness**® Xtra herbicide

### Cotton

Use **Sharpen** as an early preplant burndown treatment before planting cotton.

Not for use in cotton in California.

## Application Rates and Timings

Apply **Sharpen** as a broadcast spray at 1.0 fl oz/A plus recommended adjuvants (refer to **Additives** section for details) for the control of actively growing broadleaf weeds (refer to **Table 1** for list of weeds controlled). Wait to plant cotton until at least **42 days** and an accumulation of 1 inch of rainfall and/or irrigation occurring after application to avoid crop injury. In geographic areas with average annual rainfall less than 25 inches, the 42-day preplant interval is required after the accumulation of 1 inch of rainfall and/or irrigation. **DO NOT** apply to coarse soils classified as sand with less than 1.5% organic matter or cotton injury may occur.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply more than a maximum cumulative amount of 2.0 fl ozs/A of **Sharpen** per cropping season.

- **DO NOT** apply **Sharpen** with other **Group 14/Group E** herbicides (such as flumioxazin) as a tank mix or sequential application within 30 days of planting because crop injury may result.
- Use the most restrictive preplant interval with tank mixes of other cotton burndown herbicides.
- Cotton gin byproducts may be fed to livestock.
- Refer to **Table 4** for rotational crop planting intervals.

## Tank Mixtures

Broad-spectrum burndown control of grasses and/or additional broadleaf weeds will require a tank mix with a herbicide such as glyphosate. **Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following cotton burndown herbicide products:

- **Clarity**
- **Distinct**
- **Prowl H<sub>2</sub>O**
- glyphosate (e.g. **Roundup**)

### Fallow and Postharvest

**Sharpen** may be used as a burndown treatment to control broadleaf weeds at any time of the year during the fallow period following crop harvest and before the following crop is planted. **Sharpen** may also be used for specific postharvest uses to burndown the remaining foliage after crop harvest.

## Application Rates and Timings

Apply **Sharpen** as a broadcast burndown spray at 1.0 to 2.0 fl ozs/A plus recommended adjuvants (refer to **Additives** section for details). Best product performance is obtained when broadleaf weeds are small and actively growing (refer to **Table 1** for list of weeds controlled). Thorough coverage of existing weeds is essential and higher spray volumes may be needed for best performance.

Sequential applications may be made with a minimum of 14 days between applications; but **DO NOT** exceed a maximum seasonal cumulative amount of 6.0 fl ozs/A of **Sharpen** per cropping season.

For residual broadleaf weed control, **Sharpen® herbicide** may be applied at 2.0 to 4.0 fl ozs/A.

Specific rotational crop planting intervals must be observed between an application of **Sharpen** and planting of the following crop (see **Table 4** for rotational crop planting intervals).

**State-specific Use in California.** Apply **Sharpen** as a broadcast burndown spray at 1.0 to 2.0 fl ozs/A with the recommended adjuvants. **DO NOT** apply greater than 2.0 fl ozs/A in a single application.

**Postharvest use on tomato vines.** Apply **Sharpen** as a broadcast burndown spray at 1.0 to 2.0 fl ozs/A plus recommended adjuvants (refer to **Additives** section for details). Thorough spray coverage of existing tomato vines is essential and higher spray volumes may be needed for best performance. **DO NOT** apply before or during tomato fruit harvest. Not for use on tomato vines in California.

## Tank Mixtures

Broad-spectrum burndown control of grasses and/or additional broadleaf weeds will require a tank

mix with another herbicide. **Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Clarity® herbicide**
- **Distinct® herbicide**
- glyphosate (e.g. **Roundup® herbicide**)

## Harvest Aid/Desiccation

**Sharpen** may be used for harvest aid/desiccation in the crops listed in the table following. Uniformly apply **Sharpen** as a broadcast spray by air or ground. Ground application is recommended at a minimum spray volume of 10 gallons/A. Aerial application is recommended at a minimum spray volume of 5 gallons/A. Thorough spray coverage and an MSO plus ammonium-based adjuvant system (refer to the **Additives** section) are required for optimum desiccation activity.

**Sharpen** may be applied in a single application or sequential applications.

Refer to **Table 4** for rotational crop planting intervals.

**Table 6. Crops for Harvest Aid/Desiccation**

Crop	Application Timing	Sharpen® herbicide Use Rate (fl ozs/A)	PHI* (days)	Crop-specific Restrictions and Limitations
<p><b>Dry Edible Beans</b> including:</p> <p><i>Phaseolus</i> spp. (all types) Garbanzo bean (Chickpea)</p>	<p>Spray over the top of dry edible beans that have reached physiological maturity (beans have at least 80% yellow/brown pods and no more than 30% of leaves still green for vine-type beans, and no more than 40% of leaves still green for bush-type beans; or according to Extension Service recommendations in the use area).</p> <p>Allow up to 7 days for optimum desiccation effect depending on environmental conditions.</p>	<p>Single application: 1.0 to 2.0</p> <p>Maximum cumulative amount per cropping season for desiccation uses: 2.0</p>	<p>2 (dry edible beans)</p>	<p><b>DO NOT</b> apply on dry edible beans grown for seed production.</p> <p><b>DO NOT</b> graze or feed desiccation-treated hay or straw to livestock.</p> <p>Not for use for garbanzo bean (chickpea) desiccation in California.</p>
<p><b>Dry Peas</b> including:</p> <p>Dry field peas</p>	<p>Spray over the top of dry peas that have reached physiological maturity (peas with at least 80% yellow/brown pods and no more than 30% of leaves still green for vine-type peas, and no more than 40% of leaves still green for bush-type peas; or according to Extension Service recommendations in the use area).</p> <p>Allow up to 10 days for optimum desiccation effect depending on environmental conditions.</p>	<p>Single application: 1.0 to 2.0</p> <p>Maximum cumulative amount per cropping season for desiccation uses: 2.0</p>	<p>3 (dry peas)</p>	<p><b>DO NOT</b> apply on dry peas grown for seed production.</p> <p>Desiccation-treated pea vines may be grazed or fed to livestock.</p> <p>Not for use for dry peas desiccation in California.</p>

(continued)

**Table 6. Crops for Harvest Aid/Desiccation** (continued)

Crop	Application Timing	Sharpen® herbicide Use Rate (fl ozs/A)	PHI* (days)	Crop-specific Restrictions and Limitations
<p><b>Oilseeds Canola (Rapeseed) Subgroup 20A</b></p> <p>including:</p> <ul style="list-style-type: none"> <li>Crambe</li> <li>Gold-of-pleasure (Camelina)</li> <li>Juncea</li> <li>Mustard seed</li> <li>Rapeseed (Canola)</li> </ul>	<p>Spray over the top of crop(s) that have reached physiological maturity [seeds in the middle pods have started to turn in color for canola (rapeseed), Juncea, and mustard; or according to Extension Service recommendations in the use area for other crops].</p> <p>Allow up to 7 days for optimum desiccation effect depending on environmental conditions.</p>	<p>Single application: 1.0 to 2.0</p> <p>Maximum cumulative amount per cropping season for desiccation uses: 2.0</p>	<p>3 (seed)</p>	<p><b>DO NOT</b> apply on oilseed crops grown for seed production.</p> <p>Not for use for oilseeds canola (rapeseed) desiccation in California.</p>
<p><b>Oilseeds Cottonseed Subgroup 20C</b></p> <p>including:</p> <ul style="list-style-type: none"> <li>Cotton (Cottonseed)</li> </ul> <p>including:</p> <ul style="list-style-type: none"> <li>Spindle pick harvested cotton</li> <li>Stripper harvested cotton</li> </ul>	<p>Spray over the top of cotton that has reached physiological maturity (according to local State Agricultural Extension Service guidelines, such as nodes above cracked boll, accumulated heat units after cutout, or at least 60% to 70% boll opening).</p> <p>Allow up to 10 days for optimum desiccation and defoliation effect depending on environmental conditions.</p> <p>Large plant size, dense canopy, and environmental conditions not conducive for defoliation may require a second defoliation application 5 to 7 days later.</p>	<p>Single application: 0.5 to 2.0</p> <p>Maximum cumulative amount per cropping season for defoliation/ desiccation uses: 2.0</p>	<p>5 (cotton)</p>	<p><b>DO NOT</b> apply on cotton grown for seed production.</p> <p><b>Sharpen</b>-treated gin trash may be fed to livestock.</p> <p>Not for use for cotton desiccation in California.</p>

(continued)

**Table 6. Crops for Harvest Aid/Desiccation** (continued)

Crop	Application Timing	Sharpen® herbicide Use Rate (fl ozs/A)	PHI* (days)	Crop-specific Restrictions and Limitations
<p><b>Oilseeds Sunflower Subgroup 20B</b> including:</p> <p>Safflower Sunflower</p>	<p>Spray over the top of crop(s) that has reached physiological maturity (seed moisture less than 36% for sunflower; when heads are fully mature for safflower; or according to Extension Service recommendations in the use area). For many sunflower varieties, the back of the sunflower heads are yellow and the bracts are turning brown at this timing. Allow up to 7 days for optimum desiccation effect depending on environmental conditions.</p>	<p>Single application: 1.0 to 2.0</p> <p>Maximum cumulative amount per cropping season for desiccation uses: 4.0</p>	<p>7 (seed)</p>	<p><b>DO NOT</b> apply on oilseed crops grown for seed production.</p> <p><b>DO NOT</b> use after multiple minor frosts or single significant frost event.</p> <p>Not for use for safflower desiccation in California.</p>
<p><b>Soybean</b></p>	<p>Spray over the top of soybeans that have reached physiological maturity (all pods and seeds have no more green color).</p> <p><b>Indeterminate Varieties</b> Greater than 65% brown pods and greater than 70% leaf drop or when seed moisture is 30% or less.</p> <p><b>Determinate Varieties</b> Beans are fully developed, greater than 50% leaf drop, and remaining leaves are yellowing.</p> <p>Allow up to 10 days for optimum desiccation effect depending on environmental conditions.</p>	<p>Single application: 1.0 to 2.0</p> <p>Maximum cumulative amount per cropping season for desiccation uses: 2.0</p>	<p>3 (soybean seed)</p>	<p><b>DO NOT</b> apply on soybean grown for seed production.</p> <p><b>DO NOT</b> graze or feed desiccation-treated hay or straw to livestock.</p>

\* PHI = preharvest interval

## Tank Mixtures Cotton

**Sharpen® herbicide** may be tank mixed with other cotton defoliant products including, but not limited to: ethephon (such as **Prep™ brand Ethephon for cotton and tobacco**), ethephon plus AMADS (such as **CottonQuik® cotton harvest aid/defoliant**), ethephon plus cyclanilide (**Finish® 6 Pro harvest aid for cotton**), sodium chlorate, thidiazuron (such as **Dropp® SC cotton defoliant**), thidiazuron plus diuron (such as **GinStar® EC cotton defoliant**), tribufos (such as **Def® 6 emulsifiable defoliant** or **Folex® 6 EC cotton defoliant**), or glyphosate (e.g. **Roundup® herbicide**), or paraquat (e.g. **Gramoxone® herbicide**). Refer to the other product's label for restrictions on tank mixing, precautions, and rotational restrictions. The most restrictive labeling applies to tank mixes.

## All Other Crops

Apply **Sharpen** with a labeled rate of glyphosate (e.g. **Roundup**) for additional preharvest weed control. **Sharpen** also may be tank mixed with other registered harvest aid/desiccant products. Read and follow the applicable restrictions and limitations and directions for use on the glyphosate and other tank mix product labels, and to confirm they are labeled for the specific crop to be desiccated. The most restrictive labeling applies to tank mixes.

## Legume Vegetables [chickpea, edible pea, field pea, and lentil]

**Sharpen** may be applied preplant surface, preplant incorporated, and/or preemergence in legume crops specified in this section for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled).

## Application Rates and Timings

See the specific application rates and timings recommendations as they vary by legume crop. With burndown applications, an adjuvant system (refer to

**Additives** section for details) is required for optimum burndown activity.

Before applying **Sharpen** to any of the specified legume crops, verify the selectivity of **Sharpen** on your variety with your seed company (supplier) to help avoid potential injury to sensitive varieties.

Use of **Sharpen** may result in delayed crop emergence and stunting under certain environmental conditions including cool temperatures, excessive rainfall/irrigation, and/or persistent wet soil conditions occurring after application.

## Chickpea (garbanzo bean)

**Sharpen** is for use in all types of chickpeas.

**Burndown.** Apply **Sharpen** early preplant through preemergence at 1.0 fl oz/A for burndown broadleaf weed control before crop emergence. Sequential applications may be made with a minimum of 14 days between applications.

**Enhanced Burndown.** Apply **Sharpen** early preplant through preemergence at 2.0 fl ozs/A for enhanced burndown broadleaf weed control before crop emergence. Ensure that the seed row is closed. Soil conditions that cause poor seed furrow closure and coverage may result in delayed crop emergence or stunting. A sequential application of **Sharpen** may be made with a minimum of 30 days between applications.

**DO NOT** apply more than a maximum cumulative amount of 4.0 fl ozs/A of **Sharpen** (0.089 lb ai/A saflufenacil) per cropping season in chickpeas.

## Edible Peas

**Sharpen** is for use **ONLY** on the following edible peas:

- Edible-podded peas (sugar snap pea)
- Succulent peas (English pea, garden pea, green pea, marrowfat pea)

**State-specific Use in Idaho, Illinois, Iowa, Michigan, Minnesota, New York, Oregon, Washington, and Wisconsin.** Apply **Sharpen** preplant incorporated or preemergence at 0.75 fl oz/A in English or sugar snap peas for residual suppression of the following broadleaf weeds:

- Black nightshade
- Common lambsquarters
- Redroot pigweed
- Velvetleaf

**Preplant incorporated** - Apply **Sharpen**® herbicide up to one week before planting. **DO NOT** incorporate deeper than 3 inches.

**Preemergence** - Apply **Sharpen** up to 3 days after planting but before cracking stage or emergence, or severe crop injury will occur. Ensure that the seed row is closed. Soil conditions that cause poor seed furrow closure and coverage may result in delayed crop emergence or stunting.

A sequential application of **Sharpen** may be made with a minimum of 30 days between applications. **DO NOT** apply more than a maximum cumulative amount of 2.0 fl ozs/A of **Sharpen** (0.045 lb ai/A saflufenacil) per cropping season in edible peas.

## Field Peas

**Sharpen** is for use **ONLY** on dry field peas including Austrian winter peas.

**Burndown.** Apply **Sharpen** early preplant through preemergence at 1.0 fl oz/A for burndown broadleaf weed control before crop emergence. Sequential applications may be made with a minimum of 30 days between applications.

**Enhanced Burndown.** Apply **Sharpen** early preplant through preemergence at 2.0 fl ozs/A for enhanced burndown broadleaf weed control before crop emergence. Ensure that the seed row is closed. Soil conditions that cause poor seed furrow closure and coverage may result in delayed crop emergence or stunting. A sequential application of **Sharpen** may be made with a minimum of 30 days between applications.

**DO NOT** apply more than a maximum cumulative amount of 4.0 fl ozs/A of **Sharpen** (0.089 lb ai/A saflufenacil) per cropping season in dry field peas.

## Lentils

**Sharpen** may be applied to green-type and red-type lentils. **DO NOT** apply **Sharpen** to Spanish brown lentils.

## State-specific Use in Minnesota, Montana, North Dakota, South Dakota, and Wyoming.

Apply **Sharpen** early preplant to preemergence at 0.75 fl oz/A for burndown broadleaf weed control before crop emergence. For only limited residual activity on broadleaf weeds. **Sharpen** may also be applied preplant incorporated or preemergence at 0.75 to 2.0 fl ozs/A. Sequential applications may be made with a minimum of 14 days between applications. **DO NOT** apply more than a maximum cumulative amount of 2.0 fl ozs/A of **Sharpen** (0.045 lb ai/A saflufenacil) per cropping season in lentils.

**DO NOT** use **Sharpen** for lentils grown in any other state including California.

**Use Advisory for Lentils.** Lentil injury may be observed depending on factors including rainfall, soil type, seeding depth, and variety. Rainfall shortly after **Sharpen** application can result in slight injury to the crop. Lentils will be more susceptible to injury from **Sharpen** on coarse-texture and low-organic matter soils. Injury will usually appear as leaf tissue necrosis on the outer edges of the leaves. Lentils will grow out of injury symptoms, and yield will not be impacted at labeled rates. Soil residual herbicides may increase the sensitivity of lentils to **Sharpen** and should not be combined as a tank mix or sequential treatment within 30 days of planting in a lentil weed control program.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply **Sharpen** if cold and/or wet conditions are present or predicted to occur within 1 week of application.
- Plant legumes at least 1/2-inch deep to reduce risk of crop injury from **Sharpen** application.
- **DO NOT** apply when legumes have reached the cracking stage or after emergence or severe crop injury will occur.
- **DO NOT** apply **Sharpen** with other products containing **Group 14/Group E** herbicides (such as sulfentrazone or flumioxazin) as a tank mix partner or sequential application within 30 days of planting because crop injury may result.



- There is no required (preharvest) interval between a preplant or preemergence application of **Sharpen® herbicide** and the harvest of mature legume pods or seeds.
- Legume forage may be fed or grazed 65 or more days after application.
- Refer to **Table 4** for rotational crop planting intervals.

## Tank Mixtures

Broad-spectrum burndown control of grasses and/or additional broadleaf weeds will usually require a tank mix with another herbicide. **Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Prowl® H<sub>2</sub>O herbicide**
- **Pursuit® herbicide**
- glyphosate (e.g. **Roundup® herbicide**)

## Rice

**Sharpen** may be applied preplant, preemergence, or postemergence in rice for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled; refer to **Table 7** for additional rice-specific weeds controlled).

Rice in this label refers to drilled or dry-seeded, and water-seeded rice. Before applying **Sharpen** to rice, verify the selectivity of **Sharpen** on your variety or hybrid with your local seed company (supplier) and/or BASF representative to help avoid potential injury to sensitive varieties or hybrids.

## Application Rates and Timings Preplant and Preemergence Burndown plus Residual Weed Control

Apply **Sharpen** as a broadcast burndown spray preplant through preemergence (0 to 3 days after rice planting) at 1.0 to 2.0 fl ozs/A. An adjuvant system including MSO and AMS (refer to **Additives** section for details) is required for optimum broadleaf burndown activity. Best product performance is obtained when broadleaf weeds are small and actively

growing. Thorough spray coverage of emerged weeds is required; higher spray volumes may be needed for best performance.

For additional residual broadleaf weed control, **Sharpen** can be applied preplant through preemergence at rates of 3.0 to 4.0 fl ozs/A.

## Postemergence Weed Control

Apply **Sharpen** for postemergence control of small and actively growing emerged broadleaf weeds at 1.0 fl oz/A. Postemergence applications can be made before or after flooding when rice has reached the 2-leaf stage up to internode elongation. **DO NOT** apply **Sharpen** to rice in the spike or 1-leaf stage.

**Table 7. Additional Weeds Controlled in Rice by Postemergence Applications of Sharpen**

Common Name	Scientific Name	Maximum Height (inches)
Dayflower	<i>Commelina</i> spp.	3
Ducksalad	<i>Heteranthera limesa</i>	3
Eclipta	<i>Eclipta alba</i>	4
Flat sedge	<i>Cyperus iria</i>	3
Jointvetch, Indian	<i>Aeschynomene indica</i>	3
Jointvetch, Northern	<i>Aeschynomene virginica</i>	3
Redstem	<i>Ammannia</i> spp.	4
Sesbania	<i>Sesbania exaltata</i>	8
Texasweed	<i>Caperonia palustris</i>	3
Water hyssop	<i>Bacopa eisenii</i>	3
Woolly croton	<i>Croton capitatus</i>	3

**Adjuvant Requirements for Postemergence-specific Applications in Rice.** For optimum postemergence control of emerged broadleaf weeds in rice, the following adjuvants are required with **Sharpen**:

- Use a crop oil concentrate (COC) at a rate of 1 pint/A to 1 quart/A.

- **DO NOT** use methylated seed oil (MSO) or severe crop injury may occur.
- **DO NOT** use a nonionic surfactant (NIS) as a substitute for COC or poor performance on broadleaf weeds will occur.

**Rice Crop Response.** Temporary leaf burn and/or speckling may occur after postemergence applications; new growth and development is unaffected with rapid recovery under good growing conditions. Severe leaf burn and/or stand loss may occur under stressful growing conditions (e.g. low soil fertility, seedling/foiar disease, extreme hot or cold weather, high soil pH, high soil salt concentration, or drought).

## Sequential Applications

Sequential applications of **Sharpen**® herbicide may be made in rice, but **DO NOT** apply more than a maximum cumulative amount of 6.0 fl ozs/A of **Sharpen** per cropping season.

Separate all sequential applications by at least 14 days.

Preplant or preemergence burndown applications may be applied as part of a sequential application program where the first application is made preplant or preemergence, and the sequential application(s) is made postemergence. Postemergence applications may be sequentially made before or after flooding.

For postemergence applications, **DO NOT** apply more than a maximum cumulative amount of 2.0 fl ozs/A of **Sharpen** per cropping season.

**State-specific Use in California.** Apply **Sharpen** as a broadcast burndown spray at 1.0 to 2.0 fl ozs/A at least 15 days before rice planting and 45 days before a permanent flood is established. An adjuvant system (refer to **Additives** section for details) is required for optimum broadleaf burndown activity. **DO NOT** apply **Sharpen** after paddy flooding or postemergence to the crop. **DO NOT** apply more than 2.0 fl ozs/A in a single application. **DO NOT** apply within 45 days of permanent flooding in water-seeded rice paddies. **DO NOT** use released tailwater for irrigation of adjacent crops.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply more than a maximum cumulative amount of 6.0 fl ozs/A of **Sharpen** per cropping season.
- **Sharpen** may be applied to rice fields used for crustacean (including crayfish) production and commercial fish production.
- **DO NOT** apply **Sharpen** to rice fields that will also be used for mollusk production during the treatment year.
- **DO NOT** release flood water from treated fields for 7 days after **Sharpen** application.
- Refer to **Table 4** for rotational crop planting intervals.

## Tank Mixtures

Broad-spectrum burndown control of grasses and/or additional broadleaf weeds will require a tank mix with another herbicide. Read and follow the applicable restrictions and limitations and directions for use on the other product label. The most restrictive labeling applies to tank mixes. **Sharpen** may be tank mixed or applied sequentially with, but not limited to:

- **Beyond**® herbicide
- **Clearpath**® herbicide
- **Facet**® L herbicide
- **Newpath**® herbicide
- **Prowl**® H<sub>2</sub>O herbicide
- glyphosate (e.g. **Roundup**® herbicide)

Tank mixtures with products formulated as emulsifiable concentrates (EC) may enhance the crop injury potential of **Sharpen** in postemergence applications.

## Small Grains (barley, oats, pearl millet, proso millet, rye, triticale, and wheat)

**Sharpen® herbicide** may be applied preplant surface, preplant incorporated, or preemergence to small grains for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled). Small grains in this label refers to wheat (including durum, spring and winter), barley, oats, millet (pearl and proso), rye, and triticale. Before applying **Sharpen** to small grains, verify the selectivity of **Sharpen** on your variety with your seed company (supplier) to help avoid potential injury to sensitive varieties.

### Application Rates and Timings

**Burndown.** Apply **Sharpen** for burndown and/or residual control of broadleaf weeds early preplant through preemergence at 1.0 to 2.0 fl ozs/A. **Sharpen** at 2.0 fl ozs/A will provide limited residual control of broadleaf weeds. Performance depends on amount of rainfall for activation, soil texture, and broadleaf species/population. An adjuvant system (refer to the **Additives** section for details) is required for optimum broadleaf burndown activity.

**Sequential Applications.** **Sharpen** may be applied sequentially as needed before small grain emergence. Early preplant applications may be applied as part of a split application program where the first application is made early preplant and the second application is made preemergence. Separate sequential applications by at least 30 days in millet and by at least 14 days in all other small grains.

### Crop-specific Restrictions and Limitations

- **DO NOT** apply more than a maximum cumulative amount of 4.0 fl ozs/A of **Sharpen** per cropping season.
- **DO NOT** apply after small grain emergence or crop injury will occur.
- Small grain forage and hay can be fed or grazed 30 or more days after application.

- Ensure that the seed row is sufficiently covered with soil to avoid washing and concentration of the herbicide in the seed zone.
- **DO NOT** apply to other types of millet (e.g. foxtail millet) or severe crop injury may occur.
- **DO NOT** apply to millet grown in soils with a pH of 7.8 or above or crop injury may occur.
- Refer to **Table 4** for rotational crop planting intervals.

### Tank Mixtures

Broad-spectrum burndown control of grasses and/or additional broadleaf weeds will usually require a tank mix with another herbicide. **Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Clarity® herbicide**
- glyphosate (e.g. **Roundup® herbicide**)

## Sorghum (grain)

**Sharpen** may be applied preplant or preemergence to sorghum (grain sorghum) for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled). Before applying **Sharpen** to sorghum, verify with your local seed company (supplier) the selectivity of **Sharpen** on your hybrid or variety to help avoid potential injury to sensitive hybrids or varieties.

### Burndown Weed Control

**Sharpen** can be applied at 1.0 to 2.0 fl ozs/A (all soil types) with an adjuvant system (refer to the **Additives** section for details) any time before sorghum emergence to provide burndown of weeds listed in **Table 1**. A burndown application of **Sharpen** can be followed by residual rates of **Verdict® herbicide**. Sequential applications must be separated by at least 14 days. However, **DO NOT** exceed the cropping seasonal maximum cumulative amount per acre of saflufenacil from all product sources.

**State-specific Use in California.** Apply **Sharpen** early preplant through preemergence at 1.0 to 2.0 fl ozs/A with an adjuvant system for enhanced burndown broadleaf weed control before crop emer-

gence. Separate sequential applications by at least 14 days. **DO NOT** apply greater than 2.0 fl ozs/A in a single application.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply **Sharpen® herbicide** after sorghum emergence or severe crop injury will occur.
- **DO NOT** apply **Sharpen** at greater than 1.0 fl oz/A within 30 days of planting where an at-planting application of an organophosphate or carbamate insecticide(s) is planned and/or has occurred or severe injury may result.

**EXCEPTION:** **Sharpen** may be applied when **Aztec® 2.1% granular insecticide**, **Aztec® 4.67 G granular insecticide**, **Fortress® 5G granular insecticide**, or **SmartChoice™ 5G granular insecticide** is applied at planting as a band, T-band, or in-furrow. **Sharpen** may be applied with all other classes of at-planting insecticides including pyrethroids, neonicotinoids, and fipronil.

- **DO NOT** apply more than a maximum cumulative amount of 0.111 lb ai/A of saflufenacil per cropping season in sorghum from all product sources.
- Sorghum forage can be harvested, fed, or grazed 70 days or more after application.
- Refer to **Table 4** for rotational crop planting intervals.

## Tank Mixtures

**Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Clarity® herbicide**
- **Outlook® herbicide**
- **Verdict® herbicide**
- atrazine
- glyphosate (e.g. **Roundup® herbicide**)

**NOTE:** Sorghum seed must be properly treated with an approved chloroacetamide safener when applying a chloroacetamide herbicide or sorghum injury will occur.

## Soybean

**Sharpen** may be applied in the fall and/or in the spring as a preplant surface or preemergence burndown application in conventional-till, reduced-till, or no-till soybeans for broadleaf weed control (refer to **Table 1** and **Table 2** for lists of weeds controlled). An adjuvant system (refer to **Additives** section for details) is required for optimum burndown activity.

Use of **Sharpen** may result in delayed soybean emergence and stunting under certain environmental conditions including cool temperatures, excessive rainfall/irrigation, and/or persistent wet soil conditions occurring after application.

Not for use in soybean in California.

## Application Rates and Timings

### Fall Application

Apply **Sharpen** at 1.0 to 2.0 fl ozs/A for burndown broadleaf weed control after the prior crop is harvested. Application must be made before first killing frost. Fall application can be made to all soil types.

### Spring Application

Apply **Sharpen** early preplant through preemergence at 1.0 fl oz/A for burndown broadleaf weed control before crop emergence.

Apply **Sharpen** early preplant at 1.5 or 2.0 fl ozs/A for enhanced burndown broadleaf weed control.

### Soybean Planting Interval

Depending on **Sharpen** use rate, soil texture and organic matter, an interval between **Sharpen** application and planting may be required (see **Table 8** and **Table 9**). This interval must be observed before planting soybean or crop injury may occur.

**Table 8. Minimum Soybean Planting Intervals**

Minimum Preplant Interval (days) Required between Sharpen® herbicide Application and Soybean Planting		
Sharpen Use Rate (fl ozs/A)	Soil Texture <sup>1</sup>	
	Coarse Soils with ≤ 2.0% Organic Matter	All Other Soils
1.0	30	0
1.5	30	14
2.0	44	30

<sup>1</sup> Refer to **Table 3** for definition of soil texture groups.

**Table 9. Minimum Soybean Planting Intervals with other Group 14/Group E Herbicides**

Minimum Preplant Interval (days) Required between Sharpen Application and Soybean Planting when Tank Mixed or Sequentially Applied with a Group 14/Group E Herbicide <sup>1</sup>		
Sharpen Use Rate (fl ozs/A)	Soil Texture <sup>2</sup>	
	Coarse Soils with ≤ 2.0% Organic Matter	All Other Soils
1.0	30	14*
1.5	30	30
2.0	44	30

<sup>1</sup> **Group 14/Group E** herbicides such as sulfentrazone or flumioxazin

<sup>2</sup> Refer to **Table 3** for definition of soil texture groups.

\* Interval for reduced-till and no-till soybean only. Interval for conventional-till soybean is 30 days.

## Crop-specific Restrictions and Limitations

- **DO NOT** apply more than a maximum cumulative amount of 4.0 fl ozs/A of **Sharpen** (0.089 lb ai/A of saflufenacil) per cropping season. Sequential applications **MUST** be separated by at least 30 days.
- **DO NOT** apply more than a maximum cumulative amount of 0.089 lb ai/A of saflufenacil per cropping season in soybean from all product sources.
- **DO NOT** apply when soybean has reached the cracking stage or after emergence or severe crop injury will occur.
- **DO NOT** apply **Sharpen** within 30 days of planting where an at-planting application of an organophosphate or carbamate insecticide(s) is planned and/or has occurred because severe injury may result.
- Ensure that the seed row is sufficiently covered with soil to avoid washing and concentration of the herbicide in the seed zone.
- Always use the most restrictive preplant interval of all inclusive herbicides when applying **Sharpen** as part of a tank mix.
- Soybean forage may be fed or grazed 65 or more days after application.
- Refer to **Table 4** for rotational crop planting intervals.
- **Group 14/Group E** herbicides labeled for postemergence application in soybean may be used 14 days or more after soybean emergence.

## Tank Mixtures

Broad-spectrum burn-down control of grasses and/or additional broadleaf weeds will usually require a tank mix with a herbicide such as glyphosate. **Sharpen** may be tank mixed with or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Clarity®** herbicide
- **Extreme®** herbicide
- **Optill® PRO** herbicide
- **Prowl® H<sub>2</sub>O** herbicide
- **Pursuit®** herbicide
- **Scepter®** herbicide
- **Verdict®** herbicide
- **Zidua®** herbicide
- glyphosate (e.g. **Roundup®** herbicide)

## Noncropland Areas

**DO NOT** apply **Sharpen®** herbicide in any residential setting.

**Sharpen** may be used:

- In noncropland areas including fence rows and nonirrigation ditchbanks

### Application Rates and Timings

**Sharpen** may be applied either in a single application or sequentially with an interval of 14 days or more.

Application rates for **Sharpen** when applied alone, in tank mix, or sequentially are given in **Table 10**.

**DO NOT** apply more than a maximum cumulative total of 6.0 fl ozs of **Sharpen** (0.134 pound active ingredient saflufenacil) per acre per year.

In California, **DO NOT** apply more than 2.0 fl ozs/A of **Sharpen** in a single application.

**Table 10. Application Rates in Noncropland Areas**

Application	Application Target	Application Rate (fl ozs/A)
Postemergence	Weed size < 6 inches	2 to 4
	Weed size ≥ 6 inches and/or heavier weed infestations	4 to 6 <sup>a</sup>
Postemergence + Residual	Burndown + Residual preemergence weed control	6 <sup>b</sup>

(continued)

**Table 10. Application Rates in Noncropland Areas** (continued)

Application	Application Target	Application Rate (fl ozs/A)
<b>Tank Mixes with Glyphosate</b>		
<b>Accelerated Burndown</b>	Accelerated burndown of broadleaf weeds and/or control of glyphosate-tolerant species [such as horseweed (marestail)]	1 to 2
<b>Accelerated Burndown + Residual</b>	Accelerated burndown of broadleaf weeds plus control of glyphosate-tolerant species with residual preemergence weed control	6 <sup>b</sup>

<sup>a</sup> Partial control or suppression may result with applications to weeds > 6 inches.

<sup>b</sup> To provide effective residual control of labeled weed species, **Sharpen MUST** be used at the maximum use rate of 6 fl ozs/A.

### Spot Treatments

**Sharpen** may be applied as a spot treatment to emerged broadleaf weeds. Consult the chart following for the amount of **Sharpen** to make various gallons of spray mix to use for spot treatments applied to actively growing broadleaf weeds and sizes referenced in **Table 1**. Spray thoroughly to wet weed foliage but not to the point of runoff.

To maximize performance, refer to the **Additives** section for recommended adjuvant and rate to add to the spray mix.

Each spray mix is equivalent to applying **Sharpen® herbicide** at a use rate of 2.0 fl ozs/A in a spray volume of 100 gallons per acre. Applications of a spot spray mix should not be made to an equivalent area less than what is shown in the chart or exceed the equivalent broadcast rate of 2.0 fl ozs/A.

Spot treatments may be applied via an ATV-mounted (all-terrain vehicle-mounted) or tractor-mounted sprayer equipped for low-pressure hand wand applications. **DO NOT** apply spot treatments using high-pressure hand wands.

Spray Mix (gallons)	Spray Mix Treatment Area (sq ft)	Sharpen (fl oz)
10	4,356	0.2
25	10,890	0.5
50	21,780	1.0

## Selective Weeding

Apply **Sharpen** up to 2.0 fl ozs/A as a postemergence spray plus the recommended adjuvant (refer to **Additives** section for details) as a uniform broadcast application or spot treatment for selective broadleaf weed control.

## Tank Mixtures for Selective Weeding

Broad-spectrum postemergence and/or residual control of grasses or additional broadleaf species will require a tank mix with another herbicide. **Sharpen** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Plateau® herbicide**
- glyphosate (e.g. **Roundup® herbicide**)

## Leafy Spurge Control

**Sharpen** applied in tank mixture with **Plateau** will control leafy spurge when applied late spring/early summer in noncropland areas as described above. This tank mixture will also control additional weeds listed on the respective **Sharpen** and **Plateau** labels. **Sharpen** plus **Plateau** tank mix may be applied by either ground or air as either a uniform broadcast application or a spot treatment.

Apply **Sharpen** at 1.0 to 2.0 fl ozs/A plus **Plateau** at 4.0 to 6.0 fl ozs/A to leafy spurge when it reaches the yellow bract (pre-bloom) stage in late spring/early summer. **DO NOT** apply this tank mix as a fall application because control may not be satisfactory.

**DO NOT** feed or allow animals to graze areas of grass treated with **Sharpen** within 365 days of treatment.

**Spray Additives for Leafy Spurge Control.** **Sharpen** plus **Plateau** tank mix requires the use of an effective adjuvant system. For best results, use a nonionic surfactant at 0.25% volume/volume (v/v) plus ammonium sulfate at 8.5 to 17.0 lbs/100 gals [1% to 2% weight/volume (w/v)]. Crop oil concentrate or methylated seed oil may also be used with this tank mixture when injury (stunting, necrosis) to grasses is acceptable.

**Water Volume for Leafy Spurge Control.** For ground applications, use 10 or more gallons of water per acre. Thorough coverage of weeds is essential and higher spray volumes may be necessary for better performance on a heavy population of leafy spurge. For aerial applications, use 5 or more gallons of water per acre.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

*(continued)*

## Conditions of Sale and Warranty

*(continued)*

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108



**Beyond, Clarity, Clearpath, Distinct, Extreme, Facet, Kixor, Newpath, Optill, Outlook, Plateau, Prowl, Pursuit, Scepter, Sharpen, Status, Verdict, and Zidua** are registered trademarks of BASF.

**Aztec, Def, Dropp, Finish, and Ginstar** are registered trademarks of Bayer.

**Prep** is a trademark of Bayer.

**CottonQuik** is a registered trademark of Nufarm Americas Inc.

**Folex and Fortress** are registered trademarks of AMVAC Chemical Corporation.

**Gramoxone** is a registered trademark of a Syngenta Group Company.

**Harness and Roundup** are registered trademarks of Monsanto Technology LLC.

**SmartChoice** is a trademark of AMVAC Chemical Corporation.

© 2014 BASF Corporation  
All rights reserved.

007969-00278.20141107.**NVA 2014-04-322-0283**

Based on: NVA 2014-04-322-0154

Supersedes: NVA 2014-04-322-0192

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry

## NOTES

## NOTES



## NOTES

## NOTES

## NOTES





# Sharpen®

Group 14 Herbicide

Powered by Kixor® Herbicide

For use in selected agricultural crops

**Active Ingredient:**

sulfafenacil: N-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3, 6-dihydro-1[2H]-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide

29.74%

**Other Ingredients:**

70.26%

**Total:**

100.00%

Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-278

EPA Est. No. 241-PR-002

## KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty,** and state-specific crop and/or use site restrictions. **In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

**First Aid – If swallowed:** Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything to an unconscious person.

**If in eyes:** Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. Call a poison control center for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If**

**inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice. **HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information at 1-800-832-HELP (4357).

**Precautionary Statements**

**Hazards to Humans and Domestic Animals. CAUTION.** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

**Environmental Hazards**

For terrestrial uses, **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. See attached booklet for complete Environmental Hazards, Groundwater Advisory, Surface Water Advisory and Endangered Species Protection Requirements.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. **Pesticide Storage: DO NOT** use or store near heat or open flame. Store in original container in a well-ventilated area separately from fertilizer, feed, or foodstuffs. Avoid cross-contamination with other pesticides. **Pesticide Disposal:** Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Handling: Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. See attached booklet for complete container handling directions including triple rinsing and pressure rinsing instructions.

**Net Contents: 1 gallon**



We create chemistry

67611782 NVA 2015-05-322-0042

Product of U.S.A.

BASF Corporation, 26 Davis Drive, Research Triangle Park, NC 27709