

ACCEPTED  
FOR REGISTRATION

May 8, 2018

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

Doc id: 556999

GROUP 3 | 11 FUNGICIDES



## FUNGICIDE

*Broad spectrum fungicide for control of plant diseases*

**ACTIVE INGREDIENT:**

Azoxystrobin ..... 7.0%  
Propiconazole ..... 11.7%

**OTHER INGREDIENTS:** ..... 81.3%

**TOTAL:** ..... 100.0%

**AVARIS™** Fungicide is formulated as a suspoemulsion (SE) formulation.

**AVARIS™** Fungicide contains 1.04 lb ai propiconazole and 0.62 lb ai azoxystrobin per gallon.

### KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET.  
SEE FIRST AID STATEMENT INSIDE BOOKLET OR ON CONTAINER LABEL.**

#### FIRST AID

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

#### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency, call Chemtrec at **1-800-424-9300**

SCPSR-HEL 1178B-L1F 1016  
4089111

NET CONTENTS:  
2.5 Gallons (9.46 L)

EPA Reg. No. 100-1178-5905  
EPA Est. 100-NE-001

Manufactured For

**HELENA CHEMICAL COMPANY**

225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017

9013-13166 11-22-17 SCP 01-00 (11-29-17)  
Avaris (Helena Chemical) – 2.5 gal. booklet/base – Mech  
SCP 2590... 3213... 3819... 5338... 5776... 6861... 8742...  
9013...

**AGI 107035**  
Print size – Booklet: 5.75" wide x 6.75" high  
Base: 6.75" wide x 6.75" high  
Pantone colors – PMS 021, PMS 2612, Black  
No. of pages – 16

## NOTES

---

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

## WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes and socks

#### User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Propiconazole is toxic to shrimp.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

#### Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Helena Chemical Company immediately if you observe any adverse environmental effects due to use of this product.

---

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

---

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of HELENA CHEMICAL COMPANY or Seller. To the extent permitted by applicable law, Buyer and User agree to hold HELENA and Seller harmless for any claims relating to such factors.

HELENA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or HELENA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, HELENA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall HELENA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF HELENA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF HELENA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

HELENA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of HELENA.

---

## DIRECTIONS FOR USE

---

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

---

## AGRICULTURAL USES

---

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks
- Protective eyewear

---

## PRODUCT INFORMATION

---

**AVARIS™** Fungicide is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. **AVARIS™** may also improve the yield and/or quality of the crop. These additional benefits are due to positive effects on plant physiology. The effects may vary according to other factors such as crop, crop hybrid, or environment. **AVARIS™** may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications should be made according to the use directions that follow.

---

## PRODUCT USE PRECAUTIONS

---

Do not use in nurseries, greenhouses or landscape plantings.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult [www.epa.gov/esp/ bulletins.htm](http://www.epa.gov/esp/ bulletins.htm). Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

---

## PRODUCT USE INSTRUCTIONS

---

**Application:** Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

**Adjuvants:** When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

**Efficacy:** Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of **AVARIS™** has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

**Rotational Crops:** Soybeans may be planted as a double crop following a cereal crop which has been treated with **AVARIS™**. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding. Plant wild rice 45 days after **AVARIS™** application. To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of **AVARIS™** application to the preceding crop, unless the second crop appears on this label.

---

Rotational Crops	Planting Time From Last AVARIS™ Application
Bean, dry and succulent Bulb crops Berries, bushberry and caneberry subgroups Brassica, leafy greens (subgroup 5B) Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale, oat and rye) Corn (field, seed, popcorn, and sweet) Dill Grasses grown for seed (Cool season grasses only) Mint Peanuts Quinoa Radish Rice Sorghum Soybeans Strawberries Sugar beets Watercress	0 days

Rotational Crops ( <i>continued</i> )	Planting Time From Last AVARIS™ Application
Buckwheat Millet	12 Months
Wild rice	45 days
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/season)	75 days
All Other Crops Intended for Food and Feed	105 days

**Crop Tolerance/Phytotoxicity:** AVARIS™ Fungicide demonstrates some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone can contribute to phytotoxicity. Under certain environmental conditions, tank mixes of AVARIS™ plus herbicides and/or fertilizers may cause crop injury in barley, triticale and wheat.

**—ATTENTION—**

AVARIS™ is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray AVARIS™ where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply AVARIS™ to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

**Spray Drift Management:** To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

**Integrated Pest Management:** AVARIS™ should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The SPECIFIC USE DIRECTIONS section in this label identifies specific IPM recommendations for each crop. Consult your local agricultural authorities for additional IPM strategies established for your area. AVARIS™ may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

---

## RESISTANCE MANAGEMENT

---

GROUP	<b>3</b>	<b>11</b>	FUNGICIDES
-------	----------	-----------	------------

AVARIS™ Fungicide is a mixture of Group 3 (propiconazole) and Group 11 (azoxystrobin) fungicides. AVARIS™ has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Helena Chemical Company encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the specific crop recommendations that limit the total number of sprays on a crop and the required alternations with fungicides from other resistance management groups. In situations requiring multiple sprays, develop season long spray

programs for Group 11 (QoI) fungicides. The program should meet the goal of no more than 1/3 of the total sprays per season, when a Group 11 fungicide is used as a solo product, or 1/2 the total sprays when a Group 11 fungicide is used in a mixture. Programs that include both solo Group 11 products and/or mixes containing Group 11 products should be no more than 1/2 the total sprays.

**AVARIS™** should not be alternated or tank mixed with any fungicide to which resistance has already developed.

---

## SPRAY EQUIPMENT

---

**AVARIS™** may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

### NOZZLES

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be *16-mesh or coarser*.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

### PUMP

- Use a pump with capacity to:
  - (1) Maintain 35-40 psi at nozzles.
  - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

---

## MIXING INSTRUCTIONS

---

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

### **AVARIS™ Alone (no tank mix):**

- Add 1/2-2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add **AVARIS™** to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after **AVARIS™** has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

**AVARIS™ + Tank Mixtures:** **AVARIS™** is usually compatible with all tank mix partners listed on this label. Do not combine **AVARIS™** in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and noninjurious to the crop under your conditions of use. To determine the physical compatibility of **AVARIS™** with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water dispersible granular products first, then liquid flowables (suspoemulsions), and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

### Mixing in the Spray Tank

- Add 1/2-2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank mix partner(s) into the tank in the same order as described above in the “**AVARIS™** + Tank Mixtures” section.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the **AVARIS™** to the spray tank.
- Allow **AVARIS™** to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.
- No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product may not be mixed with any product which prohibits such mixing.

---

## APPLICATION PROCEDURES

---

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner that will result in exposure to humans or animals.

### Ground Application:

- For field crops (non-trees), apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise.

### Aerial Application:

- Use only on crops where aerial applications are indicated.
- For field crops (non-trees), apply in a minimum spray volume of 2 gallons per acre unless specified otherwise.
- For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- Thorough coverage is necessary to provide good disease control.
- **AVARIS™** is extremely phytotoxic to certain apple varieties.
- AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).
- DO NOT spray **AVARIS™** where spray drift may reach apple trees.
- DO NOT apply via ULV.

### Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches per acre of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Note:** Do not inject **AVARIS™** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part **AVARIS™**. **AVARIS™** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton®, Buna-N, Neoprene, or PVC seals.

---

## APPLICATION INSTRUCTIONS

---

Apply **AVARIS™** Fungicide at rates and timings as described in this label.

**Directions for Use Through Sprinkler and Drip Chemigation Systems (Apply AVARIS™ only to crops for which chemigation is specified on this label):**

**Spray Preparation:** Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.



## **Use Precautions for Sprinkler and Drip Irrigation Applications**

**Sprinkler Irrigation:** Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, drip (trickle), or hand move irrigation systems. Do not apply this product through any other type of irrigation system except as specified on this label.

Apply with center pivot or continuous-move equipment distributing 1/2 acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set. Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips. Do not apply when wind speed favors drift beyond the area intended for treatment. Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

### **Specific Instructions for Public Water Systems**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Cereals</b> Wheat  <b>See next section for other cereals.</b>	<b>Early season suppression of:</b> Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blight <i>(Septoria tritici)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	7 - 14	Apply <b>AVARIS™</b> in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.  If applying early in the season in mixture with fertilizers and herbicides you may see flecking and burning.
	<b>Control of Leaf Diseases:</b> Glume Blotch <i>(Stagonospora nodorum)</i> Helminthosporium Leaf Blight <i>(Drechslera tritici-repentis)</i> Kernel Blight <i>(Alternaria spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Net Blotch <i>(Pyrenophora teres)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Rust <i>(Puccinia spp.)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when <b>AVARIS™</b> is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl oz/A may be applied.  Applications may be made no closer than a 14-day interval.  <b>AVARIS™</b> can be applied through full head emergence (Feekes growth stage 10.54). Do not apply after this stage.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of <b>AVARIS™</b> plus half the rate listed of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.
<p><b>Application:</b> <b>AVARIS™</b> is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. <b>AVARIS™</b> may be applied by ground, air, or chemigation.</p>			
<p><b>Specific Use Restrictions:</b></p> <ol style="list-style-type: none"> <li>1) Do not apply more than 2 applications/A/season.</li> <li>2) <b>AVARIS™</b> can be applied through full head emergence (Feekes growth stage 10.54). Do not apply after this stage.</li> <li>3) Do not apply more than 28.0 fl oz/A/season of <b>AVARIS™</b>.</li> <li>4) Do not apply more than 0.22 lb ai propiconazole-containing products/A/season.</li> <li>5) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/season.</li> <li>6) Under certain environmental conditions, tank mixes of <b>AVARIS™</b> plus herbicides and/or fertilizers may cause crop injury.</li> <li>7) Do not harvest within 7 days (7-day PHI) for forage or hay.</li> </ol>			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Cereals</b> Barley Oats Rye Triticale	<b>Early season suppression of:</b> Glume Blotch <i>(Stagonospora nodorum)</i> Leaf Blight <i>(Septoria tritici)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	7 – 14	Apply <b>AVARIS™</b> in the spring for suppression of early season diseases. Follow up with a second application (see below) for full season control.  If applying early in the season in mixture with fertilizers and herbicides you may see flecking and burning.
	<b>Control of Leaf Diseases:</b> Barley Scald <i>(Rhynchosporium secalis)</i> Barley Stripe <i>(Pyrenophora graminea)</i> Glume Blotch <i>(Stagonospora nodorum)</i> Helminthosporium Leaf Blight <i>(Drechslera tritici-repentis)</i> Kernel Blight <i>(Alternaria spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Net Blotch <i>(Pyrenophora teres)</i> Powdery Mildew <i>(Blumeria spp., Erysiphe spp.)</i> Rust <i>(Puccinia spp.)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when <b>AVARIS™</b> is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl oz/A may be applied.  Apply no closer than a 14-day interval.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of <b>AVARIS™</b> plus half the rate specified on other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.
<p><b>Application:</b> <b>AVARIS™</b> is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions would not provide good coverage. An adjuvant may be added at recommended rates to improve canopy coverage and penetration while reducing evaporation and drift. <b>AVARIS™</b> may be applied by ground, air, or chemigation.</p>			
<p><b>Specific Use Restrictions:</b></p> <ol style="list-style-type: none"> <li>1) Do not apply more than 2 applications/A/season.</li> <li>2) <b>AVARIS™</b> can be applied through full head emergence (Feekes growth stage 10.54). Do not apply after this stage.</li> <li>3) Do not apply more than 28.0 fl oz/A/season of <b>AVARIS™</b>.</li> <li>4) Do not apply more than 0.22 lb ai propiconazole-containing products/A/season.</li> <li>5) Do not apply more than 0.40 lb ai azoxystrobin-containing products/A/season.</li> <li>6) Under certain environmental conditions, tank mixes of <b>AVARIS™</b> plus herbicides and/or fertilizers may cause crop injury.</li> <li>7) Do not apply within 7 days of harvest (7-day PHI) for forage and hay.</li> </ol>			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Corn</b> Field Pop (Includes Seed Production)	Northern Corn Leaf Blight <i>(Setosphaeria turcica)</i> Northern Corn Leaf Spot <i>(Cochliobolus carbonum)</i> Southern Corn Leaf Blight <i>(Cochliobolus heterostrophus)</i>  also known as Helminthosporium Leaf Blights <i>(H. maydis, H. turcicum,</i> <i>H. carbonum)</i>	7 - 14	For leaf blights apply <b>AVARIS™</b> when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.  Apply no more than 2 applications of <b>AVARIS™</b> or any other Group 11 fungicide per year.  Do not use adjuvants or other additives after the V8 growth stage and prior to the VT growth stage, as use during these development times may impose stress on the plant that could inhibit proper kernel development. VT is defined as when the last branch of the tassel is completely visible but silks have not yet emerged from the ear shoot.
	Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Eye Spot <i>(Aureobasidium zeae)</i> Gray Leaf Spot <i>(Cercospora zeae-maydis)</i> Rusts <i>(Puccinia spp.)</i>	10.5 - 14	Apply <b>AVARIS™</b> when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Under heavy disease pressure or if conditions are favorable for disease, apply the highest labeled rate.  Apply no more than 2 applications of <b>AVARIS™</b> or any other Group 11 fungicide per year.
<p><b>Application:</b> <b>AVARIS™</b> may be applied by ground, air, or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.</p>			
<p><b>Specific Use Restrictions:</b></p> <ol style="list-style-type: none"> <li>1) Do not apply more than 56 fl oz/A/season of <b>AVARIS™</b>.</li> <li>2) Do not apply more than 28 fl oz (0.224 lb ai propiconazole) for field corn harvested for forage.</li> <li>3) Do not apply more than 0.45 lb ai propiconazole-containing products/A/season.</li> <li>4) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/season.</li> <li>5) Do not apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.</li> </ol>			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Corn</b> Sweet corn (Includes Seed Production)	Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> )  also known as Helminthosporium Leaf Blights ( <i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> )	7 - 14	For leaf blights apply <b>AVARIS™</b> when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate.  Alternate applications of <b>AVARIS™</b> with Tilt® or another product with a different mode of action than Group 11 fungicides.
	Anthracnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leaf Spot ( <i>Cercospora zeae-maydis</i> ) Rusts ( <i>Puccinia</i> spp.)	10.5 - 14	Apply <b>AVARIS™</b> when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule.  Alternate applications of <b>AVARIS™</b> with Tilt or another product with a different mode of action than Group 11 fungicides.
<p><b>Application: AVARIS™</b> may be applied by ground, air, or chemigation. For best results, sufficient coverage is very important. Use of a crop oil concentrate is recommended for aerial applications to reduce evaporation and enhance canopy penetration and coverage. Consult your aerial applicator for recommended concentration of crop oil concentrate. Use higher water volumes for aerial applications if equipment and/or conditions will not provide good coverage.</p>			
<p><b>Specific Use Restrictions:</b></p> <ol style="list-style-type: none"> <li>1) Do not apply more than 56 fl oz/A/season of <b>AVARIS™</b>.</li> <li>2) Do not apply more than 0.45 lb ai propiconazole-containing products/A/season.</li> <li>3) Do not apply more than 2.0 lb ai azoxystrobin-containing products/A/season.</li> <li>4) Do not apply to sweet corn within 14 days of harvest (14-day PHI) for ears or forage.</li> </ol>			

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Soybeans</b>	Aerial Web Blight ( <i>Rhizoctonia solani</i> ) Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum truncatum</i> ) Brown Spot ( <i>Septoria glycines</i> ) Cercospora Blight and Leaf Spot ( <i>C. kickuchii</i> ) Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Pod and Stem Blight ( <i>Diaporthe</i> spp.) Soybean Rust ( <i>Phakopsora pachyrhizi</i> )	14 - 20.5	<p>Aerial Web Blight: Apply 14 - 20.5 fl oz/A at the first appearance of disease and repeat the application 14 - 21 days later. Under severe disease conditions use the higher rate and shorter interval.</p> <p>Other foliar diseases (except rust): Apply 20.5 fl oz/A at growth stage R3 (early pod set when pods are 1/8-1/4 inch long) and 14 - 21 days later at growth stage R5 (pod fill).</p> <p>Soybean Rust: Apply 14 - 20.5 fl oz/A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter interval when disease is present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p> <p>On certain varieties, <b>AVARIS™</b> applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to <b>AVARIS™</b> treatments.</p>
<p><b>Application: AVARIS™</b> is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Use a higher water volume for aerial application (greater than 2 GPA) if equipment and/or conditions will not provide good coverage. <b>AVARIS™</b> may be applied by ground, air or chemigation.</p>			
<p><b>Specific Use Restrictions:</b></p> <ol style="list-style-type: none"> <li>1) Do not apply more than 42 fl oz/A/season of <b>AVARIS™</b>.</li> <li>2) Do not apply more than 0.34 lb ai of propiconazole-containing products/A/season.</li> <li>3) Do not apply more than 1.5 lb ai of azoxystrobin-containing products/A/season.</li> <li>4) Apply up to Stage R6.</li> </ol>			

<b>AVARIS™ RATE CONVERSION TABLE</b>			
<b>Fl oz/A</b>	<b>Lb ai/A Propiconazole</b>	<b>Lb ai/A Azoxystrobin</b>	<b>Treated acres per gallon product</b>
3	0.025	0.015	42.6
4	0.033	0.020	32.0
7	0.056	0.034	18.3
10.5	0.085	0.051	12.2
14	0.113	0.070	9.1
15	0.122	0.073	8.5
20.5	0.167	0.100	6.2
27.5	0.224	0.134	4.6
34.5	0.280	0.169	3.7

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

### **PESTICIDE STORAGE**

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

### **PESTICIDE DISPOSAL**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance in proper disposal methods.

### **CONTAINER HANDLING [equal to or less than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

### **CONTAINER HANDLING [greater than 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

*continued...*

## **STORAGE AND DISPOSAL** *(continued)*

### **CONTAINER HANDLING [greater than 5 gallons]**

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

**AVARIS™** trademark of Helena Holding Company

Tilt® is a trademark of a Syngenta Group Company

Topsin® is a trademark of Pennwalt Corporation

Viton® is a trademark of E.I. DuPont de Nemours and Company

### **Manufactured for:**

Helena Chemical Company

225 Schilling Boulevard, Suite 300

Collierville, TN 38017

**SCPSR-HEL 1178B-L1F 1016**

**4089111**



GROUP 3 11 FUNGICIDES



## FUNGICIDE

*Broad spectrum fungicide for control of plant diseases*

### ACTIVE INGREDIENT:

Azoxystrobin	7.0%
Propiconazole	11.7%
OTHER INGREDIENTS:	81.3%

**TOTAL:** 100.0%

**AVARIS™** Fungicide is formulated as a suspoemulsion (SE) formulation.

**AVARIS™** Fungicide contains 1.04 lb ai propiconazole and 0.62 lb ai azoxystrobin per gallon.

## KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET.

SEE FIRST AID STATEMENT INSIDE BOOKLET OR ON CONTAINER LABEL.

### FIRST AID

<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency, call Chemtrec at 1-800-424-9300.

EPA Reg. No. 100-1178-5905 EPA Est. No. 100-NE-001

AVARIS™ trademark of Helena Chemical Company

NET CONTENTS: 2.5 Gallons (9.46 L)

## PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals WARNING/AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### ENVIRONMENTAL HAZARDS

Azoxystrobin and propiconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Propiconazole is toxic to shrimp.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Helena Chemical Company immediately if you observe any adverse environmental effects due to use of this product.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance in proper disposal methods.

**CONTAINER HANDLING [equal to or less than 5 gallons]:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

SCPSR-HEL 1178B-L1F 1016 4089111

Manufactured For

## HELENA CHEMICAL COMPANY

225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING ▲