

Mouse Anti-PD1/PDCD1/CD279 [PDCD1/922]: MC0909, MC0909RTU7

Intended Use: For Research Use Only

Description: Programmed death-1 (PD1) is a member of the CD28 family of receptors that includes CD28, cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4), inducible costimulator (ICOS), and B- and T-lymphocyte attenuator. These receptors play a role in the cellular immune response. PD1 is a new marker of angioimmunoblastic lymphoma and suggests a unique cell of origin for this neoplasm. Unlike CD10 and bcl-6, PD1 is expressed by few B cells, so it may be a more specific and useful diagnostic marker in angioimmunoblastic lymphoma. It also seems to stain a greater percentage of CD3-positive neoplastic cells in angioimmunoblastic lymphoma than either CD10 or bcl-6. In addition, PD1 expression provides new evidence that angioimmunoblastic lymphoma is a neoplasm derived from germinal center-associated T cells. PD1 expression in angioimmunoblastic lymphoma lends further support to this model of T-cell oncogenesis, in which specific subtypes of T cells may undergo neoplastic transformation and result in specific distinct histologic, immunophenotypic, and clinical subtypes of T-cell neoplasia.

Specifications:


Clone: PDCD1/922
 Source: Mouse
 Isotype : IgG1k
 Reactivity: Human
 Localization: Cytoplasm
 Formulation: Purified antibody in PBS pH7.4, containing < 0.2% BSA and 15mM sodium azide (NaN3).
 Storage: Store at 2°- 8°C.
 Applications: IHC, Flow Cyt., ICC/IF
 Package:

Description	Catalog No.	Size
PD1/PDCD1/CD279 Concentrated	MC0909	1 ml
PD1/PDCD1/CD279 Prediluted	MC0909RTU7	7 ml

IHC Procedure*:

Positive Control Tissue: Angioimmunoblastic lymphoma
 Concentrated Dilution: 50-200
 Pretreatment: 10mM Tris EDTA pH9.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human tonsil tissue stained with anti-PD1 using DAB

References:

1. Pattern of CD14+ follicular dendritic cells and PD1+ T cells independently predicts time to transformation in follicular lymphoma. Smeltzer JP, et al. Clin Cancer Res. Jun 1;20(11):2862-72, 2014.
2. PD-1 expression on peripheral blood T-cell subsets correlates with prognosis in non-small cell lung cancer. Waki K, et al. Cancer Sci. Oct;105(10):1229-35, 2014.
3. Immunomodulating antibodies in the treatment of metastatic melanoma: the experience with anti-CTLA-4, anti-CD137, and anti-PD1. Simeone E, et al. J Immunotoxicol. Jul-Sep;9(3):241-7, 2012.

Doc. 100- MC0909
Rev. A