

## Physico-chemical and microbial assessment of drinking water sources across world's largest glacial deposits in Karakoram Mountain Ranges, Pakistan

Maisoor Ahmed Nafees<sup>1</sup>, Khalil Ahmed<sup>1</sup>, Andrea Lami<sup>3</sup>, Azhar Hussain<sup>2</sup>, Shujaat Ali<sup>2</sup>, Tika Khan<sup>4\*</sup>

<sup>1</sup>Department of Biological Sciences Karakorum International University, Gilgit, Pakistan; <sup>2</sup>Department of Food and Agriculture Karakorum International University, Gilgit, Pakistan; <sup>3</sup>Institute of Ecosystem Studies, Verbania, Italy; <sup>4</sup>Integrated Mountain Areas Research Centre, Karakoram International University, Pakistan.

\*Corresponding author: [tika.khan@kiu.edu.pk](mailto:tika.khan@kiu.edu.pk)

**Abstract:** Karakoram Mountain Ranges, Pakistan harbor largest glacial deposits in the world outside poles. They are key source of life and livelihoods of more than 15 million people living downstream. Water samples tested for physico-chemical and microbial health revealed severe bacterial contamination. A total Bacterial count, occurrence of *E. coli* and *Enterococci* in alpine origin sources indicate their vulnerability and undesirable human interferences into fragile mountain ecosystems making them unsafe for consumption. However, chemical ingredients found within the specified water quality limitation. Water sources need proper attention to mitigate frequency and prevalence of water borne diseases in the area.

[Maisoor Ahmed Nafees, Khalil Ahmed, Andrea Lami, Azhar Hussain, Shujaat Ali, Tika Khan. **Physico-chemical and microbial assessment of drinking water sources across world's largest glacial deposits in Karakoram Mountain Ranges, Pakistan.** *Biomedicine and Nursing* 2025;11(2):19-25]. ISSN 2379-8211 (print); ISSN 2379-8203 (online). <http://www.nbmedicine.org>. 03. doi:[10.7537/marsbnj110225.03](https://doi.org/10.7537/marsbnj110225.03)

**Key words:** *E. coli*; *Enterococci*; water borne diseases; pathogen; bacteria; Himalaya; Hindukush; water quality; Gilgit-Baltistan

### 1. Introduction


In the last few decades global population has bulged so the deterioration in climate changes. This has increase pressure on natural resources in general and water availability in particular (Cropper and Griffiths 1994). More than eighty countries of the world making almost half of the total global population are suffering from acute shortage of water have entered into a crisis situation (Birks 1977). This has led to a dramatic increase in cost of water infrastructure (Rosegrant et al. 2002). Water, both in the rivers and underground has got polluted (Goel 2006). Over one billion people do not access safe drinking water, three billion lack sanitation facility

and eighty percent (80%) of infectious diseases are waterborne killing 2.2 million every year (Petersen 2003; World Bank 1999).

The Central Karakoram National Park (CKNP) within the Karakoram Mountain Ranges, Pakistan holds the major source of fresh water in the country. Out of largest glacial and river systems outside poles Baltoro is the remarkable feeding some 30 tributaries which constitute a total surface area of 1,219 sq. km. These glaciers are life-blood for Pakistan nurturing the mighty Indus and its offshoots. They meet the needs for drinking, domestic use, irrigation, wetlands and industrial use and electricity (Acreman 2001).

Water supply contaminations with sewage and human

diarrhea that occur annually (WHO, 2013). Globally, 1.1 billion



**Figure:** Map of study area showing sampling sites

excreta poses maximum risk to public health (Mara and Cairncross 1989). However, water quality testing enables to avoid such pollutions (Hrudey et al. 2003). The sporadic eruption of water borne diseases has coerced the experts to assess the relevant sources for the responsible pathogens and water quality indicators. Contaminated drinking water contributed to a number of health issues in developing countries like more than one billion cases of

people rely on unsafe drinking water sources from lakes, rivers, and open wells (WHO, 2000).

Mountain communities living close to these perennial mighty glacial masses depend for their subsistence agriculture, drinking, household use and power generation. It is important to assess water quality to know ranges of human interventions and their impact on the lives of rural communities. Present study was carried

out for the first time in history from the area. It was aimed at providing scientific insights of water quality, its fitness for consumption, level of contamination, nature of contamination, polluting agents and potential mitigation measures. Furthermore, findings will help water analysis and monitoring schemes to improve overall water quality management and protection of important water bodies and ecosystems in CKNP and beyond.

## 2. MATERIALS AND METHODS

### Sampling

A total of 60 samples were collected from Basha and Baraldu valleys in Central Karakoram National Park (**See figure**). Collection sites were divided into three categories i.e. source, mid and last. HDPE pre sterilized bottles of 500 ml and 1000 ml were used for sample collection. Ten (n=10) samples were collected from each sites with slight variation in distances including source, middle points and last consumption points. Sampling was done in closer conformity with general guide to the sampling techniques and recommendations for preservation and handling of samples (**ISO 5667-1, 5667-2**). Samples were transported in battery operated refrigerators to the 'Water Quality Lab', Karakoram International University (KIU) and processed immediately. All ecological attributes were taken into account with GPS coordinates for future monitoring and to develop a permanent system of surveillance. Water sampling and subsequent analysis campaign continued between June 2012 and November 2014.

### Bacteriological Analysis

Following the standard procedure for examination of water and waste water, samples were analyzed for various microbiological (Bacteriological) parameters (**APHA and AWWA 2000**). The parameters analyzed were *E. coli*, *Enterococci* (intestinal), Total Bacterial Count and *Salmonella*. The membrane filtration technique (ISO 7704) was used to filter 100 ml water samples to ascertain *E. coli*, *Enterococci* and *Salmonella*. The Chromogenic (EC X-GLUC Agar) was used for the detection of *E. coli* (ISO 9308-1). The conformity test was performed with triptic soya agar. The culture and isolation of *Enterococci* was carried out by filtration of 100 ml of water subsequent culturing on Slanetz and Bartley agar (ISO 7899-2).

The conformity test for *Enterococci* was done using Bile Esculin Azide agar. For detection of *Salmonella* in 1 liter Hektoen Enteric Agar, Rappaport vasillaidis Broth were used after passing the filtered paper in to enrichment media containing buffered peptone (**ISO 6579-1981**). Oxoid *salmonella* testing kits were used pathogenic conformation. Total Bacterial count (**Heterotrophic Plate Count, APHA 9215-B**) was processed through pour plate technique by pouring 1 ml sample into Petri dishes and adding yeast extract agar as a growth medium accordingly. All the bacterial cultures were aided by incubating the relevant Petri dishes to required growth temperatures and their number was counted on total Colony forming units (CFU) basis.

### Physico-chemical investigation

The pH of the each sample was determined using a pH meter (**Adwa, AD 1020**) equipped with a temperature probe (Degree of accuracy 0.01). Turbidity was determined with Turbidimeter (Velp Scientifica, Model TB- 1). Nephelometric Turbidity Units (NTU) were taken into consideration. Similarly, conductivity meter (model AD 3000, Adwa) used to determine electrical conductivity and TDS. In accordance with the ISO standards reference solutions were used for calibration of equipment and accuracy of readings (0.01).

### Spectrophotometry

Spectrophotometer (**Techcomp, UV, dual beam, Model UV2300 II**) was used for assessment of the fallowing parameters;

### Total Nitrogen

The sum of nitrate nitrogen, nitrite nitrogen, ammonium nitrogen and organically bonded nitrogen, each determined separately. Ammonium, nitrite and organic nitrogen are oxidised to nitrate using potassium peroxodisulphate in boric acid-sodium hydroxide buffer. The oxidation of the nitrogen compounds is performed in an autoclave at 120 °C, resulting in a pH change of the buffer from 9.7 to 5.0. The resulting nitrate was determined by

spectrophotometry at 220 nm (APHA, AWWA, WEF, 1998; Valderrama, 1981).

#### **Ammonium**

Pipetted 25 ml of the sample into the 50 ml polypropylene Erlenmeyer flask with screw cap and added respective reagents. Three blanks samples were prepared per batch, using 25 mL of ultra-pure water in place of the test sample. The absorbance was set at 690nm (Fresenius et al. 1988; Grasshoff and Johannsen 1972).

#### **Total Phosphorous**

The spectrophotometric determination was performed at the wavelength of 890 nm with 5 cm cuvettes optical path and the spectrophotometer was zeroed by using de-ionised water without reagents. This procedure details the determination of total phosphorus in natural freshwater, surface water, and drinking water, in the form of ortho-phosphate and organic phosphorus compounds capable of conversion to ortho-phosphate under the oxidative digestion procedure described (Tartari and Mosello 1997; Valderrama 1981).

#### **Reactive Orthophosphate**

After mixing reagents with each 25 ml sample and preparing three blanks and repeatability the spectrophotometric determination was performed at the wavelength of 890 nm with 5 cm cuvettes optical path and the spectrophotometer is zeroed by using de-ionised water without reagents (Tartari and Mosello 1997; Valderrama 1981).

#### **Ion Chromatography**

Through ion chromatography (Dionex ICS 30000) following Anions and Cations have been ascertained. The said analysis has been carried out at Chemistry Lab, Institute of Ecosystem Studies, Council of National Research (CNR), Verabania, Italy. **Anions:** chlorides, nitrates, sulfates; (ISO 10304-1. Water quality - Determination of dissolved fluoride, chloride, nitrite, orthophosphate, bromide, nitrate and sulphate ions, using liquid chromatography of ions. Part. 1: method for water with low contamination). **Cations:** sodium, potassium, magnesium and calcium. (ISO 14911 Water quality - determination of dissolved Na<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, and using ion chromatography - method for water and waste water.)

### **3. RESULTS AND DISCUSSION**

#### **Bacteriology**

Results for E. coli, Enterococci and TBE from the valleys showed some declining trend from source to last point (table 1).

#### **Physical parameters**

Turbidity was recorded higher in some samples (table 2); however other indicators taken into account did not show any alarming situation in the study area. Some points exhibited little higher TDS while rest of locations remained in the range of given standards.

**Table 2: Physical Parameters of Source Mid and Last Points (Means followed by different letter(s) in the same column are significantly different from one another at  $LSD \leq 0.05$ )**

Valleys	Points	Temp	pH	Turbidity	Conductiv	TDS
Braldo	Source	1.56B	6.96A	7.92B	320.00A	170.75A
	Mid	1.73AB	6.97A	16.303AB	335.83A	165.92A
	Last	1.81A	7.03A	23.58A	307.67A	164.25A
	LSD	0.2442	0.0921	11.378	33.443	9.4311
Basha	Source	1.85B	7.01A	8.69A	229.39AB	161.37B
	Mid	1.92A	6.98A	7.14A	260.61A	187.21AB
	Last	1.85B	6.95A	8.95A	213.13B	212.46A
	LSD	0.0646	0.0949	2.1658	31.389	39.706

### Chemistry

Details are expressed in table 3 and 4 given below showed a normal ranges of the chemical ingredients tested from all points. Both valleys (table 3, Basha valley; table 4, Baraldu valley) vary slightly from one to another in some cases; however there was no any major point of concern or deviation from the international acceptable standards.

### 4. DISCUSSION

Keeping in view WHO guidelines, water used for drinking purposes was unsafe. Contamination level was alarming in streams and surface water as compare to the spring sources. Our results were in conformation with Chapman et al. (1971), who disapproved McCarisson myth of increasing water contamination increases risk of goiter. Similarly, bacteriological investigations are in closer agreement with the studies of Craun et al 2003, where the *E.coli* has been termed as the organism of choice for monitoring and surveillance programs in addition to a disinfection indicator.

The physico-chemical parameters however haven't shown any unusual trend, in most of the samples ph level was ranging from 6.5 to 7.5 which are acceptable and the ratio of total dissolved solids and conductivity was within the limit. Average turbidity of the samples was higher. Apart from the aesthetic value of water, presence of certain silt and sand particles may pose threat of gastric disorders and precursor behind kidney stone. The ratio of patients suffering from renal infections is considerable in the region.

Current research findings have encouraged stakeholders to establish a permanent water quality monitoring system to ensure provision of safe drinking water within CKNP of Karakoram Mountain Ranges, Pakistan. Further, the type and status of drinking water distribution system can be rectified if the source of contamination is along the medium from Head to tail. The identification of an alternative source (spring) and its possible use for the local inhabitants is possible with addition of allied sources in to the database, especially in the villages where stream water is the prime source of drinking.



The population residing within CKNP especially the areas

thankful for CKNP management team for their ongoing support during field visits.

**Table 3: Chemical Parameters Source, Mid and Last Points Bhasha Valley (Means followed by different letter(s) in the same column are significantly different from one another at  $LSD \leq 0.05$ )**

Points	T.Alk	Cl	SO <sub>4</sub>	N-NO <sub>3</sub>	N-NH <sub>4</sub>	Ca	Mg	Na	K	RP	TP	TN	SI	F
Source	1.27A	0.27B	32.81A	511.43A	6.71A	30.94A	5.61A	1.62A	4.30A	6.57A	12.85A	0.65A	2.34A	0.36A
Mid	1.32A	0.38A	34.31A	542.00A	5.714A	1.61A	5.94A	1.71A	4.24A	7.00A	11.71A	0.62B	2.15B	0.31A
Last	1.32A	0.32AB	33.92A	492.14A	5.14A	31.30A	5.81A	1.81A	4.20A	6.2A	11.14A	0.63AB	2.16B	0.29A
LSD	0.07	0.09	1.66	55.48	1.59	4.59	0.82	0.26	0.91	1.43	1.72	0.02	0.13	0.09

which remain aloof from the primary and secondary health care facilities during prolonged winter may become the chronic patients carrying gastrointestinal infestations. There is a dire need to address the personal hygiene as well as the pattern of open defecation near the sources which may lead to a worst case scenario in shape of mild to

**Table 4: Chemical Parameters Source Mid and Last Points Braldo Valley (Means followed by different letter(s) in the same column are significantly different from one another at  $LSD \leq 0.05$ )**

Points	T.Alk	Cl	SO <sub>4</sub>	N-NO <sub>3</sub>	N-NH <sub>4</sub>	Ca	Mg	Na	K	RP	TP	TN	SI	F
Source	1.24A	0.67A	80.47A	274.25A	5.33A	52.75A	29.84A	3.82A	16.18A	2.41A	4.66A	0.33A	4.50A	0.47A
Mid	1.21AB	0.63B	75.87B	272.83A	4.91A	50.57B	28.18AB	3.60AB	15.69A	2.54A	4.79A	0.33A	4.37A	0.45B
Last	1.19B	0.60C	74.16B	270.33A	4.70A	50.57B	27.87B	3.35B	15.11B	2.41A	4.75A	0.32A	4.13A	0.44C
LSD	0.03	0.02	3.06	9.17	0.76	1.95	1.91	0.43	0.56	0.33	0.69	0.01	0.49	0.01

severe and chronic ailments.

### Acknowledgements

Study is part of PhD research and was possible due to generous financial grant from Italian government and the government of Pakistan through SEED project in collaboration of Karakoram International University, Gilgit-Baltistan. Authors are indebted to Mr. Gabriele Tartari of water chemistry lab and his team, Institute of Ecosystem Studies, Council of National Research, Italy. We are also

### References

1. APHA, AWWA, WEF. 1998. Standard Methods for the Examination of Water and Wastewater (Method 4500 NO<sub>3</sub><sup>-</sup> B, Ultraviolet spectrophotometric method). Amer. Publ. Health Ass., Washington. Academic Press, San Diego.

2. APHA, AWWA, WEF. 1998. Standard Methods for the Examination of Water and Wastewater (Method 4500 N C, Persulfate method). Amer. Publ. Health Ass., Washington.
3. Acreman M. 2001. Ethical aspects of water and ecosystems. Water Policy, 3(3), 257-265.
4. Ahmed K, Shakoori AR. 2002. Vibrio cholerae E1 Tor, Ogawa 01, as the main etiological agent of the two major outbreaks of gastroenteritis in the Northern Areas of Pakistan. Pak. J. Zool. 34(1), pp.73-80
5. Ahmed S, Joyia F. 2003. Northern Areas Strategy for sustainable Development, pp.67.
6. APHA. 2000. American Public Health Association, Standard Methods for the Examination of Water and Waste water 20th Edition. Part 9000, PP: 9-40.
7. Birks JS. 1977. The Reaction of Rural Populations to Drought: A Case Study from South East Arabia (Die Reaktion ländlicher Bevölkerungen auf Dürre: eine Fallstudie aus Südost-Arabien). Erdkunde, 299-305.
8. Chapman JA, Grant IS, Khalida M, Shahid MA. 1972. Endemic Goiter in the Gilgit Agency. Philosophical trends of the royal society of London. Bioscience, Vol 263, No 856.
9. Chilando M, Kelderman P, Okeeffe H. 2008. Design of a water quality monitoring network of the limpopoo river, Banin, Mozambique. Physics and chemistry of earth (Report)
10. Craun GF, Calderon RL, Nwachuku N. 2003. Causes of Waterborne Outbreaks Criteria and Other Supporting Information. International Programme on Chemical.
11. Cropper M, Griffiths C. 1994. The interaction of population growth and environmental quality. The American Economic Review, 250-254.
12. Fresenius W, Quentin KE, Schneider W. 1988. Water Analysis. Springer-Verlag, Berlin. 804 pp.
13. Goel PK. 2006. Water Pollution - Causes, Effects and Control. New Delhi: New Age International. p. 179. ISBN 978-81-224-1839-2
14. Grasshoff K, Johannsen H. 1972. A new sensitive and direct method for the determination of ammonia in sea water. J.Cons. Perm. Int. Explor. Mer., 34: 516-521
15. ISO 7899-2. International Organization for Standardization, 1999 (7899-2) Water Quality Detection and Enumeration of Intestinal Enterococci, Part2
16. ISO 7704. International Organization for Standardization (7704) Evaluation of membrane used for microbiological analysis.
17. ISO 9308-1. International Organization for Standardization. (9308-1) Water Quality - Detection and Enumeration of Escherichia coli and coliform Bacteria-Part1
18. Mara DD, Cairncross S. 1989. Guidelines for the safe use of wastewater and excreta in agriculture and aquaculture: measures for public health protection.
19. Rosegrant MW, Cai X, Cline SA. 2002. World water and food to 2025: dealing with scarcity. Intl Food Policy Res Inst.
20. Valderrama JC. 1981. The simultaneous analysis of total nitrogen and total phosphorus in natural waters. Marine Chemistry, 10: 109 - 122.
21. WHO. 2004b. Guidelines for drinking -Water Quality. Third Edition, Vol.1: Recommendation. World Health Organization, Geneva.
22. WHO. 2000. World Health organization.. Monitoring Bathing Waters - A Practical Guide to the Design and Implementation of Assessments and Monitoring Programmes. 106-157 p.
23. WHO. 2013. World Health Organization. Press release sg/sm/8707 obv/348. 'water-related diseases responsible for 80 per cent of all illnesses, deaths. In developing world', says secretary-general in environment day message. Retrieved from on august 7, 2014.

4/2/2025