

**Mouse Anti-Blood Group Antigen B [HEP-20]: MC0006**

**Intended Use:** For Research Use Only

**Description:** The antibody HEB-20 reacts with human blood group B. The specificity of the antibody HEB-20 was confirmed by comparison of specificity and reactivity to standard reagent using >5.000 samples of blood. The antibody HEB-20 shows specific staining of erythrocytes and vascular epithelium of blood group B controls and no staining in group A controls. This antibody is applicable for tissue staining in tumor patients with blood groups B and AB. Blood group antigens are generally defined as molecules formed by sequential addition of saccharides to the carbohydrate side chains of lipids and proteins detected on erythrocytes and certain epithelial cells. The A, B and H antigens are reported to undergo modulation during malignant cellular transformation. Blood group related antigens represent a group of carbohydrate determinants carried on both glycolipids and glycoproteins. They are usually mucin type, and are detected on erythrocytes, certain epithelial cells, and in secretions of certain individuals. Sixteen genetically and biosynthetically distinct but inter related specificities belong to this group of antigens, including A, B, H, Lewis A, Lewis B, Lewis X, Lewis Y, and precursor type 1 chain antigens.


**Specifications:**

Clone: HEP-20  
 Source: Mouse  
 Isotype: IgM/k  
 Reactivity: Human  
 Localization: Membrane  
 Formulation: Protein A/G purified antibody from bioreactor concentrate. Prepared in 10mM PBS with 0.2% BSA and < 0.09% sodium azide (NaN3).  
 Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles  
 Applications: IHC, ICC/IF  
 Package:

Description	Catalog No.	Size
Blood Group Antigen B Concentrated	MC0006	1 ml

**IHC Procedure\*:**

Positive Control Tissue: KG1 cells, human colorectal carcinoma tissues  
 Concentrated Dilution: 50-200  
 Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C  
 Incubation Time and Temp: 30-60 minutes @ RT  
 Detection: Refer to the detection system manual  
 \* Result should be confirmed by an established diagnostic procedure.



FFPE human colon carcinoma stained with anti-Blood Group Antigen B using DAB

Doc. 100-MC0006  
Rev. A