

Rabbit Anti-FSH (Follicle Stimulating Hormone) [EP257]: RM0292, RM0292RTU7

Intended Use: For Research Use Only

Description: The pituitary glycoprotein hormone family includes follicle-stimulating hormone, luteinizing hormone, chorionic gonadotropin, and thyroid-stimulating hormone. All of these glycoproteins consist of an identical alpha subunit and a hormone-specific beta subunit. FSH beta is the beta subunit of follicle-stimulating hormone. In conjunction with luteinizing hormone, follicle-stimulating hormone induces egg and sperm production. The FSH antibody specifically labels Gonadotrophs in pituitary. It may be useful in the classification of pituitary tumors.

Specifications


Clone: EP257
Source: Rabbit
Isotype: IgG
Localization: Cytoplasm
Formulation: Antibody in PBS pH7.2, containing < 1% BSA and 0.09% sodium azide (NaN₃).
Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles
Applications: IHC for FFPE, Frozen
Package:

Description	Catalog No.	Size
FSH Concentrated	RM0292	1 ml
FSH Prediluted	RM0292RTU7	7 ml

IHC Procedure*

Positive Control Tissue: Pituitary, pituitary carcinoma
Concentrated Dilution: 50-200
Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
Incubation Time and Temp: 30-60 minutes @ RT
Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human pituitary stained with anti-FSH using DAB

References:

1. Isolated FSH deficiency revealing a granulosa cell tumor. Decoudier B, et al. Ann Endocrinol (Paris). Dec;71(6):543-7, 2010.
2. Early alterations in ovarian surface epithelial cells and induction of ovarian epithelial tumors triggered by loss of FSH receptor. Chen X, et al. Neoplasia. Jun;9(6):521-31, 2007.
3. Assays for LH, FSH, and prolactin. Wheeler MJ. Methods Mol Biol. 324:109-24, 2006.