

P. 2/3

Because of your hip how much difficulty do you have with:

	No difficulty at all	Slight difficulty	Moderate difficulty	Extreme difficulty	Unable to do	N/A
Twisting/pivoting on involved leg	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Rolling over in bed	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Light to moderate work (standing, walking)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Heavy work (push/pulling, climbing, carrying)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Recreational activities	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

How would you rate your current level of function during your usual activities of daily living from 0 to 100 with 100 being your level of function prior to your hip problem and 0 being the inability to perform any of your usual daily activities?

.0 %