

LA LUZ ECC SICK CHILD FORM updated 8-7-25

Child's Name: _____

Date: _____

Your child had the following indications of not feeling well today _____

Child's Temperature _____

Time: _____

Person Making Report: _____

Admin Initial _____

A check mark next to a symptom means that the responsible parent/guardian needs to follow instructions:

- | | |
|--|--|
| <ul style="list-style-type: none"><input type="checkbox"/> Diarrhea or vomiting. The child may return when free from symptoms for 48 hours. Child will be sent home if they have a watery stool that cannot be contained in diaper or underpants or 3 or more soft, loose stools in a 4-hour period or 1 time vomiting.<input type="checkbox"/> Impetigo. The child may return to school after antibiotics have been administered for 24 hours, or when sores are very dry with no yellow crust, and no longer oozing (about 7-10 days), with a doctor's note and absence of other symptoms.<input type="checkbox"/> Fever. Temperatures above 100.4F or 99.4F (axillary) under the arm. Child may return when free from fever for 48 hours without the aid of fever reducing medicine.<input type="checkbox"/> General Malaise. A child who does not eat or have the energy and requires 1:1 attention does not belong at school until fully able to participate.<input type="checkbox"/> Earache. The child should be seen by a physician and return to school following a minimum of 24 hours of treatment and absence of other symptoms.<input type="checkbox"/> Flu. If a child is diagnosed with the Flu, they should remain out of school for 48 hours after symptoms (such as fever, vomiting, diarrhea, etc.) subside without the aid of fever reducing or anti-nausea medication and they are feeling well enough to fully participate in the day's activities.<input type="checkbox"/> Pediculosis (head lice). The child may return to school following a minimum of two days of treatment and no nits are observed by staff upon return. | <ul style="list-style-type: none"><input type="checkbox"/> Cold. The child may attend school if he or she is free from fever and general malaise for 48 hours. If a child has a lingering cough (a signal of an upper respiratory infection), he or she should be seen by a physician and will require a negative COVID test. If positive for COVID, child should remain out of school until they are 48 hours symptom free (not including a lingering cough – if cough hinders sleep, eating, running, etc. they should remain out until it no longer affects their normal activities) without the aid of medication. If older than 2 years of age, it is recommended that they wear a mask for 5 days upon return.<input type="checkbox"/> Ringworm or Athlete's Foot. The child may attend if the infected area is being treated for 24 hours and is covered.<input type="checkbox"/> Strep Throat. The child may return to school after 24 hours of treatment with medication and child has no fever for 48 hours.<input type="checkbox"/> Conjunctivitis (pinkeye). The child may return to school following 24 hours of treatment with prescribed medication.<input type="checkbox"/> Rash. Out of care if child has fever or until a doctor determines that it is not contagious (<u>must have a note from the doctor</u>). In some cases, such as Hand, Foot and Mouth Disease (or other illnesses which cause blisters filled with fluid), children may not return for a minimum of 3 days (without fever for at least 48 hours), no fluid filled blister, no active mouth sores and no excessive drooling, regardless of doctor's note. |
|--|--|

Parent/Guardian Signature

Date