Main Office:

Scottsdale Vein & Proctology Center
8752 E Via De Commercio #2
Scottsdale, Arizona 85258
Office: (602) 492-9919 | Mobile: (602) 920-1023

R I C K   A   S H A C K E T
D O ,   M D (H)

Name: _____ DOB: _____ Date: _____

DISPENSE ONE OF THE FOLLOWING FOUR PRESCRIPTIONS:

1) First Choice:
Analpram HC Cream 2.5% 30 X 4G Singles (Hydrocortisone Acetate 2.5%, Pramoxine HCL 1%)
- Contains #30- Net Wt. 4g single use tubes with applicators.
- Use 3-4 times daily or as directed by your physician.
- **10 day supply (30 4g Tubes), 11 Refills**

2) Second Choice:
Analpram HC Cream 2.5% 1 OZ Tube (Hydrocortisone Acetate 2.5%, Pramoxine HCL 1%)
- Dispense preferably with a rectal applicator
- Apply to affected area 3-4 times daily
- (Hydrocortisone Acetate 2.5%, Pramoxine HCL 1% Generic Available)
- **10 day supply (1 OZ Tubes), 11 Refills**
- May dispense 90 day supply: At patient's option, dispense up to nine 1-oz tubes

3) Third Choice:
Generic 2.5% Hydrocortisone cream 1 oz.
- Dispense preferably with a rectal applicator
- Apply to affected area 3-4 times daily
- **10 day supply (1 OZ Tubes), 11 Refills**
- May dispense 90 day supply: At patient's option, dispense up to nine 1-oz tubes

4) Fourth Choice:
Hydrocortisone acetate 1% and pramoxine hydrochloride 1% aerosol foam
- Insert 1 applicatorful into the rectum and or apply with tissue around the anal area, 2 or 3 times daily and after each bowel evacuation.
- Dispense 4-canisters per month (i.e. 1- canisters per 7 days)
- **11 Refills** of 4-canisters. When used correctly, the aerosol container will deliver a minimum of 14 applications)

_____
Rick A. Shacket, DO, MD (H), BS9262611
Diplomate American Osteopathic Board of Proctology