

Rabbit Anti-CD73/NT5E [D7F9A]: RM0273

Intended Use: For Research Use Only

Description: Ecto-5'-nucleotidase (NT5E, also called CD73) is a 70 kDa glycosyl phosphatidylinositol-anchored, membrane-bound glycoprotein that catalyzes the hydrolysis of extracellular nucleoside monophosphates into bioactive nucleosides. NT5E catalyzes the terminal step of extracellular adenosine formation from adenosine monophosphate, which drives the regulation of extracellular adenosine levels and the downstream activation of the four G protein-coupled adenosine receptors. Binding of hypoxia-inducible factor (HIF-1) to the NT5E gene promoter leads to upregulation of NT5E during hypoxia (3). The biological roles of NT5E include lymphocyte adhesion, fibrosis (6), and the regulation of nociception. NT5E/CD73 (D7F9A) Rabbit mAb recognizes endogenous levels of total NT5E/CD73 protein.

Specifications:

Clone: D7F9A
Source: Rabbit
Isotype: IgG
Reactivity: Human, mouse, rat
Localization: Membrane
Formulation: Antibody in PBS pH7.4, containing BSA, glycerol, and ≤0.09% sodium azide (NaN₃).
Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles
Applications: IHC, Flow Cyt., WB
Package:

Description	Catalog No.	Size
CD73/NT5E Concentrated	RM0273	1 ml

IHC Procedure*:

Positive Control Tissue: Lung ca, liver, kidney
Concentrated Dilution: 20-40
Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
Incubation Time and Temp: 30-60 minutes @ RT
Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.