## Kittitas County Prehospital Care Protocols

Subject: ASPIRIN (Off-line Medical Direction) - EMT

## INDICATIONS FOR USE IN AN ACUTE CORONARY EVENT:

- 1) Patient exhibits any of the following signs or symptoms:
  - a) Uncomfortable pressure, fullness, squeezing or pain in the center of the chest that lasts more than a few minutes, or goes away and comes back.
  - b) Pain that spreads to the shoulders, neck or arms.
  - c) Chest discomfort with light headedness, fainting, sweating, nausea or shortness of breath.

-OR-

- 2) Patient exhibits any <u>two</u> of the following signs or symptoms, and you think it is of cardiac origin:
  - a) Atypical chest pain, stomach or abdominal pain. This may include discomfort that can be localized to a point, that is "sharp" in nature, that is reproducible by palpation, or that is in the "wrong" location (such as the upper abdomen).
  - b) Unexplained nausea (without vomiting) or lightheadedness (not vertigo) without chest pain.
  - c) Shortness of breath and difficulty breathing (*with or* without chest pain).
  - d) Unexplained anxiety, weakness or fatigue.
  - e) Palpitations, cold sweat or paleness.

## **CONTRAINDICATIONS FOR USE:**

- 1) Patient is allergic to aspirin or ibuprofen (Motrin®, Advil®).
- 2) If the patient has just taken *the recommended dose* of aspirin for this event, do not administer aspirin.
- 3) Radial pulses are not present in both arms or systolic BP varies by at least 20 mm/Hg.
- 4) Head injury suspected (includes stroke) Check w/ medical control if uncertain

## PROCEDURE:

- 1) ALS/ILS upgrade and evaluation required unless ALS/ILS is unavailable.
- 2) Be sure that the patient is alert and responsive.
- 3) If the patient has his/her own nitroglycerin and meets the criteria for administration, do not delay assisting in the administration per protocol.
- 4) Check radial pulses in both arms. If both are present *and systolic BP is within < 20 mm/Hg between arms*, have the patient chew four 81 mg aspirin or equivalent (324 mg). "Chewable" aspirin are recommended.
- 5) Record all treatment, including the dosage and the time of administration.

Effective Date:	April 16, 2012	

Medical Program Director: Jackson S. Horsley, MD