

Mouse Anti-PR [PR484]: MC0920, MC0920RTU7

Intended Use: For Research Use Only

Description: The human progesterone receptor (PR), is a ligand-activated transcription factor and is a member of the steroid receptor family. PR exists in humans as two isoforms. PR is predominantly expressed in female sex steroid responsive tissues such as the mammary gland, uterus and ovary, but is also found in other tissues such as prostate stromal cells, anterior pituitary gland, and endocrine cells of the Langerhans' islets.

Specifications


Clone: PR484
 Source: Mouse
 Isotype: IgG1k
 Reactivity: Human
 Localization: Nucleus
 Formulation: Tissue culture supernatant in PBS pH7.5, containing 0.2% BSA, 15mM sodium azide (NaN3)
 Storage: Store at 2°- 8°C.
 Applications: IHC
 Package:

Description	Catalog No.	Size
PR Concentrated	MC0920	1 ml
PR Prediluted	MC0920RTU7	7 ml

IHC Procedure*

Positive Control Tissue: Breast cancer
 Concentrated Dilution: 50-200
 Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human breast ca. stained with anti-PR using DAB

References:

1. Confirmation of the progesterone receptor as an efficient marker of treatment with 17β-estradiol in veal calves. Pezzolato M, et al. Food Addit Contam Part A Chem Anal Control Expo Risk Assess. 33(1):60-5, 2016.
2. Expression of functional toll like receptor 4 in estrogen receptor/progesterone receptor-negative breast cancer.
3. Mehmeti M, et al. Breast Cancer Res. ep 22;17:130, 2015.
4. The relationship among HOXA10, estrogen receptor α, progesterone receptor, and progesterone receptor B proteins in rectosigmoid endometriosis: a tissue microarray study. Zanatta A, et al. Reprod Sci. Jan;22(1):31-7, 2015.