

HOWARD NASS MD FAAP

DIPLOMATE AMERICAN BOARD OF PEDIATRICS

76-01 113 STREET
FOREST HILLS, NY 11375
718 520 1070

www.doctornass.com

IBUPROFEN (motrin or advil) Dosage (up to every 6 hrs if needed)

(This table shows the highest recommended dose for the low end of each weight range)

weight	dosage	drops 50mg/1.25ml (use dropper provided)	suspension 100mg/5ml	chewable	Chewable or tabs	tablets or caps
pounds	mg	ml	teaspoon (milliliter)	50mg each	100mg each	200mg each
11-16	50	1.25 ml	1 / 2 tsp (2.5ml)	X	X	X
17-21	75	1.875 ml	3 / 4 tsp (3.75 ml)	X	X	X
22-32	100	2.5 ml	1 tsp (5 ml)	2	1	X
33-43	150	3.75 ml	1- 1 / 2tsp (7.5ml)	3	X	X
44-54	200	5 ml	2 tsp (10 ml)	4	2	1
55-65	250	X	2 – 1 / 2 tsp (12.5 ml)	5	2 – 1 / 2	1
66-87	300	X	3 tsp (15 ml)	6	3	1 -1/2
88 And up	400	X	4 tsp (20 ml)	8	4	2

(Dosage in table is based on prescription dosage of 10mg/kg/dose)