

Rabbit Anti-Calcitonin [SP17]: RM0222, RM0222RTU7

Intended Use: For Research Use Only

Description: Calcitonin is a 32-amino acid polypeptide hormone that is commonly expressed in the parafollicular C cells in the thyroid gland. Calcitonin is a potent plasma calcium-lowering peptide; it decreases the level of calcium and phosphate in blood by promoting the incorporation of these ions in bones. Calcitonin antibody is useful for the identification of C cell hyperplasia and medullary thyroid carcinomas.

Specifications


Clone: SP17
 Source: Rabbit
 Isotype : IgG
 Localization: Cytoplasm
 Formulation: Antibody in PBS pH7.2, containing BSA and $\leq 0.09\%$ sodium azide (NaN₃)
 Storage: Store at 2°- 8°C
 Applications: IHC
 Package:

Description	Catalog No.	Size
Calcitonin Concentrated	RM0222	1 ml
Calcitonin Prediluted	RM0222RTU7	7 ml

IHC Procedure*

Positive Control Tissue: Thyroid, medullary carcinoma of thyroid
 Concentrated Dilution: 50-100
 Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human thyroid medullary carcinoma stained with anti-Calcitonin using DAB

References:

1. Profiling of the calcitonin-calcitonin receptor axis in primary prostate cancer: clinical implications and molecular correlates. Thakkar A, et al. *Oncol Rep. Sep;30(3):1265-74, 2013.*
2. Calcitonin gene-related peptide is a promising marker in ulcerative colitis. Li FJ, et al. *Dig Dis Sci. Mar;58(3):686-93, 2013.*
3. Somatostatin, substance P and calcitonin gene-related peptide-positive intramural nerve structures of the human large intestine affected by carcinoma. Godlewski J, et al. *Folia Histochem Cytobiol. Sep 30;48(3):475-83, 2010.*