

POST-OPERATIVE INSTRUCTIONS

FOLLOWING OFFICE SURGERY

1. KEEP THE RECTAL AREA CLEAN AND DRY. After a bowel movement, clean your bottom with peri-anal cleansing pads (i.e. AloeClean®, Tucks® or **baby-wipes**), followed by blotting the area dry with white unscented toilet paper. Change your dressing after each bowel movement, or whenever soiling occurs.

Dressing: A piece of cotton (from a **cotton roll**) or non-sterile 4x4 gauze pads, placed between your buttocks and held with white paper tape, will help keep the area dry. A sanitary napkin worn inside your underwear can also help absorb excess drainage. Dressing the anal area does not effect healing, but will help to keep your underwear and bedding clean and dry.

2. REDUCE SWELLING. A sitz bath (warm water bath) several times a day will give you soothing relief. Ice the area for 10–15 min (an ice cube in a Ziploc bag works fine) at least 2 – 3 times a day for 2 to 5 days after surgery, to reduce swelling and aid in healing.

3. CALL OUR OFFICE TO MAKE AN APPOINTMENT. We definitely want to see you in approximately 100 days from the date of your operation. If you have any symptoms that are not improving, we would like to see you sooner.

4. TAKE YOUR MEDICINE AS PRESCRIBED.

Hydrocortisone/Proxamine: For swelling, inflammation, and itching of the anal tissue. Important: use hydrocortisone cream and or suppositories several times a day until you are healed. This will shorten your healing time by as much as 50%.

Hydromorphone/Oxycodone/Hydrocodone/Codeine: For pain. Do not drink or drive on this medication.

Casanthranol/Docusate/Senna/Magnesium hydroxide: To soften the stool and gently stimulate a bowel movement.