

**Rabbit Anti-p21WAF1 [EP147]: RM0152**

**Intended Use:** For Research Use Only

**Description:** p21 / WAF1, also known as cyclin-dependent kinase inhibitor 1 or CDK-interacting protein 1, is a protein that in humans is encoded by the CDKN1A gene located on chromosome 6 (6p21.2). The p21 protein binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at G1. p21 is expressed in all adult human tissues. In tumors, the expression of p21 has been studied by immunohistochemical methods in a wide range of human tumors, such as gastric carcinoma, non-small cell lung carcinoma, and thyroid carcinoma. The expression of p21 is associated with favorable prognosis in various tumors.

**Specifications:**

Clone: EP147  
Source: Rabbit  
Isotype: IgG  
Reactivity: Human  
Localization: Nucleus  
Formulation: Antibody in PBS pH7.4, containing BSA, glycerol, and ≤0.09% sodium azide (NaN<sub>3</sub>).  
Storage: Store at 2°- 8°C. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles  
Applications: IHC  
Package:

Description	Catalog No.	Size
p21WAF1 [EP147] Concentrated	RM0152	1 ml

**IHC Procedure\*:**

Positive Control Tissue: colon cancer  
Concentrated Dilution: 50-200  
Pretreatment: Citrate pH6.0 or EDTA pH8.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C  
Incubation Time and Temp: 30-60 minutes @ RT  
Detection: Refer to the detection system manual

\* Result should be confirmed by an established diagnostic procedure.