

ER/LA OPIOID REMS:

Achieving Safe Use While Improving Patient Care

Presented by CO*RE
Collaboration for REMS Education
www.core-rems.org



Collaborative for
REMS Education

Faculty Information



Doctor Carl Christensen is the Medical Director of the Michigan Health Professional Recovery Program which monitors impaired physicians, nurses and pharmacists, and the Medical Director of the Eleonore Hutzel Women's Recovery Center, an outpatient program dedicated to caring for pregnant, chemically dependent women. Doctor Christensen is also the Medical Director at the Tolan Medical Research Clinic in the Department of Psychiatry at Wayne State University and a Clinical Associate Professor in the department of OB GYN at Wayne State.

Doctor Christensen obtained his MD and PhD in Biochemistry at Wayne State University School of Medicine.

DISCLOSURE:

The content of this presentation is non-commercial and does not represent any conflict of interest.



Collaborative for REMS Education

On July 9, 2012, the Food and Drug Administration (FDA) approved a Risk Evaluation and Mitigation Strategy (REMS) for extended-release (ER) and long-acting (LA) opioid medications.

Founded in June, 2010, the Collaborative on REMS Education (CO*RE), a multi disciplinary team of 10 partners and 3 cooperating organizations, has designed a core curriculum based on needs assessment, practice gaps, clinical competencies, and learner self-assessment to meet the requirements of the FDA REMS Blueprint.

www.core-rems.org

CORE* Organizations

Founding Partners

- ◆ American Pain Society (APS)
- ◆ American Academy of Hospice and Palliative Medicine (AAHPM)
- ◆ American Association of Nurse Practitioners (AANP)
- ◆ American Academy of Physician Assistants (AAPA)
- ◆ American Osteopathic Association (AOA)
- ◆ American Society of Addiction Medicine (ASAM)
- ◆ California Academy of Family Physicians (CAFP)
- ◆ Healthcare Performance Consulting (HPC)
- ◆ Interstate Postgraduate Medical Association (IPMA)
- ◆ Nurse Practitioner Healthcare Foundation (NPHF)

Strategic Partners

- ◆ Physicians Institute for Excellence in Medicine which coordinates 15 state medical societies
- ◆ Medscape
- ◆ American Academy of Family Physicians
- ◆ American College of Emergency Physicians (***New in 2015***)




American Academy of Hospice and Palliative Medicine



AMERICAN OSTEOPATHIC ASSOCIATION
TREATING OUR FAMILY AND YOURS



RESEARCH
EDUCATION
TREATMENT
ADVOCACY



ADVANCING EMERGENCY CARE
Collaborative for REMS Education

Content Development/Planner/Reviewer Disclosures

The following individuals disclose no relevant financial relationships:

David Bazzo, MD	Professor of Family Medicine, University of California San Diego School of Medicine
Roberto Cardarelli, DO, MPH	Professor, Department of Family and Community Medicine, University of Kentucky College of Medicine COI NOTE: While Dr. Cardarelli is PI on a Pfizer grant, his employing institution (U of KY) receives those funds to compensate for Dr. Cardarelli's time.
Ronald Crossno, MD	Senior National Medical Director, Gentiva Health Services, Rockdale, TX
Katherine Galluzzi, DO	Professor and Chair, Department of Geriatrics, Philadelphia College of Osteopathic Medicine, Philadelphia, PA
Carol Havens, MD	Family physician and addiction medicine specialist, The Permanente Medical Group, Sacramento, CA
Edwin A. Salsitz, MD, FASM	Beth Israel Medical Center, Division of Chemical Dependency; Assistant Professor, Albert Einstein College of Medicine
Barbara St. Marie, PhD, ANP-BC	Supervisor, Pain and Palliative Care; Adult and Gerontology Nurse Practitioner, Pain Management, University of Minn Medical Center, Fairview, MN
Cynthia Kear, CHCP, MDiv Shelly Rodrigues, CAE, FACEHP	Senior Vice President, California Academy of Family Physicians, San Francisco, CA Deputy Executive Vice President, California Academy of Family Physicians
Robin and Neil Heyden	Staff, CO*RE Operations Team, Heyden TY, Alameda, CA
Julie Bruno, MSW LCSW	Director, Education and Training, American Academy of Hospice and Palliative Medicine, Chicago, IL
Anne Norman, DNP, APRN, FNP-BC	Associate Vice President of Education, American Association of Nurse Practitioners
Marie Michelle-Leger, MPH, PA-C Eric D. Peterson, EdM, FACEHP	Director, Clinical Education, American Academy of Physician Assistants, Alexandria, VA Senior Director, Performance Improvement CME, American Academy of Physician Assistants

CO*RE Staff Disclosures

The following individuals disclose no relevant financial relationships:

Mary Ewert, PhD Sharon McGill, MPH	Manager, Public Health, Department of Research and Development, American Osteopathic Association, Chicago, IL Director, Department of Quality and Research, American Osteopathic Association, Chicago, IL
Stephen Biddle, M Ed Catherine Underwood, MBA, CAE	Senior Education Manager, American Pain Society Chief Executive Officer, American Pain Society, Chicago, IL
Arlene Deverman, CAE, CFRE Penny Mills, MBA	Vice President, Professional Development, American Society of Addiction Medicine Executive Vice President and CEO, American Society of Addiction Medicine Chevy Chase, MD
Thomas McKeithen Jr, BS, MBA Chris Larrison	Partners, Healthcare Performance Consulting Inc., Fleming Island, FL
Kate Nisbet, BBA, MBA Mary Ales, BA	Director of Health Systems Education, Interstate Postgraduate Medical Association Executive Director, Interstate Postgraduate Medical Association, Madison, WI
Fionna Shannon, MHS, FNP Phyllis Zimmer, MN, FNP, FAAN	Director, NPHF Continuing Education Program President, Nurse Practitioner Healthcare Foundation, Bellevue, WA
Sara Bennett Adele Cohen, MS, PCMH CCE	Project Manager, Physicians' Institute for Excellence in Medicine Senior Vice President, Physicians' Institute for Excellence in Medicine, Atlanta, GA
Piyali Chatterjee Cyndi Grimes, CCMEP Sarah Williams, PhD	Director, Medical Education, Medscape, LLC New York ,NY CME/CE Director, Medscape, LLC, New York, NY Scientific Director, Medscape, LLC, New York, NY
Cynthia Singh Lori Foley	Director, Grants and Foundation Development, American College of Emergency Physicians Director, Strategic Partnerships, American College of Emergency Physicians, Irving, TX

Acknowledgement

Presented by Michigan State Medical Society, a member of the Collaborative on REMS Education (CO*RE), 13 interdisciplinary organizations working together to improve pain management and prevent adverse outcomes.

This educational activity is supported by an independent educational grant from the ER/LA Opioid Analgesic REMS Program Companies. Please see http://ce.er-la-opioidrems.com/lwgCEUI/rems/pdf/List_of_RPC_Companies.pdf for a listing of the member companies. This activity is intended to be fully compliant with the ER/LA Opioid Analgesic REMS education requirements issued by the US Food & Drug Administration.

Products Covered by this REMS

Brand Name Products

- Avinza[®] morphine sulfate ER capsules
- Butrans[®] buprenorphine transdermal system
- Dolophine[®] methadone hydrochloride tablets
- Duragesic[®] fentanyl transdermal system
- *Embeda[®] morphine sulfate/naltrexone ER capsules
- Exalgo[®] hydromorphone hydrochloride ER tablets
- Hysingla[®] ER (hydrocodone bitartrate) ER tablets
- Kadian[®] morphine sulfate ER capsules
- Methadose[™] methadone hydrochloride tablets
- MS Contin[®] morphine sulfate CR tablets
- Nucynta[®] ER tapentadol ER tablets
- Opana[®] ER oxymorphone hydrochloride ER tablets
- OxyContin[®] oxycodone hydrochloride CR tablets
- Targiniq[™] oxycodone hydrochloride/naloxone hydrochloride ER tablets
- Zohydro[®] hydrocodone bitartrate ER capsules


Generic Products

- Fentanyl ER transdermal systems
- Methadone hydrochloride tablets
- Methadone hydrochloride oral concentrate
- Methadone hydrochloride oral solution
- Morphine sulfate ER tablets
- Morphine sulfate ER capsules
- Oxycodone hydrochloride ER tablets

* Not currently available due to voluntary recall (still approved); † No longer marketed (still approved)

WHY PRESCRIBER EDUCATION IS IMPORTANT

Introduction



Prescribers of ER/LA Opioids Should Balance:

***The benefits
of prescribing
ER/LA opioids
to treat pain***



***The risks
of serious
adverse
outcomes***

ER/LA opioid analgesics should be prescribed only by health care professionals who are knowledgeable in the use of potent opioids for the management of pain

Opioid Misuse/Abuse is a Major Public Health Problem

Improper use of any opioid can result in serious AEs including overdose & death

This risk can be greater w/ ER/LA opioids

ER opioid dosage units contain more opioid than IR formulations

Methadone is a potent opioid with a long, highly variable half-life

In 2012

37 million Americans age ≥ 12 had used an opioid for nonmedical use some time in their life

In 2011

488,004 ED visits involved nonmedical use of opioids
• Methadone involved in 30% of prescription opioid deaths

SAMHSA. (2013). *Results from the 2012 National Survey on Drug Use and Health: Detailed Tables*. NSDUH Series H-46, HHS Publication No. (SMA) 13-4795. Rockville, MD. SAMHSA. (2013). *Drug Abuse Warning Network, 2011: National Estimates of Drug-Related Emergency Department Visits*. HHS Publication No. (SMA) 13-4760, DAWN Series D-39. Rockville, MD. CDC. CDC Vital Signs. *Prescription Painkiller Overdoses. Use and abuse of methadone as a painkiller*. 2012. FDA. *Questions and Answers: FDA approves a Risk Evaluation and Mitigation Strategy for Extended-Release and Long-Acting Opioid Analgesics*. www.fda.gov/Drugs/DrugSafety/InformationbyDrugClass/ucm309742.htm. 2012.

In 2011

41,340 Americans DIED FROM DRUG POISONINGS

Nearly 17,000 deaths involved prescription opioids

In 2008

For every 1 death there are:



10 treatment admissions for abuse


32 ED visits for misuse or abuse

130 people who abuse or are addicted

825 nonmedical users

NCHS Data Brief, No. 166, September 2014. <http://www.cdc.gov/nchs/data/databriefs/db166.htm> (accessed on 1/6/15).
CDC. Policy Impact: Prescription Painkiller Overdoses. <http://www.cdc.gov/homeandrecreationalafety/rxbrief/> (Historical content - 2008 data) (accessed on 1/6/15).

First-Time Use of Specific Drugs Among Persons Age ≥ 12 (2012)



SAMHSA. (2013). *Results from the 2012 National Survey on Drug Use and Health: Summary of National Findings*. NSDUH Series H-46, HHS Publication No. (SMA) 13-4795. Rockville, MD.

Learning Objectives



Describe appropriate patient assessment for treatment with ER/LA opioid analgesics, evaluating risks and potential benefits of ER/LA therapy, as well as possible misuse.



Apply proper methods to initiate therapy, modify dose, and discontinue use of ER/LA opioid analgesics, applying best practices including accurate dosing and conversion techniques, as well as appropriate discontinuation strategies.



Demonstrate accurate knowledge about how to manage ongoing therapy with ER/LA opioid analgesics and properly use evidence-based tools while assessing for adverse effects.



Employ methods to counsel patients and caregivers about the safe use of ER/LA opioid analgesics, including proper storage and disposal.




Review/assess general and product-specific drug information concerning ER/LA opioid analgesics and identifying potential adverse effects of ER/LA opioids.

Misuse, abuse, divergence and overdose of ER/LA opioids is a major public health crisis.

YOU and **YOUR TEAM** *can* have an immediate and positive impact on this crisis while also caring for your patients appropriately.

ASSESSING PATIENTS FOR TREATMENT WITH ER/LA OPIOID ANALGESIC THERAPY

Unit 1



Balance Risks Against Potential Benefits

**Conduct thorough H&P
and appropriate testing**

**Comprehensive benefit-
to-harm evaluation**

Benefits Include

- Analgesia
(adequate pain control)
- Improved Function



Risks Include

- Overdose
- Life-threatening respiratory depression
- Abuse by patient or household contacts
- Misuse & addiction
- Physical dependence & tolerance
- Interactions w/ other medications & substances
- Risk of neonatal withdrawal syndrome w/ prolonged use during pregnancy
- Inadvertent exposure/ingestion by household contacts, especially children

Adequately **DOCUMENT**
all patient interactions,
assessments, test results,
& treatment plans

Clinical Interview: Patient Medical History

Illness relevant to (1) effects or (2) metabolism of opioids

1. Pulmonary disease, constipation, nausea, cognitive impairment
2. Hepatic, renal disease

Illness possibly linked to substance abuse, e.g.:

Hepatitis

HIV

Tuberculosis

Cellulitis

STIs

Trauma,
burns

Cardiac
disease

Pulmonary
disease

Clinical Interview: Pain & Treatment History

Description of pain



Location



Intensity



Quality



Onset/
Duration



Variations /
Patterns / Rhythms

What relieves the pain?

What causes or increases pain?

Effects of pain on physical, emotional, and psychosocial function

Patient's pain & functional goals

Clinical Interview: Pain & Treatment History, cont'd

Pain Medications



Past use

Current use

- Query state **PDMP** where available to confirm patient report
- Contact past providers & obtain prior medical records
- Conduct **UDT**

Dosage

- For opioids currently prescribed: opioid, dose, regimen, & duration
 - Important to determine if patient is **opioid tolerant**

General effectiveness

Nonpharmacologic strategies & effectiveness

Perform Thorough Evaluation & Assessment of Pain

Seek objective confirmatory data

Components of patient evaluation for pain

Order diagnostic tests (appropriate to complaint)

General: vital signs, appearance, posture, gait, & pain behaviors

Neurologic exam

Musculoskeletal Exam

- Inspection
- Palpation
- Percussion
- Auscultation
- Provocative maneuvers

Cutaneous or trophic findings

Assess Risk of Abuse, Including Substance Use & Psychiatric Hx

Obtain a complete Hx of current & past substance use

- Prescription drugs
- Illegal substances
- Alcohol & tobacco
 - Substance abuse Hx does not prohibit treatment w/ ER/LA opioids but may require additional monitoring & expert consultation/referral
- Family Hx of substance abuse & psychiatric disorders
- Hx of sexual abuse

Social history also relevant

Employment, cultural background, social network, marital history, legal history, & other behavioral patterns

Risk Assessment, cont'd

Be knowledgeable about risk factors for opioid abuse

- Personal or family Hx of alcohol or drug abuse
- Younger age
- Presence of psychiatric conditions

Understand & use addiction or abuse screening tools

- Assess potential risks associated w/ chronic opioid therapy
- Manage patients using ER/LA opioids based on risk assessment

Conduct a UDT

- Understand limitations

Risk Assessment Tools: Examples

Tool	# of items	Administered By
Patients considered for long-term opioid therapy:		
ORT Opioid Risk Tool	5	patient
SOAPP ® Screener & Opioid Assessment for Patients w/ Pain	24, 14, & 5	patient
DIRE Diagnosis, Intractability, Risk, & Efficacy Score	7	clinician
Characterize misuse once opioid treatments begins:		
PMQ Pain Medication Questionnaire	26	patient
COMM Current Opioid Misuse Measure	17	patient
PDUQ Prescription Drug Use Questionnaire	40	clinician
Not specific to pain populations:		
CAGE-AID Cut Down, Annoyed, Guilty, Eye-Opener Tool, Adjusted to Include Drugs	4	clinician
RAFFT Relax, Alone, Friends, Family, Trouble	5	patient
DAST Drug Abuse Screening Test	28	patient
SBIRT Screening, Brief Intervention, & Referral to Treatment	Varies	clinician

Opioid Risk Tool (ORT)

Mark each box that applies

Female

Male

1. Family Hx of substance abuse		Female	Male	
Alcohol	<input type="checkbox"/>	1	<input type="checkbox"/>	3
Illegal drugs	<input type="checkbox"/>	2	<input type="checkbox"/>	3
Prescription drugs	<input type="checkbox"/>	4	<input type="checkbox"/>	4
2. Personal Hx of substance abuse		Female	Male	
Alcohol	<input type="checkbox"/>	3	<input type="checkbox"/>	3
Illegal drugs	<input type="checkbox"/>	4	<input type="checkbox"/>	4
Prescription drugs	<input type="checkbox"/>	5	<input type="checkbox"/>	5
3. Age between 16 & 45 yrs		Female	Male	
	<input type="checkbox"/>	1	<input type="checkbox"/>	1
4. Hx of preadolescent sexual abuse		Female	Male	
	<input type="checkbox"/>	3	<input type="checkbox"/>	0
5. Psychologic disease		Female	Male	
ADD, OCD, bipolar, schizophrenia	<input type="checkbox"/>	2	<input type="checkbox"/>	2
Depression	<input type="checkbox"/>	1	<input type="checkbox"/>	1

Scoring Totals:

Administer

On initial visit

Prior to opioid therapy

Scoring (risk)

0-3: low

4-7: moderate

≥8: high

Screeners & Opioid Assessment for Patients with Pain (SOAPP)[®]

Identifies patients as at high, moderate, or low risk for misuse of opioids prescribed for chronic pain

How is SOAPP[®] administered?

Usually self-administered in waiting room, exam room, or prior to an office visit

May be completed as part of an interview w/ a nurse, physician, or psychologist

Prescribers should have a completed & scored SOAPP[®] while making opioid treatment decisions

When to Consider a Trial of an Opioid



Potential benefits are likely to outweigh risks

Failed to adequately respond to nonopioid & nondrug interventions


Continuous, around-the-clock opioid analgesic is needed for an extended period of time

Pain is chronic and severe

No alternative therapy is likely to pose as favorable a balance of benefits to harms

Chou R, et al. *J Pain*. 2009;10:113-30. Department of Veterans Affairs, Department of Defense. *VA/DoD Clinical Practice Guideline for Management of Opioid Therapy for Chronic Pain*. 2010.

When to Consider a Trial of an Opioid, cont'd



60-yr-old w/ chronic disabling OA pain

- Nonopioid therapies not effective, IR opioids provided some relief but experienced end-of-dose failure
- No psychiatric/medical comorbidity or personal/family drug abuse Hx
 - High potential benefits relative to potential risks
 - Could prescribe opioids to this patient in most settings w/ routine monitoring

30-yr-old w/ fibromyalgia & recent IV drug abuse

- High potential risks relative to benefits (opioid therapy not 1st line for fibromyalgia)
- Requires intensive structure, monitoring, & management by clinician w/ expertise in both addiction & pain
 - Not a good candidate for opioid therapy



When to Consider a Trial of an Opioid, cont'd

Selection of patients between these 2 extremes requires:

**Careful
assessment &
characterization
of patient risk**



**Structuring of care
to match risk**

In patients w/ Hx of substance abuse or a psychiatric comorbidity, this may require assistance from experts in managing pain, addiction, or other mental health concerns

In some cases opioids may not be appropriate or should be deferred until the comorbidity has been adequately addressed

– Consider referral

Referring High-Risk Patients

Prescribers should

Understand when to appropriately refer high-risk patients to pain management or addiction specialists

Also check your state regulations for requirements



Special Considerations: Elderly Patients



Does patient have medical problems that increase risk of opioid-related AEs?

Respiratory depression more likely in elderly, cachectic, or debilitated patients

- Altered PK due to poor fat stores, muscle wasting, or altered clearance
- Monitor closely, particularly when
 - Initiating & titrating ER/LA opioids
 - Given concomitantly w/ other drugs that depress respiration
- Reduce starting dose to 1/3 to 1/2 the usual dosage in debilitated, non-opioid-tolerant patients
- Titrate dose cautiously

Older adults more likely to develop constipation

- Routinely initiate a bowel regimen before it develops

Is patient/caregiver likely to manage opioid therapy responsibly?

Special Considerations: Pregnant Women



Managing chronic pain in pregnant women is challenging, & affects both mother and fetus

Potential risks of opioid therapy to the newborn include:

- Low birth weight
- Premature birth
- Hypoxic-ischemic brain injury
- Neonatal death
- Prolonged QT syndrome
- Neonatal opioid withdrawal syndrome

Given these potential risks, clinicians should:

- Counsel women of childbearing potential about risks & benefits of opioid therapy during pregnancy & after delivery
- Encourage minimal/no opioid use during pregnancy, unless potential benefits outweigh risks

If chronic opioid therapy is used during pregnancy, anticipate & manage risks to the patient and newborns



Special Considerations: Children (<18 years)

Safety & effectiveness of most ER/LA opioids unestablished

Pediatric analgesic trials pose challenges
Transdermal fentanyl approved in children aged ≥ 2 yrs

Most opioid studies focus on inpatient safety

Opioids are common sources of drug error

Opioid indications are primarily life-limiting conditions

Few children with chronic pain due to non-life-limiting conditions should receive opioids

When prescribing opioids to children:

Consult pediatric palliative care team or pediatric pain specialist or refer to a specialized multidisciplinary pain clinic

Berde CB, et al. *Pediatrics*. 2012;129:354-64. Gregoire MC, et al. *Pain Res Manag* 2013;18:47-50.
Mc Donnell C. *Pain Res Manag*. 2011;16:93-8. Slater ME, et al. *Pain Med*. 2010;11:207-14.

Challenge: The Friday Afternoon Patient

Red Flag:

Adjusting a prescription without performing appropriate evaluation or screening

It is 4 pm on Friday and you are four patients behind schedule. Mr. Kingston asks you to increase his current dosage of hydrocodone, because he says it is not relieving his pain. It would take you two minutes to say yes.

Action: Check your local PDMP. Employ practice management strategies that maximize efficiency.

- Patient-administered screening tools
- Office staff to administer and score tools, document results, and communicate to the prescriber

Challenge: The Delayed Surgery

Red Flag:

Patient may be stalling to continue an opioid regimen

Ms. Van Buskirk says she needs opioids to manage her pain until she can have surgery. She reports continued delays in getting to surgery. You phone the surgeon and discover that no date has been set and that she has cancelled several appointments.

Action: Set expectations for time limitations. Offer non-medicine and non-opioid options for pain management. Consider referral to addiction specialist.

Pearls for Practice



Document EVERYTHING

Conduct a Comprehensive H&P

General and pain-specific

Assess Risk of Abuse

Compare Risks with Expected Benefits

Determine Whether a Therapeutic Trial is Appropriate

INITIATING THERAPY, MODIFYING DOSING, & DISCONTINUING USE OF ER/LA OPIOID ANALGESICS

Unit II



Federal & State Regulations

Comply w/ federal & state laws & regulations that govern the use of opioid therapy for pain



Federal

- Code of Federal Regulations, Title 21 Section 1306: rules governing the issuance & filling of prescriptions pursuant to section 309 of the Act (21 USC 829)
 - www.deadiversion.usdoj.gov/21cfr/cfr/2106cfrt.htm
- United States Code (USC) - Controlled Substances Act, Title 21, Section 829: prescriptions
 - www.deadiversion.usdoj.gov/21cfr/21usc/829.htm



State

- Database of state statutes, regulations, & policies for pain management
 - www.medscape.com/resource/pain/opioid-policies
 - www.painpolicy.wisc.edu/database-statutes-regulations-other-policies-pain-management

Initiating Treatment

Prescribers should regard initial treatment as a therapeutic trial

May last from several weeks to several months

Decision to proceed w/ long-term treatment should be intentional & based on careful consideration of outcomes during the trial


Progress toward meeting therapeutic goals

Presence of opioid-related AEs

Changes in underlying pain condition

Changes in psychiatric or medical comorbidities

Identification of aberrant drug-related behavior, addiction, or diversion



ER/LA Opioid-Induced Respiratory Depression

Chief hazard of opioid agonists, including ER/LA opioids

- If not immediately recognized & treated, may lead to respiratory arrest & death
- Greatest risk: initiation of therapy or after dose increase

Manifested by reduced urge to breathe & decreased respiration rate

- Shallow breathing
- CO₂ retention can exacerbate opioid sedating effects

Instruct patients/family members to call 911*

- Managed w/ close observation, supportive measures, & opioid antagonists, depending on patient's clinical status

Chou R, et al. *J Pain*. 2009;10:113-30. FDA. *Blueprint for Prescriber Education for Extended-Release and Long-Acting Opioid Analgesics*. 08/2014.
www.fda.gov/downloads/Drugs/DrugSafety/PostmarketDrugSafety/InformationforPatientsandProviders/UCM311290.pdf

ER/LA Opioid-Induced Respiratory Depression

More likely to occur

- In elderly, cachectic, or debilitated patients
 - **Contraindicated** in patients w/ respiratory depression or conditions that increase risk
- If given concomitantly w/ other drugs that depress respiration

Reduce risk

- Proper dosing & titration are essential
- **Do not overestimate** dose when converting dosage from another opioid product
 - Can result in fatal overdose w/ first dose
- Instruct patients to swallow tablets/capsules whole
 - Dose from cut, crushed, dissolved, or chewed tablets/capsules may be fatal, particularly in opioid-naïve individuals

Initiating & Titrating: Opioid-Naïve Patients

Drug & dose selection is critical

Some ER/LA opioids or dosage forms are only recommended for **opioid-tolerant** patients

- ANY strength of transdermal fentanyl or hydromorphone ER
- Certain strengths/doses of other ER/LA products (check drug PI)

Monitor patients closely for respiratory depression

Especially within 24-72 h of initiating therapy & increasing dosage

Individualize dosage by titration based on efficacy, tolerability, & presence of AEs

Check ER/LA opioid product PI for minimum titration intervals

Supplement w/ IR analgesics (opioids & nonopioid) if pain is not controlled during titration

Initiating: Opioid-Tolerant Patients

***If opioid tolerant –
no restrictions on which products can be used***

Patients considered opioid tolerant are taking at least

- 60 mg oral morphine/day
- 25 mcg transdermal fentanyl/hr
- 30 mg oral oxycodone/day
- 8 mg oral hydromorphone/day
- 25 mg oral oxymorphone/day
- An equianalgesic dose of another opioid

Still requires caution when rotating a patient on an IR opioid to a different ER/LA opioid



**For 1 Wk
Or Longer**



Opioid Rotation



Definition:

Change from an existing opioid regimen to another opioid w/ the goal of improving therapeutic outcomes or to avoid AEs attributed to the existing drug, e.g., myoclonus

Rationale:

Differences in pharmacologic or other effects make it likely that a switch will improve outcomes

- Effectiveness & AEs of different mu opioids vary among patients
- Patients show incomplete cross-tolerance to new opioid
 - Patient tolerant to 1st opioid can have improved analgesia from 2nd opioid at a dose lower than calculated from an EDT

Equianalgesic Doses

Opioid rotation requires calculation of an approximate equianalgesic dose

Equianalgesic dose is a construct derived from relative opioid potency estimates

- Potency refers to dose required to produce a given effect

Relative potency estimates

- Ratio of doses necessary to obtain roughly equivalent effects
- Calculate across drugs or routes of administration
- Relative analgesic potency is converted into an equianalgesic dose by applying the dose ratio to a standard

Equianalgesic Dose Tables (EDT)

Many different versions:

Published

Online

Online Interactive

Smart-phone apps



Vary in terms of:



Equianalgesic values

Whether ranges are used

Which opioids are included:

May or may not include transdermal opioids, rapid-onset fentanyl, ER/LA opioids, or opioid agonist-antagonists

Example of an EDT for Adults



Drug	Equianalgesic Dose		Usual Starting Doses	
	SC/IV	PO	Parenteral	PO
Morphine	10 mg	30 mg	2.5-5 mg SC/IV q3-4hr (◆ 1.25 – 2.5mg)	5-15 mg q3-4hr (IR or oral solution) (◆ 2.5-7.5 mg)
Oxycodone	NA	20 mg	NA	5-10 mg q3-4 (◆ 2.5 mg)
Hydrocodone	NA	30 mg	NA	5 mg q3-4h (◆ 2.5 mg)
Hydromorphone	1.5 mg	7.5 mg	0.2-0.6 mg SC/IV q2-3hr (◆ 0.2mg)	1-2 mg q3-4hr (◆ 0.5-1 mg)

Limitations of EDTs

Single-dose potency studies using a specific route, conducted in patients w/ limited opioid exposure



Did Not Consider

Chronic dosing

High opioid doses

Other routes

Different pain types

Comorbidities or organ dysfunction

Gender, ethnicity, advanced age, or concomitant medications

Direction of switch from 1 opioid to another

Inter-patient variability in pharmacologic response to opioids

Incomplete cross-tolerance among mu opioids

Utilizing Equianalgesic Doses

Incomplete cross-tolerance & inter-patient variability require use of conservative dosing when converting from one opioid to another

Equianalgesic dose a starting point for opioid rotation

Intended as General Guide

Calculated dose of new drug based on EDT must be reduced, then titrate the new opioid as needed

Closely follow patients during periods of dose adjustments

Follow conversion instructions in individual ER/LA opioid PI, when provided

Guidelines for Opioid Rotation



Reduce calculated equianalgesic dose by 25%-50%*

Select % reduction based on clinical judgment

Closer to 50% reduction if patient is

- Receiving a relatively high dose of current opioid regimen
- Elderly or medically frail

Closer to 25% reduction if patient

- Does not have these characteristics
- Is switching to a different administration route of same drug

***75%-90% reduction for methadone**

Calculate equianalgesic dose of new opioid from EDT

Guidelines for Opioid Rotation, cont'd

If switching to **methadone**:



- Standard EDTs are less helpful in opioid rotation to methadone
- In opioid tolerant patients, methadone doses should not exceed 30-40 mg/day upon rotation.
 - Consider inpatient monitoring, including serial EKG monitoring
- In opioid-naïve patients, methadone should not be given as an initial drug

If switching to **transdermal**:

- **Fentanyl**, calculate dose conversion based on equianalgesic dose ratios included in the PI
- **Buprenorphine**, follow instructions in the PI

Guidelines for Opioid Rotation,

cont'd



Have a strategy to frequently assess analgesia, AEs and withdrawal symptoms

Titrate new opioid dose to optimize outcomes & safety

Dose for breakthrough pain (BTP) ***using a short-acting, immediate release preparation*** is 5%-15% of total daily opioid dose, administered at an appropriate interval

If oral transmucosal fentanyl product is used for BTP, begin dosing lowest dose irrespective of baseline opioid dose

NEVER use ER/LA opioids for BTP

Breakthrough Pain in Chronic Pain Patients

Patients on stable ATC opioids may experience BTP

Disease progression or a new or unrelated pain

Therapies

- Directed at cause of BTP or precipitating factors
- Nonspecific symptomatic therapies to lessen impact of BTP

Consider adding

- PRN IR opioid trial based on analysis of benefit versus risk
 - Risk for aberrant drug-related behaviors
 - High-risk: only in conjunction w/ frequent monitoring & follow-up
 - Low-risk: w/ routine follow-up & monitoring
- Nonopioid drug therapies
- Nonpharmacologic treatments

Reasons for Discontinuing ER/LA Opioids



No progress toward therapeutic goals

Intolerable & Unmanageable AEs

Pain level decreases in stable patients

Nonadherence or unsafe behavior

- 1 or 2 episodes of increasing dose without prescriber knowledge
- Sharing medications
- Unapproved opioid use to treat another symptom (e.g., insomnia)

Aberrant behaviors suggestive of addiction &/or diversion

- Use of illicit drugs or unprescribed opioids
- Repeatedly obtaining opioids from multiple outside sources
- Prescription forgery
- Multiple episodes of prescription loss

Challenge: The Broken Stereotype

Red Flag:

Making assumptions about a patient's risk factors without objective evidence

Ms. Yeun seems like a "good" patient. She has never abused opioids previously. She has been in the practice a long time, has never been a problem, and in fact, is rather enjoyable. She always brings Christmas cookies for the staff around the holidays.

Action: Require all patients receiving opioids to follow a treatment plan and adhere to defined expectations. Evaluate risk in all patients. Use patient-provider agreements, contracts, or other tools.

Challenge: The Early Refill

Optional Slide

Red Flag:

Patient requests an early refill every month.

You have prescribed Mr. Arias a long-acting opioid for low back pain and a short-acting PRN opioid for breakthrough pain. Every month he requests a refill for both prescriptions 3-8 days early. Upon questioning, Mr. Arias tells you that he takes both pills whenever he feels he needs them.

Action: Make sure that patients understand each medication's dosage, time of day, and maximum daily dose. Ask them to repeat these instructions back to you. Avoid clinical terms such as "PRN" that the patient may not understand.

Pearls for Practice



Treat Initiation of Opioids as a Therapeutic Trial

Anticipate ER/LA Opioid-Induced Respiratory Depression

It can be immediately life-threatening

Be Conservative and Thoughtful In Dosing


When initiating, titrating, and rotating opioids

First calculate equianalgesic dose, then reduce dose appropriately

Discontinue ER/LA opioids slowly and safely

MANAGING THERAPY WITH ER/LA OPIOID ANALGESICS

Unit III



Informed Consent

Before initiating a trial of opioid analgesic therapy, confirm patient understanding of informed consent to establish:

Analgesic & functional goals of treatment

Expectations

Potential risks

Alternatives to opioids

The potential for & how to manage:

- Common opioid-related AEs (e.g., constipation, nausea, sedation)
- Other serious risks (e.g., abuse, addiction, respiratory depression, overdose)
- AEs after long-term or high-dose opioid therapy (e.g., hyperalgesia, endocrinologic or sexual dysfunction)

Patient-Prescriber Agreement (PPA)

Document signed by both patient & prescriber at time an opioid is prescribed

Clarify treatment plan & goals of treatment w/ patient, patient's family, & other clinicians involved in patient's care

Assist in patient education

Inform patients about the risks & benefits

Document patient & prescriber responsibilities

Consider a PPA

Reinforce expectations for appropriate & safe opioid use

- Obtain opioids from a single prescriber
 - Fill opioid prescriptions at a designated pharmacy
 - Safeguard opioids
 - Do not store in medicine cabinet
 - Keep locked (e.g., use a medication safe)
 - Do not share or sell medication
 - Instructions for disposal when no longer needed
- Commitments to return for follow-up visits
 - Comply w/ appropriate monitoring
 - E.g., random UDT & pill counts
 - Frequency of prescriptions
 - Enumerate behaviors that may lead to opioid discontinuation
 - An exit strategy

Monitor Patients During Opioid Therapy



Therapeutic risks & benefits do not remain static

Affected by change in underlying pain condition, coexisting disease, or psychologic/ social circumstances

Identify patients

- Who are benefiting from opioid therapy
- Who might benefit more w/ restructuring of treatment or receiving additional services (e.g., addiction treatment)
- Whose benefits from treatment are outweighed by risks

Periodically assess continued need for opioid analgesic

Re-evaluate underlying medical condition if clinical presentation changes

Monitor Patients During Opioid Therapy, cont'd



Periodically evaluate:

- Pain control
 - Document pain intensity, pattern, & effects
- Functional outcomes
 - Document level of functioning
 - Assess progress toward achieving therapeutic goals
- Health-related QOL
- AE frequency & intensity
- Adherence to prescribed therapies

Patients requiring more frequent monitoring include:

- High-risk patients
- Patients taking high opioid doses

Anticipate & Treat Common AEs

Constipation

most common AE; does not resolve with time

- **Initiate a bowel regimen before constipation develops**
- **Increase fluid & fiber intake, stool softeners, & laxatives**
- **Opioid antagonists may help prevent/treat opioid-induced bowel dysfunction**

Nausea & vomiting

tend to diminish over days or weeks

Oral & rectal antiemetic therapies as needed

Drowsiness & sedation

tend to wane over time

Counsel patients about driving, work & home safety as well as risks of concomitant exposure to other drugs & substances w/ sedating effects

Pruritus & myoclonus

tend to diminish over days or weeks

Treatment strategies for either condition largely anecdotal

Monitor Adherence and Aberrant Behavior



Routinely monitor patient adherence to treatment plan

- Recognize & document aberrant drug-related behavior
 - In addition to patient self-report also use:
 - State PDMPs, where available
 - UDT
 - Positive for nonprescribed drugs
 - Positive for illicit substance
 - Negative for prescribed opioid
- Family member or caregiver interviews
- Monitoring tools such as the COMM, PADT, PMQ, or PDUQ
- Medication reconciliation (e.g., pill counts)

PADT=Pain Assessment & Documentation Tool

Address Aberrant Drug-Related Behavior

Behavior outside the boundaries of agreed-on treatment plan:

Behaviors that are **less** indicative of aberrancy

Unsanctioned dose escalations or other noncompliance w/ therapy on 1 or 2 occasions

Unapproved use of the drug to treat another symptom

Openly acquiring similar drugs from other medical sources

Behaviors that are **more** indicative of aberrancy

Multiple dose escalations or other noncompliance w/ therapy despite warnings

Prescription forgery

Obtaining prescription drugs from nonmedical sources

Prescription Drug Monitoring Programs (PDMPs)


48 states have an operational PDMP

1 state & DC have enacted PDMP legislation, not yet operational

1 state has no legislation

Individual state laws determine

- Who has access to PDMP information
- Which drug schedules are monitored
- Which agency administers the PDMP
- Whether prescribers are required to register w/ the PDMP
- Whether prescribers are required to access PDMP information in certain circumstances
- Whether unsolicited PDMP reports are sent to prescribers



PDMP Benefits



Record of a patient's controlled substance prescriptions

- Some are available online 24/7
- Opportunity to discuss w/ patient

Provide warnings of potential misuse/abuse

- Existing prescriptions not reported by patient
- Multiple prescribers/pharmacies
- Drugs that increase overdose risk when taken together
- Patient pays for drugs of abuse w/ cash



Prescribers can check their own prescribing Hx

PDMP Unsolicited Patient Threshold Reports

Reports automatically generated on patients who cross certain thresholds when filling prescriptions. Available in some states.

E-mailed to prescribers to whom prescriptions were attributed

Prescribers review records to confirm it is your patient & you wrote the prescription(s) attributed to you

If inaccurate, contact PDMP

If you wrote the prescription(s), patient safety may dictate need to discuss the patient w/ other prescribers listed on report

- Decide who will continue to prescribe for the patient & who might address drug abuse concerns.

Rationale for Urine Drug Testing (UDT)

Help to identify drug misuse/addiction

- Prior to starting opioid treatment

Assist in assessing adherence during opioid therapy

- As requirement of therapy w/ an opioid
- Support decision to refer

UDT frequency is based on clinical judgment

Depending on patient's display of aberrant behavior and whether it is sufficient to document adherence to treatment plan

Check state regulations for requirements



Main Types of UDT Methods

Initial testing w/ IA drug panels:



- Classify substance as present or absent according to cutoff
- Many do not identify individual drugs within a class
- Subject to cross-reactivity
- Either lab based or at POC

Identify specific drugs &/or metabolites w/ sophisticated lab-based testing; e.g., GC/MS or LC/MS*



- Specifically confirm the presence of a given drug
 - e.g., morphine is the opiate causing a positive IA*
- Identify drugs not included in IA tests
- When results are contested

* GC/MS=gas chromatography/ mass spectrometry
IA=immunoassay
LC/MS=liquid chromatography/ mass spectrometry

Detecting Opioids by UDT

Most common opiate IA drug panels

- Detect “opiates” morphine & codeine, but doesn’t distinguish
- Do not reliably detect semisynthetic opioids
 - Specific IA panels can be ordered for some
- Do not detect synthetic opioids (e.g., methadone, fentanyl)
 - Only a specifically directed IA panel will detect synthetics

GC/MS or LC/MS will identify specific opioids

- Confirm presence of a drug causing a positive IA
- Identify opioids not included in IA drug panels, including semisynthetic & synthetic opioids
- Identify opioids not included in IA drug panels, including semisynthetic & synthetic opioids

Interpretation of UDT Results

Positive Result



Demonstrates recent use

- Most drugs in urine have detection times of 1-3 d
- Chronic use of lipid-soluble drugs: test positive for ≥ 1 wk

Does not diagnose

- Drug addiction, physical dependence, or impairment

Does not provide enough information to determine

- Exposure time, dose, or frequency of use

Negative Result



Does not diagnose diversion

- More complex than presence or absence of a drug in urine

May be due to maladaptive drug-taking behavior

- Bingeing, running out early
- Other factors: eg, cessation of insurance, financial difficulties

Interpretation of UDT Results, cont'd



Be aware

Testing technologies & methodologies evolve

Differences exist between IA test menu panels vary


- Cross-reactivity patterns
 - Maintain list of all patient's prescribed & OTC drugs
 - Assist to identify false-positive result
- Cutoff levels

Time taken to eliminate drugs

- Document time of last use & quantity of drug(s) taken

Opioid metabolism may explain presence of apparently unprescribed drugs

Examples of Metabolism of Opioids



*6-MAM=6-monoacetylmorphine

Interpretation of UDT Results



Use UDT results in conjunction w/ other clinical information

Investigate unexpected results

Discuss w/ the lab

Schedule appointment
w/ patient to discuss
unexpected/abnormal results

Chart results, interpretation, & action

Do not ignore the *unexpected* positive result

May necessitate closer monitoring
&/or referral to a specialist



ER/LA Opioid Use in Pregnant Women



No adequate & well-controlled studies

Only use if potential benefit justifies the risk to the fetus

Be aware of the pregnancy status of your patients

- If prolonged use is required during pregnancy:
- Advise patient of risk of neonatal withdrawal syndrome
 - Ensure appropriate treatment will be available

Be Ready to Refer

Be familiar w/ referral sources for abuse or addiction that may arise from use of ER/LA opioids

SAMHSA substance abuse treatment facility locator

<http://findtreatment.samhsa.gov/TreatmentLocator/faces/quickSearch.jspx>

SAMHSA mental health treatment facility locator

<http://findtreatment.samhsa.gov/MHTreatmentLocator/faces/quickSearch.jspx>

Challenge: The Insistent Patient

Red Flag:

Patient refuses to consider non-opioid treatment options

Mr. Lee's daily function has improved significantly over the past two years. You suggest titrating his dosage down or trying alternative pain management options. He is extremely resistant and tells you "Nothing else relieves my pain."

Action: Work with your patient to set treatment goals and expectations. Select and document a therapy plan or use a patient-provider agreement. Evaluate Mr. Lee for potential addiction; consider referral to psychiatry or addiction medicine.

Pearls for Practice



Anticipate and Treat Common Adverse Effects

Use Informed Consent and Patient Provider Agreements

Use UDT and PDMP as Valuable Sources of Data About your Patient

However, know their limitations

Monitor Patient Adherence, Side Effects, Aberrant Behaviors, and Clinical Outcomes

Refer Appropriately if Necessary

COUNSELING PATIENTS & CAREGIVERS ABOUT THE SAFE USE OF ER/LA OPIOID ANALGESICS

Unit IV



Counsel Patients About Proper Use



Explain

- Product-specific information about the prescribed ER/LA opioid
- How to take the ER/LA opioid as prescribed
- Importance of adherence to dosing regimen, handling missed doses, & contacting their prescriber if pain cannot be controlled

Instruct patients/ caregivers to

- Read the ER/LA opioid **Medication Guide** received from pharmacy **every time** an ER/LA opioid is dispensed
- At every medical appointment explain all medications they take

Counsel Patients About Proper Use, cont'd

Counsel patients/caregivers:

- On the most common AEs of ER/LA opioids
- About the risk of falls, working w/ heavy machinery, & driving
- Call the prescriber for advice about managing AEs
- Inform the prescriber about AEs



Prescribers should report serious AEs to the FDA:
www.fda.gov/downloads/AboutFDA/ReportsManualsForms/Forms/UCM163919.pdf
or **1-800-FDA-1088**

Warn Patients



Never break, chew, crush or snort an oral ER/LA tablet/capsule, or cut or tear patches prior to use



- May lead to rapid release of ER/LA opioid causing overdose & death
- When a patient cannot swallow a capsule whole, prescribers should refer to PI to determine if appropriate to sprinkle contents on applesauce or administer via feeding tube



Use of CNS depressants or alcohol w/ ER/LA opioids can cause overdose & death



- Use with alcohol may result in rapid release & absorption of a potentially fatal opioid dose
- Other depressants include sedative-hypnotics & anxiolytics, illegal drugs



Warn Patients, cont'd

Misuse of ER/LA opioids can lead to death

- Take **exactly** as directed*
- Counsel patients/caregivers on risk factors, signs, & symptoms of overdose & opioid-induced respiratory depression, GI obstruction, & allergic reactions
- Call **911** or poison control
1-800-222-1222

***Serious side effects, including death, can occur even when used as recommended**

Do not abruptly stop or reduce the ER/LA opioid use

- Discuss how to safely taper the dose when discontinuing

Be safe. Be sure. Read the label.

Check your name.

ABC Pharmacy SMITH, JOHN
123 MAIN STREET
SPRINGFIELD, US 01234

Rx. 587123

TAKE 1 TABLET BY MOUTH
EVERY 12 HOURS

OXYCONTIN 10 MG

Qty: 60 TABLETS

Date Filled: 01/12 Discard After: 07/12

CAUTION: FEDERAL LAW PROHIBITS THE TRANSFER OF THIS DRUG TO ANY PERSON OTHER THAN THE PATIENT FOR WHOM IT WAS PRESCRIBED.

Check any warnings.

Check the directions.

TAKE OR USE THIS EXACTLY AS DIRECTED. DO NOT SKIP DOSES OR DISCONTINUE.

POISON Help
1-800-222-1222

Did you take the wrong medicine? Did you take too much? Call your Poison Center. Expert advice is available 24/7.

Consider Prescribing Naloxone

Naloxone:

- An opioid antagonist
- Antidote to acute opioid toxicity
- Instruct patients to use in event of known or suspected overdose, **in addition to calling emergency services**

Candidates for naloxone include those:

- Taking high-doses of opioids
- Taking opioid preparations that may increase risk for overdose; eg, ER/LA opioids
- Undergoing opioid rotation
- Discharged from emergency medical care following opioid intoxication/poisoning
- Legitimate medical need for analgesia, coupled with suspected/confirmed substance abuse

Available as:

- Naloxone kit (w/ syringes & needles)
- EVZIO™ (naloxone HCl) auto-injector

Encourage patients to:

- Create an “overdose plan”
- Involve friends, family members, partners, &/or caregivers

Protecting the Community



Caution Patients

- **Sharing ER/LA opioids w/ others may cause them to have serious AEs**
 - Including death
- **Selling or giving away ER/LA opioids is against the law**
- **Store medication safely and securely**
- **Protect ER/LA opioids from theft**
- **Dispose of any ER/LA opioids when no longer needed**
 - Read product-specific disposal information included w/ ER/LA opioid

Know Your Poison Center's Number.




POISON
Help
1-800-222-1222

**You could save
a life.**

1-800-222-1222

Source of Most Recent Rx Opioids Among Past-Year Users (2011-2012)



Educate Parents: Not in My House

Step 1: Monitor

- Note how many pills in each prescription bottle or pill packet
- Keep track of refills for all household members
- If your teen has been prescribed a drug, coordinate & monitor dosages & refills
- Make sure friends & relatives—especially grandparents—are aware of the risks
- If your teen visits other households, talk to the families about safeguarding their medications

Rx Opioid Disposal



New "Disposal Act" expands ways for patients to dispose of unwanted/expired opioids

Decreases amount of opioids introduced into the environment, particularly into water

Collection receptacles

Call DEA Registration Call Center at **1-800-882-9539** to find a local collection receptacle



Mail-back packages

Obtained from authorized collectors




Local take-back events

- Conducted by Federal, State, tribal, or local law enforcement
- Partnering w/ community groups

Voluntarily maintained by:

- Law enforcement
- Authorized collectors, including:
 - Manufacturer
 - Distributer
 - Reverse distributor
 - Retail or hospital/clinic pharmacy
 - Including long-term care facilities

Last DEA National Prescription Drug Take-Back Day on September 27, 2014



Other Methods of Opioid Disposal

If collection receptacle, mail-back program, or take-back event unavailable, throw out in household trash

- Take drugs out of original containers
- Mix w/ undesirable substance, e.g., used coffee grounds or kitty litter
 - Less appealing to children/pets, & unrecognizable to people who intentionally go through your trash
- Place in sealable bag, can, or other container
 - Prevent leaking or breaking out of garbage bag
- Before throwing out a medicine container
 - Scratch out identifying info on label



Prescription Drug Disposal

**FDA lists especially harmful medicines –
in some cases fatal w/ just 1 dose –
if taken by someone other than the patient**

- Instruct patients to check medication guide



Flush down sink/toilet if no collection receptacle, mail-back program, or take-back event available

- **As soon as they are no longer needed**
 - So cannot be accidentally taken by children, pets, or others
- **Includes transdermal adhesive skin patches**
 - Used patch worn for 3d still contains enough opioid to harm/kill a child
 - Dispose of used patches immediately after removing from skin
- **Fold patch in half so sticky sides meet, then flush down toilet**
- **Do NOT place used or unneeded patches in household trash**
 - Exception is Butrans: can seal in Patch-Disposal Unit provided & dispose of in the trash

Challenge: The Offended Patient

Red Flag:

You decide not to request routine risk assessment for fear of creating conflict

Mrs. Jorgensen has been your patient for eight years and has never caused any problems. When you ask her to undergo urine drug testing, she becomes upset and accuses you of not trusting her.

Action: Describe UDT as a routine part of medication monitoring rather than a “drug test”. Create an office policy for performing UDT on all ER/LA opioid patients. Practice by following universal precautions. Use a patient-provider agreement to clarify expectations of treatment.

Challenge: The Daughter's Party

Red Flag:

Patients do not safeguard their opioid medications correctly

Your patient's daughter, Jody, stole her father's opioids from his bedside drawer to take to a "fishbowl party". Her best friend consumed a mix of opioids and alcohol and died of an overdose.

Action: Always counsel patients about safe drug storage; warn patients about the serious consequences of theft, misuse, and overdose. Tell your patients that taking another person's medication, even once, is against the law.

Pearls for Practice



Establish Informed Consent

Counsel Patients about Proper Use

Appropriate use of medication
Consequences of inappropriate use

Educate the Whole Team


Patients, families, caregivers

Tools and Documents Can Help with Counseling

Use them!

GENERAL DRUG INFORMATION FOR ER/LA OPIOID ANALGESIC PRODUCTS

Unit V



General ER/LA Opioid Drug Information

Prescribers should be knowledgeable about general characteristics, toxicities, & drug interactions for ER/LA opioid products:

ER/LA opioid analgesic products are scheduled under the Controlled Substances Act & can be misused & abused

Respiratory depression is the most serious opioid AE

Can be immediately life-threatening

Constipation is the most common long-term AE

Should be anticipated

For Safer Use: Know Drug Interactions, PK, & PD



CNS depressants can potentiate sedation & respiratory depression

Some ER/LA products rapidly release opioid (dose dump) when exposed to alcohol

Some drug levels may increase without dose dumping

Use w/ MAOIs may increase respiratory depression

Certain opioids w/ MAOIs can cause serotonin syndrome

Can reduce efficacy of diuretics

Inducing release of antidiuretic hormone

Methadone & buprenorphine can prolong QTc interval

Drugs that inhibit or induce CYP enzymes can increase or lower blood levels of some opioids

Opioid Tolerant

Tolerance to sedating & respiratory-depressant effects is critical to safe use of certain ER/LA opioid products, dosage unit strengths, or doses

Patients must be opioid tolerant before using

- Any strength of transdermal fentanyl or hydromorphone ER
- Certain strengths or daily doses of other ER products

Opioid-tolerant patients are those taking at least

- 60 mg oral morphine/day
- 25 mcg transdermal fentanyl/hr
- 30 mg oral oxycodone/day
- 8 mg oral hydromorphone/day
- 25 mg oral oxymorphone/day
- An equianalgesic dose of another opioid

**FOR 1 WK
OR LONGER**

Key Instructions: ER/LA Opioids



Individually titrate to a dose that provides adequate analgesia & minimizes adverse reactions

Times required to reach steady-state plasma concentrations are product-specific

Refer to product information for titration interval

Continually re-evaluate to assess maintenance of pain control & emergence of AEs

Key Instructions: ER/LA Opioids,



cont'd

During chronic therapy, especially for non-cancer-related pain, periodically reassess the continued need for opioids

If pain increases, attempt to identify source, while adjusting dose

When an ER/LA opioid is no longer required, gradually titrate dose downward to prevent signs & symptoms of withdrawal in physically dependent patients

Do not abruptly discontinue

Common Drug Information for This Class

Limitations of usage

- Reserve for when alternative options (eg, non-opioids or IR opioids) are ineffective, not tolerated, or otherwise inadequate
- Not for use as an as-needed analgesic
- Not for mild pain or pain not expected to persist for an extended duration
- Not for acute pain

Dosage reduction for hepatic or renal impairment

See individual drug PI

Relative potency to oral morphine

- Intended as general guide
- Follow conversion instructions in individual PI
- Incomplete cross-tolerance & inter-patient variability require conservative dosing when converting from 1 opioid to another
 - Halve calculated comparable dose & titrate new opioid as needed

Transdermal Dosage Forms

Do not cut, damage, chew, or swallow



Exertion or exposure to external heat can lead to fatal overdose

Rotate location of application

Prepare skin: clip - not shave - hair & wash area w/ water

Monitor patients w/ fever for signs or symptoms of increased opioid exposure

Metal foil backings are not safe for use in MRIs

Drug Interactions Common to this Class

Concurrent use w/ other CNS depressants can increase risk of respiratory depression, hypotension, profound sedation, or coma

Reduce initial dose of one or both agents

Avoid concurrent use of partial agonists* or mixed agonist/antagonists† with full opioid agonist
May reduce analgesic effect &/or precipitate withdrawal

May enhance neuromuscular blocking action of skeletal muscle relaxants & increase respiratory depression

Concurrent use w/ anticholinergic medication increases risk of urinary retention & severe constipation
May lead to paralytic ileus

*Buprenorphine; †Pentazocine, nalbuphine, butorphanol

Drug Information Common to This Class

Use in opioid-tolerant patients

- See individual PI for products which:
 - Have strengths or total daily doses only for use in opioid-tolerant patients
 - Are only for use in opioid-tolerant patients at all strengths

Contraindications

- Significant respiratory depression
- Acute or severe asthma in an unmonitored setting or in absence of resuscitative equipment
- Known or suspected paralytic ileus
- Hypersensitivity (e.g., anaphylaxis)
- See individual PI for additional contraindications

Pearls for Practice



Patients **MUST** be opioid-tolerant in order to safely take most ER/LA opioid products

Be familiar with drug-drug interactions, pharmacokinetics and pharmacodynamics of ER/LA opioids

Central nervous system depressants (alcohol, sedatives, hypnotics, tranquilizers, tricyclic antidepressants) can have a potentiating effect on the sedation and respiratory depression caused by opioids.

Challenge: The Patient in the ER


Red Flag:

You are woken by a telephone call at 2 am reporting that your patient, Mr. Diallo, is in the ER with apparent respiratory depression.

Action: Be familiar with risk factors for respiratory depression and know when opioids are contra-indicated. Anticipate possible risks and develop contingency plans. Teach patients, family, and caregivers about respiratory depression and its symptoms.

SPECIFIC DRUG INFORMATION FOR ER/LA OPIOID ANALGESIC PRODUCTS

Unit VI



Specific Characteristics

Know for opioid products you prescribe:

**Drug
substance**

Formulation

Strength

**Dosing
interval**

**Key
instructions**

**Use in opioid-
tolerant
patients**

**Product-
specific safety
concerns**

**Relative
potency to
morphine**

**Specific information about
product conversions, if available**

Specific drug interactions

For detailed information, refer to online PI:

DailyMed at www.dailymed.nlm.nih.gov Drugs@FDA at www.fda.gov/drugsatfda

Morphine Sulfate ER Capsules (Avinza)

Dosing interval	<ul style="list-style-type: none">• Once a day
Key instructions	<ul style="list-style-type: none">• Initial dose in opioid non-tolerant patients is 30 mg• Titrate in increments of not greater than 30 mg using a minimum of 3-4 d intervals• Swallow capsule whole (do not chew, crush, or dissolve)• May open capsule & sprinkle pellets on applesauce for patients who can reliably swallow without chewing; use immediately• MDD:* 1600 mg (renal toxicity of excipient, fumaric acid)
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications w/ alcohol may result in rapid release & absorption of potentially fatal dose• P-gp* inhibitors (e.g., quinidine) may increase absorption/exposure of morphine by ~2-fold
Opioid-tolerant	<ul style="list-style-type: none">• 90 mg & 120 mg capsules for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• None

* MDD=maximum daily dose; P-gp= P-glycoprotein

Buprenorphine Transdermal System (Butrans)

Dosing interval

- One transdermal system every 7 d

Key instructions

- Initial dose in opioid non-tolerant patients on <30 mg morphine equivalents & in mild-moderate hepatic impairment: 5 mcg/h
- When converting from 30 mg-80 mg morphine equivalents, first taper to 30 mg morphine equivalent, then initiate w/ 10 mcg/h
- Titrate in 5 or 10 mcg/h increments by using no more than 2 patches of the 5 or 10 mcg/h system(s) w/ minimum of 72 h prior between dose adjustments. Total dose from all patches should be ≤ 20 mcg/h
- Maximum dose: 20 mcg/h due to risk of QTc prolongation
- Application
 - Apply only to sites indicated in PI
 - Apply to intact/non-irritated skin
 - Prep skin by clipping hair; wash site w/ water only
 - Rotate application site (min 3 wks before reapply to same site)
 - Do not cut
- Avoid exposure to heat
- Dispose of patches: fold adhesive side together & flush down toilet

Buprenorphine Transdermal System (Butrans) cont'd

Drug interactions

- CYP3A4 inhibitors may increase buprenorphine levels
- CYP3A4 inducers may decrease buprenorphine levels
- Benzodiazepines may increase respiratory depression
- Class IA & III antiarrhythmics, other potentially arrhythmogenic agents, may increase risk of QTc prolongation & torsade de pointe

Opioid-tolerant

- 7.5 mcg/h, 10 mcg/h, 15 mcg/h, & 20 mcg/h for use in opioid-tolerant patients only

Drug-specific safety concerns

- QTc prolongation & torsade de pointe
- Hepatotoxicity
- Application site skin reactions

Relative potency: oral morphine

- Equipotency to oral morphine not established

Methadone Hydrochloride Tablets (Dolophine)

NOTE: While the dosing information below reflects the 8/20/14 FDA Blue Print, the CO*RE Expert Clinical Faculty believe it to be too aggressive and perhaps a risky approach. CO*RE Expert Clinical Faculty discourages methadone for opioid naive patients as an initial drug and recommends 4-5 d intervals for dosing adjustments.

Dosing interval

- Every 8 to 12 h

Key instructions

- Initial dose in opioid non-tolerant patients: 2.5 – 10 mg
- Conversion of opioid-tolerant patients using equianalgesic tables can result in overdose & death. Use low doses according to table in full PI
- Dosage adjustments using a minimum of 1-2 d intervals
- High inter-patient variability in absorption, metabolism, & relative analgesic potency
- Opioid detoxification or maintenance treatment only provided in a federally certified opioid (addiction) treatment program (CFR, Title 42, Sec 8)

Drug interactions

- Pharmacokinetic drug-drug interactions w/ methadone are complex
 - CYP 450 inducers may decrease methadone levels
 - CYP 450 inhibitors may increase methadone levels
 - Anti-retroviral agents have mixed effects on methadone levels
- Potentially arrhythmogenic agents may increase risk for QTc prolongation & torsade de pointe
- Benzodiazepines may increase respiratory depression

Methadone Hydrochloride Tablets (Dolophine) cont'd

Opioid-tolerant

- Refer to full PI

Drug-specific safety concerns

- QTc prolongation & torsade de pointe
- Peak respiratory depression occurs later & persists longer than analgesic effect
- Clearance may increase during pregnancy
- False-positive UDT possible

Relative potency: oral morphine

- Varies depending on patient's prior opioid experience

Fentanyl Transdermal System

(Duragesic)

12, 25, 37.5*, 50, 62.5*, 75, 87.5*, and 100 mcg/hr
(*These strengths are available only in generic form)

Dosing interval

- Every 72 h (3 d)

Key instructions

- Use product-specific information for dose conversion from prior opioid
- Hepatic or renal impairment: use 50% of dose if mild/moderate, avoid use if severe
- Application
 - Apply to intact/non-irritated/non-irradiated skin on a flat surface
 - Prep skin by clipping hair, washing site w/ water only
 - Rotate site of application
 - Titrate using a minimum of 72 h intervals between dose adjustments
 - Do not cut
- Avoid exposure to heat
- Avoid accidental contact when holding or caring for children
- Dispose of used/unused patches: fold adhesive side together & flush down toilet

Fentanyl Transdermal System (Duragesic), cont'd

Key instructions	Specific contraindications: <ul style="list-style-type: none">• Patients who are not opioid-tolerant• Management of<ul style="list-style-type: none">– Acute or intermittent pain, or patients who require opioid analgesia for a short time– Post-operative pain, out-patient, or day surgery– Mild pain
Drug interactions	<ul style="list-style-type: none">• CYP3A4 inhibitors may increase fentanyl exposure• CYP3A4 inducers may decrease fentanyl exposure• Discontinuation of concomitant CYP P450 3A4 inducer may increase fentanyl plasma concentration
Opioid-tolerant	<ul style="list-style-type: none">• All doses indicated for opioid-tolerant patients only
Drug-specific safety concerns	<ul style="list-style-type: none">• Accidental exposure due to secondary exposure to unwashed/unclothed application site• Increased drug exposure w/ increased core body temp or fever• Bradycardia• Application site skin reactions
Relative potency: oral morphine	<ul style="list-style-type: none">• See individual PI for conversion recommendations from prior opioid

Morphine Sulfate ER-Naltrexone Tablets (Embeda)

Dosing interval	<ul style="list-style-type: none">• Once a day or every 12 h
Key instructions	<ul style="list-style-type: none">• Initial dose as first opioid: 20 mg/0.8 mg• Titrate using a minimum of 1-2 d intervals• Swallow capsules whole (do not chew, crush, or dissolve)• Crushing or chewing will release morphine, possibly resulting in fatal overdose, & naltrexone, possibly resulting in withdrawal symptoms• May open capsule & sprinkle pellets on applesauce for patients who can reliably swallow without chewing, use immediately
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications w/ alcohol may result in rapid release & absorption of potentially fatal dose• P-gp inhibitors (e.g., quinidine) may increase absorption/exposure of morphine by ~2-fold
Opioid-tolerant	<ul style="list-style-type: none">• 100 mg/4 mg capsule for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• None

Hydromorphone Hydrochloride ER Tablets (Exalgo)

Dosing interval	<ul style="list-style-type: none">• Once a day
Key instructions	<ul style="list-style-type: none">• Use conversion ratios in individual PI• Start patients w/ moderate hepatic impairment on 25% dose prescribed for patient w/ normal function• Renal impairment: start patients w/ moderate on 50% & patients w/ severe on 25% dose prescribed for patient w/ normal function• Titrate in increments of 4-8 mg using a minimum of 3-4 d intervals• Swallow tablets whole (do not chew, crush, or dissolve)• Do not use in patients w/ sulfite allergy (contains sodium metabisulfite)
Drug interactions	<ul style="list-style-type: none">• None
Opioid-tolerant	<ul style="list-style-type: none">• All doses are indicated for opioid-tolerant patients only
Product-specific adverse reactions	<ul style="list-style-type: none">• Allergic manifestations to sulfite component
Relative potency: oral morphine	<ul style="list-style-type: none">• ~5:1 oral morphine to hydromorphone oral dose ratio, use conversion recommendations in individual product information

Hydrocodone Bitartrate (Hysingla ER)

Extended-Release Tablets, 20 mg, 30 mg, 40 mg, 60 mg, 80 mg, 100 mg, and 120mg

Dosing interval

- Once a day

Key instructions

- Opioid-naïve patients: initiate treatment with 20 mg orally once daily.
- During titration, adjust the dose in increments of 10 mg to 20 mg every 3 to 5 days until adequate analgesia is achieved.
- Swallow tablets whole (do not chew, crush, or dissolve).
- Consider use of an alternative analgesic in patients who have difficulty swallowing or have underlying gastrointestinal disorders that may predispose them to obstruction.
- Take one tablet at a time, with enough water to ensure complete swallowing immediately after placing in the mouth.
- Use 1/2 of the initial dose and monitor closely for adverse events, such as respiratory depression and sedation, when administering Hysingla ER to patients with severe hepatic impairment or patients with moderate to severe renal impairment.

Hydrocodone Bitartrate (Hysingla ER), cont'd

Drug interactions	<ul style="list-style-type: none">• CYP3A4 inhibitors may increase hydrocodone exposure.• CYP3A4 inducers may decrease hydrocodone exposure.• Concomitant use of Hysingla ER with strong laxatives (e.g., Lactulose) that rapidly increase GI motility may decrease hydrocodone absorption and result in decreased hydrocodone plasma levels.• The use of MAO inhibitors or tricyclic antidepressants with Hysingla ER may increase the effect of either the antidepressant or Hysingla ER.
Opioid-tolerant	<ul style="list-style-type: none">• 80 mg is only for use in opioid tolerant patients.
Drug-specific safety concerns	<ul style="list-style-type: none">• Use with caution in patients with difficulty swallowing the tablet or underlying gastrointestinal disorders that may predispose patients to obstruction.• Esophageal obstruction, dysphagia, and choking have been reported with Hysingla ER.• In nursing mothers, discontinue nursing or discontinue drug. QTc prolongation has been observed with Hysingla ER following daily doses of 160 mg.• Avoid use in patients with congenital long QTc syndrome. This observation should be considered in making clinical decisions regarding patient monitoring when prescribing Hysingla ER in patients with congestive heart failure, bradyarrhythmias, electrolyte abnormalities, or who are taking medications that are known to prolong the QTc interval.• In patients who develop QTc prolongation, consider reducing the dose.
Relative potency: oral morphine	<ul style="list-style-type: none">• See individual PI for conversion recommendations from prior opioid



Morphine Sulfate ER Capsules (Kadian)

Dosing interval	<ul style="list-style-type: none">• Once a day or every 12 h
Key instructions	<ul style="list-style-type: none">• PI recommends not using as first opioid• Titrate using minimum of 2-d intervals• Swallow capsules whole (do not chew, crush, or dissolve)• May open capsule & sprinkle pellets on applesauce for patients who can reliably swallow without chewing, use immediately
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications w/ alcohol may result in rapid release & absorption of potentially fatal dose of morphine• P-gp inhibitors (e.g., quinidine) may increase absorption/exposure of morphine by ~2-fold
Opioid-tolerant	<ul style="list-style-type: none">• 100 mg & 200 mg capsules for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• None

Morphine Sulfate CR Tablets (MS Contin)

Dosing interval	<ul style="list-style-type: none">• Every 8 h or every 12 h
Key instructions	<ul style="list-style-type: none">• Product information recommends not using as first opioid.• Titrate using a minimum of 1-2 d intervals• Swallow tablets whole (do not chew, crush, or dissolve)
Drug interactions	<ul style="list-style-type: none">• P-gp inhibitors (e.g., quinidine) may increase absorption/exposure of morphine by ~2-fold
Opioid-tolerant	<ul style="list-style-type: none">• 100 mg & 200 mg tablet strengths for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• None

Tapentadol ER Tablets (Nucynta ER)

Dosing interval	<ul style="list-style-type: none">• Every 12 h
Key instructions	<ul style="list-style-type: none">• 50 mg every 12 h is initial dose in opioid non-tolerant patients• Titrate by 50 mg increments using minimum of 3-d intervals• MDD: 500 mg• Swallow tablets whole (do not chew, crush, or dissolve)• Take 1 tablet at a time w/ enough water to ensure complete swallowing immediately after placing in mouth• Dose once/d in moderate hepatic impairment (100 mg/d max)• Avoid use in severe hepatic & renal impairment
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications w/ alcohol may result in rapid release & absorption of a potentially fatal dose of tapentadol• Contraindicated in patients taking MAOIs
Opioid-tolerant	<ul style="list-style-type: none">• No product-specific considerations
Product-specific safety concerns	<ul style="list-style-type: none">• Risk of serotonin syndrome• Angio-edema
Relative potency: oral morphine	<ul style="list-style-type: none">• Equipotency to oral morphine has not been established

Oxymorphone Hydrochloride ER Tablets (Opana ER)

Dosing interval	<ul style="list-style-type: none">• Every 12 h dosing, some may benefit from asymmetric (different dose given in AM than in PM) dosing
Key instructions	<ul style="list-style-type: none">• Use 5 mg every 12 h as initial dose in opioid non-tolerant patients & patients w/ mild hepatic impairment & renal impairment (creatinine clearance <50 mL/min) & patients >65 yrs• Swallow tablets whole (do not chew, crush, or dissolve)• Take 1 tablet at a time, w/ enough water to ensure complete swallowing immediately after placing in mouth• Titrate in increments of 5-10 mg using a minimum of 3-7 d intervals• Contraindicated in moderate & severe hepatic impairment
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications w/ alcohol may result in absorption of a potentially fatal dose of oxymorphone
Opioid-tolerant	<ul style="list-style-type: none">• No product-specific considerations
Product-specific safety concerns	<ul style="list-style-type: none">• Use with caution in patients who have difficulty swallowing or underlying GI disorders that may predispose to obstruction (e.g. small gastrointestinal lumen)
Relative potency: oral morphine	<ul style="list-style-type: none">• Approximately 3:1 oral morphine to oxymorphone oral dose ratio

Oxycodone Hydrochloride CR Tablets (OxyContin)

Dosing interval	<ul style="list-style-type: none">• Every 12 h
Key instructions	<ul style="list-style-type: none">• Initial dose in opioid non-tolerant patients: / 10 mg every 12 h• Titrate using a minimum of 1-2 d intervals• Hepatic impairment: start w/ $\frac{1}{3}$-$\frac{1}{2}$ usual dosage• Renal impairment (creatinine clearance <60 mL/min): start w/ $\frac{1}{2}$ usual dosage• Consider other analgesics in patients w/ difficulty swallowing or underlying GI disorders that predispose to obstruction. Swallow tablets whole (do not chew, crush, or dissolve)• Take 1 tablet at a time, w/ enough water to ensure complete swallowing immediately after placing in mouth
Drug interactions	<ul style="list-style-type: none">• CYP3A4 inhibitors may increase oxycodone exposure• CYP3A4 inducers may decrease oxycodone exposure
Opioid-tolerant	<ul style="list-style-type: none">• Single dose >40 mg or total daily dose >80 mg for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• Choking, gagging, regurgitation, tablets stuck in throat, difficulty swallowing tablet• Contraindicated in patients w/ GI obstruction
Relative potency: oral morphine	<ul style="list-style-type: none">• Approximately 2:1 oral morphine to oxycodone oral dose ratio

Oxycodone Hydrochloride/Naloxone Hydrochloride ER Tablets (Targiniq ER)

Dosing interval	<ul style="list-style-type: none">• Every 12 h
Key instructions	<ul style="list-style-type: none">• Opioid-naïve patients: initiate treatment w/ 10mg/5mg every 12 h• Titrate using min of 1-2 d intervals• Do not exceed 80 mg/40 mg total daily dose (40 mg/20 mg q12h)• May be taken w/ or without food• Swallow whole. Do not chew, crush, split, or dissolve: this will release oxycodone (possible fatal overdose) & naloxone (possible withdrawal)• Hepatic impairment: contraindicated in moderate-severe impairment. In patients w/ mild impairment, start w/ $\frac{1}{3}$-$\frac{1}{2}$ usual dosage• Renal impairment (creatinine clearance <60 mL/min): start w/ $\frac{1}{2}$ usual dosage
Drug interactions	<ul style="list-style-type: none">• CYP3A4 inhibitors may increase oxycodone exposure• CYP3A4 inducers may decrease oxycodone exposure
Opioid-tolerant	<ul style="list-style-type: none">• Single dose >40 mg/20 mg or total daily dose of 80 mg/40 mg for opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• Contraindicated in patients w/ moderate-severe hepatic impairment
Relative potency: oral morphine	<ul style="list-style-type: none">• See individual PI for conversion recommendations from prior opioids

Hydrocodone Bitartrate ER Capsules (Zohydro ER)

Dosing interval	<ul style="list-style-type: none">• Every 12 h
Key instructions	<ul style="list-style-type: none">• Initial dose in opioid non-tolerant patient is 10 mg• Titrate in increments of 10 mg using a min of 3-7 d intervals• Swallow capsules whole (do not chew, crush, or dissolve)
Drug interactions	<ul style="list-style-type: none">• Alcoholic beverages or medications containing alcohol may result in rapid release & absorption of a potentially fatal dose of hydrocodone• CYP3A4 inhibitors may increase hydrocodone exposure• CYP3A4 inducers may decrease hydrocodone exposure
Opioid-tolerant	<ul style="list-style-type: none">• Single dose >40 mg or total daily dose >80 mg for use in opioid-tolerant patients only
Product-specific safety concerns	<ul style="list-style-type: none">• None
Relative potency: oral morphine	<ul style="list-style-type: none">• Approximately 1.5:1 oral morphine to hydrocodone oral dose ratio

Summary



Prescription opioid abuse & overdose is a national epidemic. Clinicians must play a role in prevention

Understand how to assess patients for treatment w/ ER/LA opioids

Be familiar w/ how to initiate therapy, modify dose, & discontinue use of ER/LA opioids

Know how to manage ongoing therapy w/ ER/LA opioids

Know how to counsel patients & caregivers about the safe use of ER/LA opioids, including proper storage & disposal

Be familiar w/ general & product-specific drug information concerning ER/LA opioids

IMPORTANT!

Thank you for completing the post-activity assessment for this CO*RE session.

Your participation in this assessment allows CO*RE to report de-identified numbers to the FDA.

A strong show of engagement will demonstrate that clinicians have voluntarily taken this important education and are committed to patient safety and improved outcomes.

THANK YOU!

Thank you!

www.core-remms.org

