Main Office:

Scottsdale Vein & Proctology Center 8752 E Via De Commercio #2 Scottsdale, Arizona 85258

Office: (602) 492-9919 | Mobile: (602) 920-1023



Name:		DOB:	Date:	
	DISPENSE ONE OF THE FOLLOWING FOUR PRESCRIPTIONS:			

## 1) First Choice:

**ProCort** (Hydrocortisone Acetate 1.85%, Pramoxine HCL 1.15% with rectal applicators)

- Fully insert applicator before using, and clean immediately afterwards.
- Use 3-4 times daily or as directed by your physician.
- 14 day supply (60 g Tubes), 11 Refills

## 2) Second Choice:

**Analpram HC Cream 2.5% 1 OZ Tube** (Hydrocortisone Acetate 2.5%, Pramoxine HCL 1%)

- Dispense preferably with a rectal applicator
- Apply to affected area 3-4 times daily
- (Hydrocortisone Acetate 2.5%, Pramoxine HCL 1% Generic Available)
- 10 day supply (1 OZ Tubes), 11 Refills
- May dispense 90 day supply: At patient's option, dispense up to nine 1-oz tubes

## 3) Third Choice:

### Generic 2.5% Hydrocortisone cream 1 oz.

- Dispense preferably with a rectal applicator
- Apply to affected area 3-4 times daily
- 10 day supply (1 OZ Tubes), 11 Refills
- May dispense 90 day supply: At patient's option, dispense up to nine 1-oz tubes

#### 4) Fourth Choice:

# Hydrocortisone acetate 1% and pramoxine hydrochloride 1% aerosol foam

- Insert 1 applicatorful into the rectum and or apply with tissue around the anal area, 2 or 3 times daily and after each bowel evacuation.
- Dispense 4-canisters per month (i.e. 1- canisters per 7 days)
- 11 Refills of 4-canisters. When used correctly, the aerosol container will deliver a minimum of 14 applications)

Rick A. Shacket, DO, MD (H), BS9262611 Diplomate American Osteopathic Board of Proctology

#### **LOCATIONS**