

Rabbit Anti-CD19 [SP110]: RM0230

Intended Use: For Research Use Only

Description: CD19 is a pan B cell marker expressed in a wide range of maturational stages including pre B cells. CD19 labels the membrane of B cells in germinal centers including B cells and follicular dendritic cells, mantle zone cells and cells in the interfollicular areas. It is negative in plasma cells. CD19 is found in the majority of B cell-derived malignancies.

Specifications


Clone: SP110
 Source: Rabbit
 Isotype: IgG
 Localization: Membrane
 Formulation: Antibody in PBS pH7.4, containing BSA, and <= 0.09% sodium azide (NaN3)
 Storage: Store at 2°- 8°C
 Applications: IHC
 Package:

Description	Catalog No.	Size
CD19 Concentrated	RM0230	1 ml

IHC Procedure*

Positive Control Tissue: Lymph node, tonsil
 Concentrated Dilution: 50-100
 Pretreatment: Citrate pH6.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



Human diffused large B-cell lymphoma FFPE tissue stained with anti-CD19 using DAB

References:

1. Immunophenotyping of chronic B-cell neoplasms: flow cytometry versus immunohistochemistry. Abdel-Ghafar AA, et al. Hematol Rep. Jan 2;4(1):e3, 2012.
2. Composite T lymphoblastic leukemia/lymphoma and diffuse large B-cell lymphoma: case report. Niino D, et al. Pathol Int. Jun;61(6):363-8, 2011.
3. Potent antitumor activity of the anti-CD19 auristatin antibody drug conjugate hBU12-vcMMAE against rituximab-sensitive and -resistant lymphomas. Gerber HP, et al. Kung-Sutherland M, et al. Blood. Apr 30;113(18):4352-61, 2009.
4. CD19 regulates B cell maturation, proliferation, and positive selection in the FDC zone of murine splenic germinal centers. Wang Y, et al. Immunity. Jun;22(6):749-61, 2005.