

Informed Consent for Endoscopic Retrograde Cholangiopancreatogrophy (ERCP)

Name:	Procedure Date:	Time:
1. I,		ian) give consent for Dr.
ERCP with possible biopsy, dilation, s	her associates to perform an endoscopic retrophincterotomy (widening the sphincter), stone ssue, and control of bleeding if necessary.	
physician to visualize the interior of the Using a combination of endoscopic are therapy can be performed as deemed (sphincterotomy), followed by removato drain properly. Sedation and pain reprocedure. These medications may care	volves the passage of a digital optic instrume e esophagus, stomach, and duodenum (first sid x-ray techniques, visualization of the bile dinecessary, including widening the opening of of stones or placement of a stent or tube to a slieving medications may be given to minimize the localized irritation and/or a drug reaction is able to drive the remainder of the day and I HAVE A DRIVER take me home.	several inches of the small intestines). ucts is possible. Appropriate f the bile ducts or pancreatic ducts allow bile or pancreatic enzymes e discomfort and relax me for the . I understand that with the anesthesia/
	he procedure which have been adequately ex I regularly see my physician with any question uestions before signing this consent.	
perforation of the esophagus, stomac require surgery, hospitalization, repeat but rare complications which can occur can continue up to two weeks after the pancreas, caused by the procedure. It managed with pain medications for a but serious or possibly fatal risks incluprocedure is unsuccessful due to ana	s of this procedure include, but are not limited in, small intestines, or bile ducts. These complete ERCP, and/or a transfusion. Perforation of the at a rate of 1 per 1,000 endoscopies. Bleed the procedure. There is also a risk of infection at his occurs at a rate of 10 in 100 cases and called the development of the services of the services and called the services are the services of the service	ications, should they occur, may ne bowels or bile ducts are known, ing, usually after a sphincterotomy, and pancreatitis, or inflammation of the an range from mild abdominal pain, ich are very rare. Other extremely rare, ke. There is also a possibility that the ning and cannulating, or passing a
	guarantees regarding the results of this proced and may include radiologic imaging and pleir own limitations and benefits.	
have not been answered to my satisfa HAVE ANY QUESTIONS AS TO THE	nd this consent form, and understand that I s ction or if I do not understand any of the word RISKS OR HAZARDS OF THE PROPOSED IGNING THIS CONSENT FORM. DO NOT S FORM.	ds or terms used in this form. IF YOU PROCEDURE OR TREATMENT, ASK
Patient/Legal Representative signatur	e Date	Time
Witness signature	Date	Time