

June 4, 2015



New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration
Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

PROLINE® 480 SC Fungicide

EPA Reg. No. 264-825

This supplemental label expires on 01/16/2018 and must not be used or distributed after this date.

Supplemental Label

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for PROLINE® 480 SC Fungicide attached to the container.

USE DIRECTIONS FOR SPECIFIC CROPS

APPLICATION DIRECTIONS			
CROP	DISEASE CONTROLLED	RATE OF PROLINE 480 SC FUNGICIDE	APPLICATION TIMING
Corn (field corn, field corn grown for seed and popcorn)	Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye spot (<i>Aureobasidium zeae</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>) Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>)* Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>)* Rust (<i>Puccinia spp.</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)* *The above diseases are also known as Helminthosporium leaf blights	5.7 fl oz per acre	Apply PROLINE 480 SC at the first sign of disease. Repeat applications as needed on a 7 – to 14-day interval if favorable conditions for disease development persist. Do not use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel is visible but the silks have not yet emerged).
	For the suppression of Fusarium, Gibberella and Aspergillus ear rots (<i>Fusarium spp.</i> , <i>Gibberella spp.</i> and <i>Aspergillus spp.</i>)	5.7 fl oz per acre	For optimum suppression of Fusarium, Gibberella and Aspergillus ear rots, apply PROLINE 480 SC Fungicide from the R1 (initial silk emergence) to the R2 (brown silk) corn growth stages. PROLINE 480 SC Fungicide will reduce both disease symptoms and levels of mycotoxin in the grain.

PROLINE is a Registered trademark of Bayer.

SAL 02/10/2015 – PRLV2

Other requirements: PROLINE 480 SC Fungicide may be applied by either ground, aerial or chemigation application equipment.

For aerial applications, apply PROLINE 480 SC Fungicide using a minimum of 3 gpa spray solution. An adjuvant may be used to improve spray coverage. Refer to the adjuvant product label for specific use directions.

Application of PROLINE 480 SC Fungicide is not recommended at times when corn is under severe environmental stress conditions. Do not exceed 22.8 fl oz total from all uses, including soil and foliar applications, per acre of PROLINE 480 SC Fungicide per year. Do not apply within 14 days of harvest for grain and fodder. Forage may be harvested the same day of application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).