

Patient Name: _____ DOB _____ Age _____ Date _____

PERI-OPERATIVE CONCIOUS SEDATION

☐ Medical Assistant – Certified observing: _____ ☐ Time _____ Minutes

☐ Pulse Oximetry Average PaO2 _____ % | ☐ Oxygen _____ % | Nitrogen _____ % | Flow _____ Liter/Min

15 Minute Vitals: BP: _____ / _____ P: _____ RR: _____ | 30 Minute Vitals: BP: _____ / _____ P: _____ RR: _____

45 Minute Vitals: BP: _____ / _____ P: _____ RR: _____ | _____ Minute Vitals: BP: _____ / _____ P: _____ RR: _____

DIAGNOSES: ☐ R10.2 AnoRectal Pain ☐ F40.9 Phobia

POST-PROCEDURE

PUDENDAL NERVE BLOCK FOR POST-OPERATIVE ANALGESIA

☐ Medical Assistant – Certified observing: _____

☐ The purpose and indication for the regional anesthesia of the pudendal nerve, is for postoperative analgesia.

☐ Modifier -59 is required to distinguish the nerve block from the intraoperative anesthesia.

☐ This regional block is as separate from routine postoperative pain management, which consists merely of daily PO narcotic analgesia.

☐ The time spent on perioperative placement of the block is separate from, and was not included in any reported anesthesia time.

☐ PROCEDURE NOTE: Using a transcutaneous perineal approach, the ischial tuberosity is palpated and a 25 gage needle was introduced slightly medial to the tuberosity. The needle is advanced approximately 2.5 cm. Aspiration was performed to ensure that the needle is not in a blood vessel and then the anesthetic is injected. The needle is withdrawn and directed into the deep superficial tissue of the anus and anesthetic is again injected to block the perineal, inferior rectal, ilioinguinal, and genitofemoral components of the pudendal nerve. → ☐ This was repeated on the opposite side.

☐ During the injection procedure, this patient was monitored with pulse oximetry, pulse rate, and blood pressure, by other qualified trained observer(s) in the operating room. The motor response during the injection was negligible, unless otherwise stated below:

☐ Drug Used:

☐ Bupivacaine & Adrenaline Injection B.P. 0.25% w/ 1 in 200,000, ☐ 5ml ☐ 10ml ☐ 15ml ☐ 20ml

☐ Exparel 1 Unit, 266mg, 20ml, (C9290)

☐ Amount of drug not administered, wasted:

☐ Post Procedure Vital Signs: BP: _____ P: _____ RR: _____ PaO2: _____ %

DIAGNOSES: ☐ G89.18 Acute Post Procedural Pain ☐ G89.28 Chronic Post Procedural Pain

8752 E Via De Commercio #2, Scottsdale, Arizona 85258, drshacket@yahoo.com, 602.492.9919

Diplomate American Osteopathic Board of Proctology, Rick Shacket, DO, MD(H) _____