

**Mouse Anti-PAX5 [A11]: MC0203, MC0203RTU7**

**Intended Use:** For Research Use Only

**Description:** PAX5 is a B-cell lineage specific activator protein (BSAP) that is essential for maintaining the identity and function of mature B cells during late B lymphopoiesis. It also plays a role in neural development and spermatogenesis. PAX5 is expressed in pro-, pre-, and mature B cells, and it is expressed in the vast majority of B-cell malignancies. Anti-PAX5 is a specific marker for the B cell lineage. PAX5 is thus useful for a panel of antibodies for the identification of cellular origin of undifferentiated tumors. The expression of PAX5 in endocrine tumors has been shown to be high in Merkel cell carcinoma and small cell carcinoma, but not carcinoid tumor. PAX5 is also a marker for neuronendocrine carcinomas.

**Specifications**


Clone: A11  
 Source: Mouse  
 Isotype: IgG2a/λ  
 Reactivity: Human, mouse, rat  
 Localization: Nucleus  
 Formulation: Antibody in PBS pH7.4, containing BSA ≤ 0.09% sodium azide (NaN3)  
 Storage: Store at 2°- 8°C  
 Applications: IHC, ELISA, IF, IP, WB  
 Package:

Description	Catalog No.	Size
PAX5 Concentrated	MC0203	1 ml
PAX5 Prediluted	MC0203RTU7	7 ml

**IHC Procedure\***

Positive Control Tissue: Tonsil  
 Concentrated Dilution: 50-200  
 Pretreatment: Citrate pH6.0 or EDTA pH8.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C  
 Incubation Time and Temp: 30-60 minutes @ RT  
 Detection: Refer to the detection system manual

\* Result should be confirmed by an established diagnostic procedure.



FFPE human tonsil stained with anti-PAX5 using DAB

**References:**

1. Diagnostic utility of PAX2 and PAX5 in distinguishing non-small cell lung cancer from small cell lung cancer. Ren Y, et al. Int J Clin Exp Pathol 8:14709-16, 2015.
2. Protein Expression for Novel Prognostic Markers (Cyclins D1, D2, D3, B1, B2, ITGβ7, FGFR3, PAX5) Correlate With Previously Reported Gene Expression Profile Patterns in Plasma Cell Myeloma. Mansoor A, et al. Appl Immunohistochem Mol Morphol. May-Jun;23(5):327-33, 2015.
3. A recurrent germline PAX5 mutation confers susceptibility to pre-B cell acute lymphoblastic leukemia. Shah S, et al. Nat Genet 45:1226-31, 2013.

Doc. 100-MC0203  
Rev. A