

Kittitas County Prehospital EMS Protocols

SUBJECT: PEDIATRIC CARDIOPULMONARY ARREST

- A. Establish and maintain airway.
- B. Verify cardiopulmonary arrest.
- C. Initiate CPR and ventilate per pediatric BVM with supplemental O₂ @ 12-15 lpm.
- D. Place ET tube and continue ventilations with BVM.
- E. Establish cardiac monitor.
- F. Establish peripheral IV access with **Isotonic Crystalloid @ TKO.**
- G. Consider intraosseous route if patient < 12 years old, or external jugular.

Asystole / Pulseless Electrical Activity (PEA)

- A. Attempt to identify underlying cause of PEA, and treat as per protocols.
- B. Administer **Epinephrine**, 1:10,000, 0.01 mg/kg q 3 minutes, IV or IO.
- C. If only ET route available, administer **Epinephrine**, 1:1,000, 0.1 mg/kg q 3 minutes.
- D. In PEA, consider fluid challenge of **Isotonic Crystalloid @ 20 cc/kg.**

Ventricular fibrillation (Pulseless VT)

- A. Defibrillate ASAP @ 2 joules/kg. Second and subsequent shocks shall be at 4 joules/kg.
- B. Administer **Epinephrine**, 1:10,000, 0.01 mg/kg q 3 minutes, IV or IO.
- C. If only ET route available, administer **Epinephrine**, 1:1,000, 0.1 mg/kg q 3 minutes.
- D. Immediately following first dose of **Epinephrine**, administer **Amiodarone**, 5 mg/kg, IV or IO.
- E. Repeat defibrillation @ 4 joules/kg as necessary.