

**Rabbit Anti-BOB.1 [EPR17685]: RM0010, RM0010RTU7**

**Intended Use:** For Research Use Only

**Description:** BOB.1, also known as B-cell-specific coactivator OBF-1 or OCA-B, is a lymphoid-specific transcriptional coactivator that interacts with the transcription factors Oct-1 and Oct-2. BOB.1 has been shown to be critical for the development of a normal immune response, where it mediates octamer-dependent transcriptional activity in B lymphocytes. It is also critically involved in the formation of germinal centers in secondary lymphoid organs. BOB.1 levels have been observed to be massively upregulated in germinal center B cells, as compared with resting B cells. The BOB.1 antibody labels B lymphocytes and plasma cells. It is expressed in various B cell derived lymphomas and Hodgkin's lymphomas (HL). The expression of BOB.1 is high in Nodular lymphocyte predominant Hodgkin lymphoma (NLPHL), but low in classic HL.

**Specifications:**


Clone: EPR17685  
 Source: Rabbit  
 Isotype: IgG  
 Reactivity: Human  
 Immunogen: Recombinant fragment within human BOB1 aa 1 to the C-terminus  
 Localization: Nucleus  
 Formulation: Antibody in PBS pH7.4, containing BSA and ≤ 0.09% sodium azide (NaN3)  
 Storage: Store at 2°- 8°C  
 Applications: IHC, WB  
 Package:

| Description        | Catalog No. | Size |
|--------------------|-------------|------|
| BOB.1 Concentrated | RM0010      | 1 ml |
| BOB.1 Prediluted   | RM0010RTU7  | 7 ml |

**IHC Procedure\*:**

Positive Control Tissue: Tonsil, lymphoma  
 Concentrated Dilution: 25-50  
 Pretreatment: Tris EDTA pH9.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C  
 Incubation Time and Temp: 30-60 minutes @ RT  
 Detection: Refer to the detection system manual

\* Result should be confirmed by an established diagnostic procedure.



FFPE human non-hodgkin lymphoma stained with anti-BOB.1 using DAB

**References:**

1. Analysis of Transcription Factor OCT.1, OCT.2 and BOB.1 Expression Using Tissue Arrays in Classical Hodgkin's Lymphoma. Mónica García-Cosío, et al. Mod Pathol, 17 (12), 1531-8 Dec 2004.
2. Differential Emu Enhancer Activity and Expression of BOB.1/OBF.1, Oct2, PU.1, and Immunoglobulin in Reactive B-cell Populations, B-cell non-Hodgkin Lymphomas, and Hodgkin Lymphomas. Christoph Loddenkemper, et al. J Pathol, 202 (1), 60-9 Jan 2004.
3. The B Lymphocyte-Specific Coactivator BOB.1/OBF.1 Is Required at Multiple Stages of B-cell Development. J Hess, et al. Mol Cell Biol, 21 (5), 1531-9 Mar 2001.

Doc. 100-RM0010  
Rev. B