

Rabbit Anti-HIF-1 alpha [EP118]: RM0374

Intended Use: For Research Use Only

Description: The HIF-1 Alpha subunit of hypoxia-inducible factor 1 is a transcription factor that functions as a master transcriptional regulator of the adaptive response to hypoxia. HIF-1 activates the transcription of many genes, thus playing a role in various biological processes, including cardiovascular development, angiogenesis, energy metabolism, and cell survival.


Specifications

Clone: EP118
Source: Rabbit
Isotype: IgG
Reactivity: Human
Localization: Nucleus, cytoplasm
Formulation: Antibody in PBS pH7.5, containing 0.2% BSA and <0.1% sodium azide (NaN₃)
Storage: Store at 2°- 8°C
Applications: IHC
Package:

Description	Catalog No.	Size
HIF-1 alpha Concentrated	RM0374	1 ml

IHC Procedure

Positive Control: Breast carcinoma
Concentrated Dilution: 50-200
Pretreatment: Citrate pH6.0 or EDTA pH8.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
Incubation Time and Temp: 30-60 minutes @ RT
Detection: Refer to the detection system manual
* Result should be confirmed by an established diagnostic procedure.



FFPE human breast carcinoma tissue stained with anti-HIF-1 alpha using DAB

References:

1. Hypoxia-Inducible Factor-1 α Is Associated With Sprouting Angiogenesis in the Murine Laser-Induced Choroidal Neovascularization Model. André H, et al. Invest Ophthalmol Vis Sci. Oct;56(11):6591-604, 2015.
2. CXCR7 and CXCR4 Expressions in Infiltrative Astrocytomas and Their Interactions with HIF1 α Expression and IDH1 Mutation. Bianco AM, et al. Pathol Oncol Res. Apr;21(2):229-40, 2015.
3. HIF-1 α , VEGF and WT-1 are protagonists in bilateral primary angiosarcoma of breast: a case report and review of literature. Al-Salam S, et al. Int J Clin Exp Pathol. 5(3):247-53, 2012.
4. Hypoxia and the expression of HIF-1alpha and HIF-2alpha in the retina of streptozotocin-injected mice and rats. Wright WS, et al. Exp Eye Res. Mar;90(3):405-12, 2010.