



Pharmacy

Prepared for: Chris Tester

Prescriber: Norman Medman, MD

DOB: 1/10/1957

Pharmacist: Bob Farmer

BETROPROLEN 100 MG TAB Refills: NO

Date: 11/22/2018

**TAKE ONE TABLET BY MOUTH
TWICE DAILY IN THE MORNING
AND EVENING**

Rx #: 5900476

BETROPROLEN (Be-TROE-PROE-LEN)

General Uses: Betroprolen is a beta-blocker used primarily to treat high blood pressure and angina. It may be used after a heart attack to stabilize heart functioning. Your doctor may also prescribe this medicine for other uses.

Directions for Use: Be sure to follow your doctor's directions for use of this medication. The dosage and schedule are described above. Take the tablets at the same time each day, with or without food as prescribed. Store this medicine in a cool, dark place. If you miss a dose of this medication take it right away unless it is nearly time for your next dose. If it is close to the time for your next dose, skip the missed dose and resume your normal schedule. **DO NOT DOUBLE THE DOSE UNLESS YOUR DOCTOR TELLS YOU TO DO SO.**

Precautions: Always take the exact dose at the prescribed time. Check with your doctor before changing your schedule or discontinuing the use of this medicine. Before you have any medical procedures or surgery (including dental care), tell your medical provider about the use of this medicine. This medication may cause drowsiness or vertigo, especially initially. Do not drive or operate dangerous machinery until you know how you tend to react to this medication. Do not combine this medicine with any other medicine including over-the-counter medication. Inform your doctor if you are pregnant or if you plan to become pregnant in the next six months. This medicine is transferred to breast milk, so do not use this medication without checking with your doctor, if you are breast feeding. Occasionally this medication causes hot flashes, loose bowels, and rapid heart rate.

Side effects: Side effects of this medication include drowsiness, dizziness, fatigue, slight weakness, and decreased endurance. If any of these persist for more than one week or are bothersome to you, call your pharmacist or health care provider. If you experience severe shortness of breath, call your local emergency number immediately.

Other adverse effects may rarely occur. Any questions or comments should be directed to your health care provider.



Pharmacy Prepared for: Chris Tester

Prescriber: Norman Medman, MD

DOB: 1/10/1957

Pharmacist: Bob Farmer

CYCLONOPROL 500 MG TAB Refills: NO

Date: 11/22/2018

**TAKE ONE TABLET BY MOUTH
THREE TIMES DAILY WITH FOOD**

Rx #: 5900477

CYCLONOPROL (Sye-cloe-NOE-prol)

General Uses: Cyclonoprol is a narcotic and acetaminophen combination. It is generally used to combat moderate to severe pain. It works by binding to the opioid receptors in the central nervous system and decreases the formation of cortisol. Your doctor may also prescribe this medicine for other uses.

Directions for Use: Be sure to follow your doctor's directions for use of this medication. The dosage and schedule are described above. Take the tablets by mouth at the same time each day, with food as prescribed. If taking it with food upsets your stomach contact the prescriber. Store this medicine in a cool, dark place in a tightly closed container. Keep the tablets out of sunlight. If you miss a dose of this medication take it right away unless it is nearly time for your next dose. If it is close to the time for your next dose, skip the missed dose and resume your normal schedule. **DO NOT DOUBLE THE DOSE UNLESS YOUR DOCTOR TELLS YOU TO DO SO.**

Precautions: Always take the exact dose at the prescribed time. If you are allergic to codeine or hydrocodone do not take this medication. Check with your doctor before changing your schedule or discontinuing the use of this medicine. Do not exceed the recommended dose. Before you have any medical procedures or surgery (including dental care), tell your medical provider about the use of this medicine. This medication may cause drowsiness or vertigo, especially initially. To avoid dizziness, be sure to get up slowly and allow yourself time to adjust to position changes. Do not drive or operate dangerous machinery until you know how you react to this medication. Do not combine this medicine with any other medicine including over-the-counter medication. Do not consume alcohol in combination with this medication. Inform your doctor if you are pregnant or if you plan to become pregnant in the next six months. This medicine is transferred to breast milk, so do not use this medication without checking with your doctor, if you are breast feeding. Occasionally this medication causes hot flashes and rapid heart rate. Elderly people may be more sensitive to the effects of this medication.

Side effects: Side effects of this medication include mild nausea, vomiting, drowsiness, dizziness, mild anxiety, and jitteriness. Visual changes should be reported to your doctor. If any of these persist for more than one week or are bothersome to you, call your pharmacist or health care provider. If you experience severe shortness of breath, significant mood change, or loss of hearing, call your local emergency number immediately. Other adverse effects may rarely occur. Any questions or comments should be directed to your health care provider.



Pharmacy Prepared for: Chris Tester
DOB: 1/10/1957
Refills: NO
Rx #: 5900477

Prescriber: Norman Medman, MD
Pharmacist: Bob Farmer
Date: 11/22/2018

ANDROCIPLIN
TAKE ONE CAPSULE BY MOUTH
TWICE DAILY

ANDROCIPLIN (An-DROE-si-plin)

General Uses: Androciplin is a penicillin based antibiotic. It is generally used to combat bacterial infections.

Directions for Use: Be sure to follow your doctor's directions for use of this medication. The dosage and schedule are described above. Take the tablets by mouth at the same time each day, either one hour before meals or two hours after eating on an empty stomach. If taking it with food upsets your stomach contact the prescriber. Store this medicine in a tightly closed container at room temperature below 80 degrees Fahrenheit. Keep the tablets out of bright sunlight. If you miss a dose of this medication, take it right away unless it is nearly time for your next dose. If it is close to the time for your next dose, skip the missed dose and resume your normal schedule. **DO NOT DOUBLE THE DOSE UNLESS YOUR DOCTOR TELLS YOU TO DO SO. BE SURE TO CONTINUE TAKING THIS MEDICATION UNTIL YOU HAVE TAKEN ALL THE PRESCRIBED PILLS, EVEN IF YOU FEEL BETTER IN A FEW DAYS.**

Precautions: Always take the exact dose at the prescribed time. If you are allergic to Penicillin, do not take this medication. Check with your doctor before changing your schedule or discontinuing the use of this medicine. Take all the pills in the container as prescribed. Do not exceed the recommended dose. Before you have any medical procedures or surgery (including dental care), tell your medical provider about the use of this medicine. Do not combine this medicine with any other medicine including over-the-counter medication. Do not consume alcohol in combination with this medication. Inform your doctor if you are pregnant or if you plan to become pregnant in the next six months. **THIS MEDICATION MAY DECREASE THE EFFECTIVENESS OF BIRTH CONTROL PILLS.** This medicine is transferred to breast milk, so do not use this medication without checking with your doctor, if you are breast feeding.

Side effects: Side effects of this medication include mild nausea, vomiting, diarrhea, mouth and lip irritations. If any of these persist for more than one or two days or are bothersome to you, call your pharmacist or health care provider. If you experience severe shortness of breath, a rash or hives, or vaginal discharge, call your doctor as soon as you can. Other adverse effects and drug interactions may rarely occur. Contact your health care provider with questions or comments, or if you would like more information.