

Rabbit Anti-CD38 Recombinant [MD71R]: RM0459, RM0459RTU7

Intended Use: For Research Use Only

Description: CD38, also called ADP-ribosyl cyclase, is a druggable ectoenzyme that uses nicotinamide adenine dinucleotide (NAD) as a substrate to generate second messengers. In particular, it synthesizes cyclic ADP-ribose, a second messenger for glucose-induced insulin secretion. CD38 also has cADPR hydrolase activity. It is preferentially expressed at both early and late stages of B- and T-cell maturation. In normal lymph nodes and tonsils, the antigen is detected mainly on B cells in germinal centers and plasma cells. An antibody to CD38 is helpful in the identification of plasma cells and tumors with plasmablastic differentiation. A prognostic value of CD38 in B-cell chronic lymphocytic leukemia (CLL) has been reported. Expression of CD38 is linked to unmutated IgVH genes and a worse prognosis. CD38 is involved in the generation of adenosine, which is implicated in tumor immune evasion. CD38 mRNA expression in metastatic castration-resistant prostate cancer (mCRPC) was most significantly associated with upregulated immune signaling pathways, and with interleukin (IL)-12, IL-23, and IL-2.

Specifications:


Clone: MD71R
 Source: Rabbit
 Isotype: IgG
 Reactivity: Human
 Immunogen: Synthetic peptide to CD38 aa 200-300
 Localization: Membrane
 Formulation: Antibody in PBS pH7.4, containing BSA and $\leq 0.09\%$ sodium azide (NaN₃)
 Storage: Store at 2°- 8°C
 Applications: IHC
 Package:

Description	Catalog No.	Size
CD38 Recombinant Concentrated	RM0459	1 ml
CD38 Recombinant Prediluted	RM0459RTU7	7 ml

IHC Procedure*:

Positive Control Tissue: Tonsil, spleen. Raji cells
 Concentrated Dilution: 50-200
 Pretreatment: Tris EDTA pH9.0, 15 minutes using Pressure Cooker, or 30-60 minutes using water bath at 95°-99°C
 Incubation Time and Temp: 30-60 minutes @ RT
 Detection: Refer to the detection system manual

* Result should be confirmed by an established diagnostic procedure.



FFPE human tonsil stained with anti-CD38 Recombinant using DAB

References:

1. Multi-cellular 3D human primary liver cell culture elevates metabolic activity under fluidic flow. Esch MB, et al. Lab Chip 15:2269-77, 2015.
2. Clinicopathological study of primary biliary cirrhosis with interface hepatitis compared to autoimmune hepatitis. World Kobayashi M, et al. J Gastroenterol 20:3597-608, 2014.

Doc. 100-RM0459
Rev. A